



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 06:50 AM UTC

PDB ID : 6JML / pdb_00006jml
Title : Re-refined structure of R-state L-lactate dehydrogenase from *Lactobacillus casei*
Authors : Arai, K.; Miyanaga, A.; Uchikoba, H.; Fushinobu, S.; Taguchi, H.
Deposited on : 2019-03-12
Resolution : 2.30 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	5% 86% 7% • 5%
1	B	326	3% 88% 8% • •
1	C	326	2% 82% 8% • 9%
1	D	326	9% 84% 7% • 7%
1	E	326	4% 87% 8% • •

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	326	2% 87% 8% . .

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 14535 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-lactate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	309	Total	C	N	O	S	0	0	0
			2371	1515	393	457	6			
1	B	315	Total	C	N	O	S	0	0	0
			2414	1541	401	466	6			
1	C	296	Total	C	N	O	S	0	0	0
			2277	1456	377	438	6			
1	D	304	Total	C	N	O	S	0	0	0
			2327	1484	387	450	6			
1	E	314	Total	C	N	O	S	0	0	0
			2409	1538	400	465	6			
1	F	315	Total	C	N	O	S	0	0	0
			2414	1541	401	466	6			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0
2	F	1	Total O S 5 4 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	44	Total O 44 44	0	0
3	B	54	Total O 54 54	0	0
3	C	39	Total O 39 39	0	0
3	D	36	Total O 36 36	0	0
3	E	47	Total O 47 47	0	0
3	F	43	Total O 43 43	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-lactate dehydrogenase




- Molecule 1: L-lactate dehydrogenase




- Molecule 1: L-lactate dehydrogenase





- Molecule 1: L-lactate dehydrogenase







• Molecule 1: L-lactate dehydrogenase



• Molecule 1: L-lactate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	164.38Å 82.89Å 179.77Å 90.00° 91.21° 90.00°	Depositor
Resolution (Å)	34.10 – 2.30 34.10 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (34.10-2.30) 99.9 (34.10-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, R_{free}	0.194 , 0.222 0.201 , 0.227	Depositor DCC
R_{free} test set	5411 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	40.2	Xtriage
Anisotropy	0.502	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.008 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14535	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	2/2413 (0.1%)	1.42	6/3273 (0.2%)
1	B	1.21	6/2458 (0.2%)	1.46	8/3335 (0.2%)
1	C	1.15	1/2317 (0.0%)	1.44	6/3141 (0.2%)
1	D	1.15	1/2366 (0.0%)	1.42	6/3208 (0.2%)
1	E	1.17	1/2453 (0.0%)	1.44	5/3328 (0.2%)
1	F	1.20	7/2458 (0.3%)	1.41	6/3335 (0.2%)
All	All	1.17	18/14465 (0.1%)	1.43	37/19620 (0.2%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	124	ALA	C-O	-8.09	1.13	1.24
1	B	181	HIS	CE1-NE2	7.54	1.40	1.32
1	C	187	PRO	C-O	-6.72	1.16	1.23
1	F	187	PRO	C-O	-6.54	1.17	1.23
1	F	59	PRO	C-O	-6.28	1.15	1.24
1	F	191	HIS	CE1-NE2	6.21	1.38	1.32
1	B	63	PRO	C-O	-6.16	1.16	1.23
1	B	84	ALA	N-CA	6.12	1.55	1.46
1	A	84	ALA	C-O	5.92	1.31	1.24
1	B	83	THR	C-O	5.83	1.31	1.24
1	D	59	PRO	C-O	-5.82	1.16	1.24
1	F	8	ASP	C-O	5.58	1.30	1.23
1	E	63	PRO	C-O	-5.45	1.17	1.23
1	B	174	HIS	CE1-NE2	5.43	1.38	1.32
1	F	192	ALA	C-O	5.32	1.30	1.23
1	F	259	PRO	C-O	-5.31	1.18	1.23
1	A	59	PRO	C-O	-5.25	1.17	1.24
1	F	84	ALA	C-O	5.09	1.30	1.24

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	83	THR	CA-C-N	-12.01	105.48	122.41
1	B	83	THR	C-N-CA	-12.01	105.48	122.41
1	A	166	VAL	N-CA-CB	-10.63	100.51	112.21
1	C	166	VAL	N-CA-CB	-10.26	100.92	112.21
1	D	166	VAL	N-CA-CB	-9.84	101.38	112.21
1	F	166	VAL	N-CA-CB	-9.38	101.89	112.21
1	B	166	VAL	N-CA-CB	-8.25	101.62	112.26
1	E	166	VAL	N-CA-CB	-8.03	101.90	112.26
1	B	288	ASN	CA-CB-CG	7.45	120.05	112.60
1	B	126	ASN	CA-CB-CG	-6.76	105.83	112.60
1	D	183	ASP	CB-CA-C	-6.25	98.34	110.46
1	C	83	THR	CA-C-O	-6.06	110.50	120.80
1	A	212	ASP	CB-CA-C	6.05	122.30	110.67
1	D	182	GLY	CA-C-O	-5.91	116.61	121.76
1	F	126	ASN	CA-CB-CG	-5.90	106.70	112.60
1	C	126	ASN	CB-CA-C	5.77	117.01	108.87
1	A	126	ASN	CA-CB-CG	-5.68	106.92	112.60
1	D	248	LYS	N-CA-C	-5.67	105.18	111.36
1	E	126	ASN	CB-CA-C	5.66	116.84	108.87
1	E	183	ASP	CB-CA-C	-5.66	99.49	110.46
1	D	166	VAL	CB-CA-C	5.64	117.89	110.84
1	F	166	VAL	CB-CA-C	5.60	117.84	110.84
1	A	103	LYS	CB-CG-CD	5.51	123.96	111.30
1	A	213	LYS	CB-CG-CD	5.46	123.86	111.30
1	F	137	LYS	CA-CB-CG	5.46	125.02	114.10
1	A	182	GLY	CA-C-O	-5.33	117.12	121.76
1	C	222	ARG	NE-CZ-NH1	-5.33	116.17	121.50
1	B	83	THR	CA-C-O	-5.28	112.91	118.77
1	C	166	VAL	CB-CA-C	5.27	117.43	110.84
1	B	83	THR	O-C-N	-5.23	116.59	122.34
1	E	236	PHE	CA-CB-CG	5.23	119.03	113.80
1	B	126	ASN	CB-CA-C	5.19	116.19	108.87
1	F	84	ALA	CA-C-O	-5.14	113.15	120.51
1	D	126	ASN	CB-CA-C	5.12	116.08	108.87
1	E	48	LYS	CB-CA-C	5.11	119.54	110.85
1	F	317	PHE	CB-CA-C	5.10	119.18	110.01
1	C	304	LYS	CB-CG-CD	5.06	122.93	111.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2371	0	2392	9	0
1	B	2414	0	2435	7	0
1	C	2277	0	2301	9	0
1	D	2327	0	2345	13	0
1	E	2409	0	2430	13	0
1	F	2414	0	2435	10	0
2	A	10	0	0	0	0
2	B	10	0	0	0	0
2	C	10	0	0	0	0
2	D	10	0	0	0	0
2	E	10	0	0	0	0
2	F	10	0	0	0	0
3	A	44	0	0	2	0
3	B	54	0	0	0	0
3	C	39	0	0	0	0
3	D	36	0	0	1	0
3	E	47	0	0	0	0
3	F	43	0	0	2	0
All	All	14535	0	14338	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:94:ARG:CG	1:D:94:ARG:HH21	2.03	0.71
1:A:235:THR:HG23	3:A:502:HOH:O	1.88	0.71
1:D:200:ALA:O	1:D:203:VAL:HG12	1.93	0.68
1:F:137:LYS:HD2	3:F:525:HOH:O	1.91	0.68
1:C:161:SER:OG	1:C:217:MET:HG3	1.95	0.67
1:D:42:ILE:HG13	3:D:523:HOH:O	1.97	0.65
1:D:94:ARG:HH21	1:D:94:ARG:HG2	1.63	0.63
1:C:4:ILE:O	1:C:7:LYS:HG3	2.01	0.61
1:C:231:LEU:HD13	1:E:48:LYS:HG3	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:4:ILE:HG12	1:D:7:LYS:HD3	1.89	0.53
1:A:161:SER:OG	1:A:217:MET:HG3	2.09	0.52
1:B:263:TYR:CZ	1:B:272:ASP:HA	2.47	0.50
1:E:263:TYR:CZ	1:E:272:ASP:HA	2.47	0.50
1:F:161:SER:OG	1:F:217:MET:HG3	2.10	0.49
1:C:124:ALA:HB2	1:C:243:LEU:HD21	1.95	0.49
1:D:263:TYR:CZ	1:D:272:ASP:HA	2.48	0.49
1:F:263:TYR:CZ	1:F:272:ASP:HA	2.48	0.49
1:A:263:TYR:CZ	1:A:272:ASP:HA	2.48	0.49
1:C:263:TYR:CZ	1:C:272:ASP:HA	2.49	0.47
1:B:137:LYS:HD2	1:B:137:LYS:HA	1.71	0.46
1:F:58:LEU:N	1:F:59:PRO:CD	2.79	0.46
1:E:58:LEU:N	1:E:59:PRO:CD	2.79	0.46
1:D:293:PRO:HG2	1:E:201:GLU:CG	2.47	0.45
1:E:4:ILE:HG12	1:E:7:LYS:HD3	1.97	0.45
1:A:58:LEU:N	1:A:59:PRO:CD	2.80	0.45
1:C:58:LEU:N	1:C:59:PRO:CD	2.80	0.44
1:D:58:LEU:N	1:D:59:PRO:CD	2.80	0.44
1:B:163:ALA:HB1	1:B:168:VAL:O	2.17	0.44
1:E:163:ALA:HB1	1:E:168:VAL:O	2.17	0.44
1:D:253:ASP:OD1	1:D:282:ASN:HB2	2.18	0.44
1:D:110:ASP:HB2	1:D:111:PRO:HD3	2.00	0.43
1:F:110:ASP:HB2	1:F:111:PRO:HD3	2.00	0.43
1:C:163:ALA:HB1	1:C:168:VAL:O	2.19	0.43
1:E:137:LYS:HD2	1:E:137:LYS:HA	1.88	0.43
1:B:58:LEU:N	1:B:59:PRO:CD	2.80	0.43
1:C:253:ASP:OD1	1:C:282:ASN:HB2	2.19	0.43
1:D:293:PRO:HG2	1:E:201:GLU:CD	2.44	0.43
1:B:253:ASP:OD1	1:B:282:ASN:HB2	2.19	0.42
1:A:163:ALA:HB1	1:A:168:VAL:O	2.19	0.42
1:B:110:ASP:HB2	1:B:111:PRO:HD3	2.02	0.42
1:F:137:LYS:CD	3:F:525:HOH:O	2.62	0.42
1:A:110:ASP:HB2	1:A:111:PRO:HD3	2.02	0.42
1:E:110:ASP:HB2	1:E:111:PRO:HD3	2.02	0.42
1:F:163:ALA:HB1	1:F:168:VAL:O	2.19	0.42
1:B:270:LEU:HD22	1:B:306:ALA:CB	2.50	0.41
1:C:228:ILE:HB	1:C:235:THR:HG22	2.01	0.41
1:F:106:LYS:HD2	1:F:138:LEU:HD22	2.02	0.41
1:E:124:ALA:HB2	1:E:243:LEU:HD21	2.02	0.41
1:E:253:ASP:OD1	1:E:282:ASN:HB2	2.21	0.41
1:A:253:ASP:OD1	1:A:282:ASN:HB2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:201:GLU:CD	1:E:293:PRO:HG2	2.45	0.41
1:A:137:LYS:HD2	1:A:137:LYS:HA	1.90	0.40
1:D:196:GLY:O	1:E:191:HIS:HD2	2.05	0.40
1:F:253:ASP:OD1	1:F:282:ASN:HB2	2.20	0.40
1:F:124:ALA:HB2	1:F:243:LEU:HD21	2.03	0.40
1:A:42:ILE:HG13	3:A:538:HOH:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	305/326 (94%)	296 (97%)	9 (3%)	0	100	100
1	B	313/326 (96%)	301 (96%)	12 (4%)	0	100	100
1	C	292/326 (90%)	282 (97%)	10 (3%)	0	100	100
1	D	298/326 (91%)	288 (97%)	9 (3%)	1 (0%)	36	46
1	E	312/326 (96%)	303 (97%)	9 (3%)	0	100	100
1	F	313/326 (96%)	303 (97%)	10 (3%)	0	100	100
All	All	1833/1956 (94%)	1773 (97%)	59 (3%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	204	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/269 (95%)	245 (96%)	10 (4%)	28	43
1	B	259/269 (96%)	250 (96%)	9 (4%)	32	48
1	C	246/269 (91%)	239 (97%)	7 (3%)	38	56
1	D	250/269 (93%)	240 (96%)	10 (4%)	28	42
1	E	259/269 (96%)	252 (97%)	7 (3%)	39	58
1	F	259/269 (96%)	250 (96%)	9 (4%)	32	48
All	All	1528/1614 (95%)	1476 (97%)	52 (3%)	32	49

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ILE
1	A	137	LYS
1	A	166	VAL
1	A	204	LYS
1	A	208	GLU
1	A	213	LYS
1	A	235	THR
1	A	265	ASP
1	A	307	SER
1	A	308	GLN
1	B	4	ILE
1	B	48	LYS
1	B	137	LYS
1	B	166	VAL
1	B	204	LYS
1	B	208	GLU
1	B	216	LYS
1	B	235	THR
1	B	308	GLN
1	C	4	ILE
1	C	137	LYS
1	C	166	VAL
1	C	208	GLU
1	C	265	ASP
1	C	307	SER
1	C	308	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	42	ILE
1	D	94	ARG
1	D	137	LYS
1	D	166	VAL
1	D	204	LYS
1	D	213	LYS
1	D	265	ASP
1	D	300	GLU
1	D	307	SER
1	D	308	GLN
1	E	137	LYS
1	E	166	VAL
1	E	204	LYS
1	E	208	GLU
1	E	216	LYS
1	E	265	ASP
1	E	308	GLN
1	F	106	LYS
1	F	126	ASN
1	F	166	VAL
1	F	204	LYS
1	F	208	GLU
1	F	211	GLU
1	F	265	ASP
1	F	307	SER
1	F	308	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	284	ASN
1	E	191	HIS
1	F	284	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	D	401	-	4,4,4	0.37	0	6,6,6	0.49	0
2	SO4	C	402	-	4,4,4	0.42	0	6,6,6	0.54	0
2	SO4	B	402	-	4,4,4	0.35	0	6,6,6	0.36	0
2	SO4	B	401	-	4,4,4	0.36	0	6,6,6	0.69	0
2	SO4	C	401	-	4,4,4	0.31	0	6,6,6	0.37	0
2	SO4	F	402	-	4,4,4	0.40	0	6,6,6	0.46	0
2	SO4	F	401	-	4,4,4	0.29	0	6,6,6	0.63	0
2	SO4	D	402	-	4,4,4	0.90	0	6,6,6	0.42	0
2	SO4	E	402	-	4,4,4	0.48	0	6,6,6	0.32	0
2	SO4	A	402	-	4,4,4	0.34	0	6,6,6	0.25	0
2	SO4	E	401	-	4,4,4	0.42	0	6,6,6	0.68	0
2	SO4	A	401	-	4,4,4	0.33	0	6,6,6	0.30	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	309/326 (94%)	0.18	15 (4%) 35 36	28, 47, 76, 92	0
1	B	315/326 (96%)	-0.02	10 (3%) 50 52	29, 45, 69, 87	0
1	C	296/326 (90%)	-0.02	6 (2%) 65 66	30, 44, 70, 95	0
1	D	304/326 (93%)	0.48	29 (9%) 14 15	30, 48, 93, 117	0
1	E	314/326 (96%)	0.05	14 (4%) 38 40	29, 42, 69, 94	0
1	F	315/326 (96%)	0.10	8 (2%) 58 60	31, 46, 68, 87	0
All	All	1853/1956 (94%)	0.13	82 (4%) 39 41	28, 45, 73, 117	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	206	HIS	5.7
1	D	203	VAL	5.6
1	A	84	ALA	5.5
1	B	318	ALA	5.4
1	F	318	ALA	5.1
1	D	87	PRO	5.0
1	D	5	THR	4.8
1	D	205	ALA	4.7
1	D	84	ALA	4.5
1	C	98	VAL	4.3
1	D	95	LEU	4.3
1	D	93	THR	4.2
1	D	200	ALA	4.2
1	A	87	PRO	3.8
1	B	84	ALA	3.7
1	A	4	ILE	3.7
1	A	93	THR	3.5
1	D	103	LYS	3.4
1	E	226	TYR	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	204	LYS	3.4
1	E	5	THR	3.4
1	F	89	LYS	3.4
1	D	211	GLU	3.2
1	D	316	ALA	3.2
1	D	297	HIS	3.2
1	C	313	LEU	3.2
1	D	202	TRP	3.1
1	D	213	LYS	3.1
1	D	314	THR	3.0
1	A	317	PHE	2.9
1	E	48	LYS	2.9
1	E	207	PRO	2.8
1	A	85	GLY	2.8
1	D	102	LEU	2.8
1	E	204	LYS	2.8
1	A	5	THR	2.7
1	D	94	ARG	2.6
1	D	199	ILE	2.6
1	E	209	ILE	2.6
1	D	4	ILE	2.6
1	D	99	ASN	2.5
1	A	213	LYS	2.5
1	B	7	LYS	2.5
1	D	226	TYR	2.5
1	A	102	LEU	2.5
1	A	311	LYS	2.5
1	B	288	ASN	2.5
1	F	84	ALA	2.4
1	E	196	GLY	2.4
1	C	311	LYS	2.4
1	B	5	THR	2.4
1	A	137	LYS	2.4
1	E	137	LYS	2.3
1	E	210	LYS	2.3
1	C	217	MET	2.3
1	A	303	GLN	2.3
1	B	42	ILE	2.3
1	D	42	ILE	2.3
1	F	42	ILE	2.3
1	A	95	LEU	2.3
1	B	304	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	212	ASP	2.2
1	E	200	ALA	2.2
1	E	248	LYS	2.2
1	E	199	ILE	2.2
1	B	248	LYS	2.2
1	F	297	HIS	2.2
1	C	292	ILE	2.1
1	B	204	LYS	2.1
1	A	94	ARG	2.1
1	D	133	TYR	2.1
1	E	42	ILE	2.1
1	F	288	ASN	2.1
1	E	317	PHE	2.1
1	F	85	GLY	2.1
1	D	137	LYS	2.1
1	F	4	ILE	2.1
1	C	137	LYS	2.0
1	D	304	LYS	2.0
1	A	126	ASN	2.0
1	D	85	GLY	2.0
1	B	103	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	F	401	5/5	0.88	0.13	55,56,64,65	0
2	SO4	B	401	5/5	0.93	0.09	49,56,56,65	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	E	402	5/5	0.95	0.08	48,49,51,52	0
2	SO4	E	401	5/5	0.96	0.07	47,48,52,54	0
2	SO4	A	401	5/5	0.96	0.07	49,52,56,59	0
2	SO4	D	401	5/5	0.96	0.08	46,52,56,57	0
2	SO4	C	402	5/5	0.98	0.06	39,40,42,46	0
2	SO4	C	401	5/5	0.98	0.05	50,50,52,53	0
2	SO4	D	402	5/5	0.98	0.07	39,43,45,46	0
2	SO4	F	402	5/5	0.98	0.06	39,40,42,42	0
2	SO4	A	402	5/5	0.99	0.05	39,39,42,44	0
2	SO4	B	402	5/5	0.99	0.05	41,41,43,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.