



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 01:49 PM UTC

PDB ID : 7JMP / pdb_00007jmp
Title : Crystal structure of SARS-CoV-2 receptor binding domain in complex with neutralizing antibody COVA2-39
Authors : Wu, N.C.; Yuan, M.; Liu, H.; Zhu, X.; Wilson, I.A.
Deposited on : 2020-08-02
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	14% 63% 8% 29%
2	H	232	2% 87% 6% 6%
3	L	216	2% 94% ••

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4953 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	165	Total	C	N	O	S	0	0	0
			1274	822	210	238	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	542	SER	-	expression tag	UNP P0DTC2
A	543	GLY	-	expression tag	UNP P0DTC2
A	544	HIS	-	expression tag	UNP P0DTC2
A	545	HIS	-	expression tag	UNP P0DTC2
A	546	HIS	-	expression tag	UNP P0DTC2
A	547	HIS	-	expression tag	UNP P0DTC2
A	548	HIS	-	expression tag	UNP P0DTC2
A	549	HIS	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called COVA2-39 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	218	Total	C	N	O	S	0	1	0
			1620	1017	276	320	7			

- Molecule 3 is a protein called COVA2-39 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	L	211	Total	C	N	O	S	0	4	0
			1563	980	258	319	6			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	87	Total	O	0	0
			87	87		
5	H	186	Total	O	0	0
			186	186		
5	L	209	Total	O	0	0
			209	209		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1



- Molecule 2: COVA2-39 heavy chain



- Molecule 3: COVA2-39 light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.91Å 80.44Å 72.01Å 90.00° 104.88° 90.00°	Depositor
Resolution (Å)	34.82 – 1.71 34.82 – 1.71	Depositor EDS
% Data completeness (in resolution range)	96.6 (34.82-1.71) 96.7 (34.82-1.71)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.71Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.177 , 0.209 0.179 , 0.211	Depositor DCC
R_{free} test set	3845 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.183	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4953	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	2/1309 (0.2%)	0.63	0/1778
2	H	0.71	4/1656 (0.2%)	0.77	0/2258
3	L	0.54	0/1611	0.70	0/2203
All	All	0.61	6/4576 (0.1%)	0.71	0/6239

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	108	ARG	CD-NE	-8.59	1.34	1.46
2	H	109	VAL	C-O	-7.70	1.15	1.24
1	A	483	VAL	C-O	-6.92	1.16	1.24
2	H	107	THR	C-O	-6.73	1.16	1.24
1	A	484	GLU	C-O	-5.53	1.17	1.23
2	H	108	ARG	C-O	-5.22	1.17	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1274	0	1140	12	0
2	H	1620	0	1574	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	L	1563	0	1503	7	0
4	A	14	0	13	0	0
5	A	87	0	0	1	0
5	H	186	0	0	2	0
5	L	209	0	0	2	0
All	All	4953	0	4230	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:129:LYS:NZ	5:L:301:HOH:O	1.92	1.01
1:A:355:ARG:HD3	1:A:398:ASP:OD1	1.72	0.89
2:H:199:ASN:HD21	2:H:201:LYS:HE2	1.48	0.78
1:A:381:GLY:HA3	1:A:430:THR:HG22	1.73	0.70
2:H:199:ASN:ND2	2:H:201:LYS:HE2	2.18	0.58
1:A:417:LYS:HD2	5:H:469:HOH:O	2.05	0.54
3:L:194:GLN:HG2	3:L:203:GLU:HG3	1.90	0.54
2:H:87:THR:CG2	2:H:108:ARG:NH2	2.72	0.53
1:A:424:LYS:HD3	5:A:1104:HOH:O	2.08	0.52
2:H:4:LEU:HD22	2:H:22:CYS:SG	2.50	0.52
1:A:353:TRP:O	1:A:466:ARG:NH1	2.44	0.51
2:H:167:PRO:HG2	3:L:165:SER:OG	2.13	0.48
2:H:87:THR:HG22	2:H:108:ARG:NH2	2.28	0.48
2:H:42:GLY:HA3	3:L:163:THR:HG21	1.97	0.46
1:A:396:TYR:HB2	1:A:514:SER:OG	2.15	0.45
3:L:120:PRO:HD3	3:L:132:LEU:HD12	1.99	0.45
2:H:42:GLY:C	2:H:43:LYS:HD2	2.42	0.45
3:L:129:LYS:HE3	3:L:129:LYS:HB2	1.68	0.45
1:A:376:THR:HB	1:A:435:ALA:HB3	1.98	0.44
1:A:431:GLY:HA2	1:A:515:PHE:CE2	2.52	0.44
2:H:171:GLN:HG3	2:H:175:LEU:O	2.18	0.44
1:A:354:ASN:O	1:A:398:ASP:HA	2.18	0.43
1:A:498:GLN:H	1:A:501:ASN:ND2	2.17	0.42
2:H:199:ASN:HD21	2:H:201:LYS:CE	2.27	0.42
1:A:431:GLY:HA2	1:A:515:PHE:CD2	2.55	0.42
3:L:53:LYS:NZ	5:L:304:HOH:O	2.41	0.42
2:H:43:LYS:HA	2:H:43:LYS:HE2	2.03	0.41
2:H:13:GLN:H	2:H:13:GLN:CD	2.30	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:TRP:HZ3	1:A:355:ARG:HE	1.70	0.40
2:H:100(C):GLU:HB3	5:H:374:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/231 (68%)	149 (95%)	8 (5%)	0	100	100
2	H	215/232 (93%)	211 (98%)	4 (2%)	0	100	100
3	L	213/216 (99%)	208 (98%)	5 (2%)	0	100	100
All	All	585/679 (86%)	568 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/203 (62%)	126 (100%)	0	100	100
2	H	179/194 (92%)	178 (99%)	1 (1%)	78	68
3	L	174/181 (96%)	172 (99%)	2 (1%)	65	50
All	All	479/578 (83%)	476 (99%)	3 (1%)	84	68

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	149	PRO
3	L	137[A]	SER
3	L	137[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	493	GLN
1	A	501	ASN
2	H	39	GLN
3	L	38	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1001	1	14,14,15	1.58	1 (7%)	17,19,21	2.18	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1001	1	-	4/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1001	NAG	O5-C1	5.81	1.53	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1001	NAG	C1-O5-C5	8.75	123.92	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1001	NAG	O5-C5-C6-O6
4	A	1001	NAG	C4-C5-C6-O6
4	A	1001	NAG	C8-C7-N2-C2
4	A	1001	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	165/231 (71%)	1.08	33 (20%) 3 5	21, 45, 81, 94	0
2	H	218/232 (93%)	0.06	4 (1%) 67 74	16, 28, 42, 54	1 (0%)
3	L	211/216 (97%)	-0.07	4 (1%) 66 73	15, 27, 40, 51	4 (1%)
All	All	594/679 (87%)	0.30	41 (6%) 23 28	15, 30, 61, 94	5 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	367	VAL	5.5
1	A	355	ARG	5.3
1	A	368	LEU	5.2
1	A	384	PRO	5.0
1	A	383	SER	4.8
3	L	3	ALA	4.7
1	A	369	TYR	4.6
1	A	395	VAL	4.3
1	A	366	SER	4.2
3	L	209	THR	3.9
2	H	133	GLY	3.8
1	A	392	PHE	3.6
1	A	359	SER	3.5
1	A	393	THR	3.5
1	A	338	PHE	3.4
2	H	127	SER	3.2
1	A	382	VAL	3.2
1	A	379	CYS	3.2
1	A	415	THR	3.1
1	A	445	VAL	2.9
1	A	377	PHE	2.8
1	A	358	ILE	2.8
1	A	380	TYR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	214	LYS	2.7
1	A	513	LEU	2.5
1	A	515	PHE	2.5
2	H	92	CYS	2.5
1	A	446	GLY	2.5
1	A	371	SER	2.4
1	A	514	SER	2.4
1	A	394	ASN	2.3
1	A	357	ARG	2.3
1	A	372	ALA	2.3
1	A	432	CYS	2.3
1	A	381	GLY	2.3
1	A	413	GLY	2.3
3	L	126	GLN	2.2
1	A	441	LEU	2.2
1	A	505	TYR	2.2
3	L	153	SER	2.1
1	A	396	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	A	1001	14/15	0.74	0.14	80,83,88,89	0

6.5 Other polymers [i](#)

There are no such residues in this entry.