



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 09:17 AM UTC

PDB ID : 1JNV / pdb_00001jnv
Title : The Conformation of the Epsilon and Gamma Subunits within the E. coli F1 ATPase
Authors : Hausrath, A.C.; Capaldi, R.A.; Matthews, B.W.
Deposited on : 2001-07-25
Resolution : 4.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1135 (4.90-3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	492	 87% 12% .
1	B	492	 86% 13% .
1	C	492	 84% 16%
2	D	467	 91% 9%
2	E	467	 86% 14%
2	F	467	 91% 8%
3	Y	135	 86% 10% .
4	Z	287	 81% 13% . 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13072 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP SYNTHASE ALPHA CHAIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	487	Total	C	N	O	0	0	0
			1948	974	487	487			
1	B	487	Total	C	N	O	32	0	0
			1948	974	487	487			
1	C	492	Total	C	N	O	0	0	0
			1968	984	492	492			

- Molecule 2 is a protein called ATP SYNTHASE BETA CHAIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	467	Total	C	N	O	0	0	0
			1868	934	467	467			
2	E	466	Total	C	N	O	0	0	0
			1864	932	466	466			
2	F	466	Total	C	N	O	0	0	0
			1864	932	466	466			

- Molecule 3 is a protein called ATP SYNTHASE EPSILON CHAIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	Y	130	Total	C	N	O	0	0	0
			520	260	130	130			


- Molecule 4 is a protein called ATP SYNTHASE GAMMA CHAIN.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	Z	273	Total	C	N	O	0	0	0
			1092	546	273	273			

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP SYNTHASE ALPHA CHAIN






• Molecule 2: ATP SYNTHASE BETA CHAIN

Chain E: 86% 14%




• Molecule 2: ATP SYNTHASE BETA CHAIN

Chain F: 91% 8%




• Molecule 3: ATP SYNTHASE EPSILON CHAIN

Chain Y: 86% 10%



• Molecule 4: ATP SYNTHASE GAMMA CHAIN

Chain Z: 81% 13%



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	180.68Å 197.52Å 237.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 4.40 25.00 – 4.40	Depositor EDS
% Data completeness (in resolution range)	64.5 (25.00-4.40) 64.1 (25.00-4.40)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.58 (at 4.40Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.416 , (Not available) 0.396 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	168.3	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.14 , 999.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.81	EDS
Total number of atoms	13072	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
2	D	0	1
2	F	0	1
3	Y	0	3
4	Z	0	15
All	All	0	22

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 22 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	379	UNK	Mainchain
1	C	379	UNK	Mainchain
2	D	82	UNK	Mainchain
2	F	82	UNK	Mainchain
3	Y	13	UNK	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1948	0	21	34	0
1	B	1948	0	20	36	0
1	C	1968	0	19	48	17
2	D	1868	0	13	32	0
2	E	1864	0	17	37	0
2	F	1864	0	12	38	0
3	Y	520	0	15	17	17
4	Z	1092	0	20	22	0
All	All	13072	0	137	242	17

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 242 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:UNK:C	1:A:96:UNK:H2	1.07	1.61
1:A:95:UNK:C	1:A:96:UNK:N	1.75	1.50
2:D:363:UNK:C	2:D:364:UNK:N	1.85	1.39
2:F:392:UNK:C	3:Y:131:UNK:CA	2.03	1.36
2:F:392:UNK:CA	3:Y:131:UNK:C	2.19	1.20

The worst 5 of 17 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:20:UNK:C	3:Y:115:UNK:O[8_456]	0.78	1.42
1:C:20:UNK:CA	3:Y:115:UNK:O[8_456]	1.11	1.09
1:C:20:UNK:O	3:Y:115:UNK:C[8_456]	1.24	0.96
1:C:20:UNK:O	3:Y:115:UNK:O[8_456]	1.31	0.89
1:C:20:UNK:N	3:Y:116:UNK:C[8_456]	1.42	0.78

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
4	Z	4
1	C	2
2	F	2
2	D	2
1	A	2
1	B	2
2	E	1

The worst 5 of 15 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	Z	248:UNK	C	249:UNK	N	3.57
1	C	95:UNK	C	96:UNK	N	2.43
1	F	363:UNK	C	364:UNK	N	2.24
1	D	363:UNK	C	364:UNK	N	1.85
1	A	95:UNK	C	96:UNK	N	1.75

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB($\text{\AA}^2$)	Q<0.9
1	A	0/492	-	-	-	-
1	B	0/492	-	-	-	-
1	C	0/492	-	-	-	-
2	D	0/467	-	-	-	-
2	E	0/467	-	-	-	-
2	F	0/467	-	-	-	-
3	Y	0/135	-	-	-	-
4	Z	0/287	-	-	-	-
All	All	0/3299	-	-	-	-

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.