



Full wwPDB NMR Structure Validation Report ⓘ

Mar 9, 2026 – 08:22 PM UTC

PDB ID : 2JN9 / pdb_00002jn9
Title : NMR solution structure of YkvR protein from Bacillus subtilis: NESG target SR358
Authors : Gurla, S.V.T.; Aramini, J.M.; Chi Ho, K.; Cunningham, K.; Ma, L.-C.; Xiao, R.; Baran, M.C.; Acton, T.B.; Liu, J.; Rost, B.; Montelione, G.T.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2006-12-29

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	105	 64% 13% • 22%

2 Ensemble composition and analysis

This entry contains 10 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *minimized average structure*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:71, A:82-A:92 (82)	1.44	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 3, 4, 5, 7, 8
2	1, 9, 10
Single-model clusters	6

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1694 atoms, of which 832 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called YkvR protein.

Mol	Chain	Residues	Atoms						Trace
1	A	105	Total	C	H	N	O	S	0
			1694	543	832	144	172	3	

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	cloning artifact	UNP O31683
A	2	VAL	-	cloning artifact	UNP O31683
A	98	LEU	-	cloning artifact	UNP O31683
A	99	GLU	-	cloning artifact	UNP O31683
A	100	HIS	-	expression tag	UNP O31683
A	101	HIS	-	expression tag	UNP O31683
A	102	HIS	-	expression tag	UNP O31683
A	103	HIS	-	expression tag	UNP O31683
A	104	HIS	-	expression tag	UNP O31683
A	105	HIS	-	expression tag	UNP O31683

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: YkvR protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: YkvR protein



4.2.2 Score per residue for model 2


- Molecule 1: YkvR protein



H105


4.2.3 Score per residue for model 3

- Molecule 1: YkvR protein




4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: YkvR protein




4.2.5 Score per residue for model 5

- Molecule 1: YkvR protein



4.2.6 Score per residue for model 6

- Molecule 1: YkvR protein



H100
H101
H102
H103
H104
H105

4.2.7 Score per residue for model 7

- Molecule 1: YkvR protein

Chain A: 59% 14% • • 22%

M1 V2 K3 T4 L5 N8 L12 E13 M14 Y17 E20 S21 E22 P23 K24 R25 Y46 Y59 A60 F61 M65 T66 L72 T73 N74 L75 Y76 E77 P78 G79 A80 V81 E87 I88 T89 D92 K93 N94 A95 D96 S97 L98 H100 H101 H102 H103 H104 H105

4.2.8 Score per residue for model 8

- Molecule 1: YkvR protein

Chain A: 61% 14% • 22%

M1 V2 K3 Q18 R25 T30 V33 T34 S35 L45 Y46 R57 T66 L72 T73 N74 L75 Y76 E77 P78 G79 A80 V81 S82 E83 F84 Y85 I86 E87 I88 T89 E90 I91 D92 K93 N94 A95 D96 S97 L98 H100 H101 H102 H103 H104 H105

4.2.9 Score per residue for model 9

- Molecule 1: YkvR protein

Chain A: 60% 17% • 22%

M1 L5 R25 F29 N32 V33 T34 T37 I41 L45 Y46 E47 V52 L59 E64 M65 I66 N67 L72 T73 N74 L75 Y76 E77 P78 G79 A80 V81 S82 I86 E90 K93 N94 A95 D96 S97 L98 H100 H101 H102 H103 H104 H105

4.2.10 Score per residue for model 10

- Molecule 1: YkvR protein

Chain A: 48% 27% • • 22%

M1 V2 R6 L7 N8 N9 Y10 T11 L12 E13 M14 Y17 Q18 E19 E22 P23 K24 K26 F29 N32 V33 T34 T37 I41 A42 Y43 L44 L45 Y46 E47 M51 E56 R57 D58 L59 M65 L72 T73 N74 L75 Y76 E77 P78 G79 A80 V81 S82 I86 K93 N94 A95 D96 S97 L98 H100 H101 H102 H103 H104 H105

5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry*.

Of the 56 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
AutoStructure	refinement	
AutoStructure	structure solution	
DYANA	structure solution	
DYANA	refinement	
CNS	refinement	1.1
X-PLOR	refinement	2.0.6

No chemical shift data was provided.

6 Model quality

6.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.46±0.01	0±0/685 (0.0± 0.0%)	0.74±0.05	1±1/929 (0.1± 0.1%)
All	All	0.46	0/6850 (0.0%)	0.74	8/9290 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	0.4±0.5
All	All	0	4

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	22	GLU	CA-C-N	6.66	128.17	119.84	7	3
1	A	22	GLU	C-N-CA	6.66	128.17	119.84	7	3
1	A	92	ASP	CB-CA-C	-5.69	108.38	115.89	7	1
1	A	47	GLU	CB-CA-C	-5.15	110.62	116.54	5	1

There are no chirality outliers.

All unique planar outliers are listed below.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	22	GLU	Peptide	4

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	673	660	659	11±3
All	All	6730	6600	6590	114

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:48:LYS:HA	1:A:65:MET:HG3	0.76	1.55	5	1
1:A:14:MET:HG3	1:A:17:TYR:HB2	0.71	1.60	3	1
1:A:20:GLU:O	1:A:25:ARG:HA	0.70	1.85	1	2
1:A:22:GLU:HB3	1:A:23:PRO:HD2	0.70	1.63	7	2
1:A:20:GLU:HG3	1:A:23:PRO:HD3	0.70	1.61	6	1
1:A:22:GLU:HB3	1:A:23:PRO:CD	0.69	2.18	1	3
1:A:46:TYR:HB3	1:A:65:MET:SD	0.66	2.31	10	1
1:A:71:SER:HB2	1:A:85:TYR:CD2	0.65	2.27	6	1
1:A:32:ASN:HA	1:A:82:SER:O	0.63	1.94	10	3
1:A:33:VAL:HG21	1:A:84:PHE:CD1	0.62	2.30	4	4
1:A:56:GLU:HG3	1:A:57:ARG:HD2	0.60	1.73	10	1
1:A:64:GLU:HG2	1:A:91:ILE:HA	0.58	1.75	3	1
1:A:48:LYS:HG3	1:A:49:THR:H	0.58	1.58	3	1
1:A:46:TYR:CD1	1:A:65:MET:HE2	0.58	2.33	3	1
1:A:14:MET:SD	1:A:17:TYR:HB2	0.58	2.39	7	2
1:A:29:PHE:CE1	1:A:86:ILE:HB	0.58	2.34	9	2
1:A:37:THR:O	1:A:41:ILE:HG12	0.57	2.00	2	1
1:A:48:LYS:N	1:A:48:LYS:HD2	0.57	2.15	2	1
1:A:8:ASN:HA	1:A:51:ASN:ND2	0.56	2.16	5	1
1:A:8:ASN:HA	1:A:51:ASN:HD21	0.54	1.63	5	1
1:A:59:LEU:HD22	1:A:61:PHE:HB2	0.54	1.80	7	2
1:A:24:LYS:HB3	1:A:90:GLU:HB3	0.54	1.80	6	1
1:A:17:TYR:CD2	1:A:29:PHE:HB3	0.53	2.38	10	1
1:A:46:TYR:HB2	1:A:65:MET:SD	0.52	2.43	4	1
1:A:66:THR:OG1	1:A:89:THR:HB	0.52	2.05	8	2
1:A:61:PHE:CZ	1:A:90:GLU:HA	0.50	2.40	4	1
1:A:18:GLN:HB3	1:A:28:ALA:O	0.50	2.07	6	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:45:LEU:HG	1:A:46:TYR:CE1	0.50	2.42	4	1
1:A:71:SER:HB3	1:A:85:TYR:HD2	0.49	1.67	5	1
1:A:45:LEU:O	1:A:46:TYR:HB2	0.49	2.08	9	1
1:A:22:GLU:CB	1:A:23:PRO:HD2	0.49	2.35	7	2
1:A:45:LEU:O	1:A:45:LEU:HG	0.48	2.08	8	2
1:A:61:PHE:HZ	1:A:90:GLU:HA	0.48	1.68	4	1
1:A:71:SER:CB	1:A:85:TYR:HB2	0.48	2.39	1	1
1:A:33:VAL:HG13	1:A:38:TYR:HA	0.48	1.86	6	1
1:A:65:MET:HG2	1:A:88:ILE:HG12	0.48	1.85	6	1
1:A:45:LEU:HA	1:A:84:PHE:HE2	0.48	1.69	8	1
1:A:47:GLU:N	1:A:65:MET:SD	0.48	2.87	9	1
1:A:50:PHE:HB2	1:A:88:ILE:HD13	0.47	1.85	1	1
1:A:24:LYS:HE2	1:A:90:GLU:HB3	0.47	1.84	3	1
1:A:30:THR:CG2	1:A:83:GLU:HB3	0.47	2.39	4	3
1:A:8:ASN:HB2	1:A:51:ASN:HB2	0.47	1.86	6	1
1:A:20:GLU:HG3	1:A:23:PRO:CD	0.47	2.38	6	1
1:A:45:LEU:H	1:A:45:LEU:HD23	0.47	1.69	8	1
1:A:47:GLU:HB2	1:A:50:PHE:CZ	0.46	2.45	2	1
1:A:6:ARG:HB3	1:A:11:THR:HA	0.46	1.88	10	1
1:A:22:GLU:CB	1:A:23:PRO:CD	0.46	2.93	7	3
1:A:65:MET:SD	1:A:88:ILE:HG12	0.46	2.51	3	1
1:A:53:GLU:HA	1:A:59:LEU:O	0.46	2.11	1	1
1:A:6:ARG:HG3	1:A:53:GLU:HB2	0.46	1.88	4	1
1:A:3:LYS:HB2	1:A:3:LYS:NZ	0.46	2.26	8	2
1:A:33:VAL:HB	1:A:82:SER:O	0.46	2.11	6	1
1:A:12:LEU:HD21	1:A:29:PHE:CE1	0.45	2.46	6	1
1:A:26:LYS:HD2	1:A:87:GLU:CD	0.45	2.36	2	1
1:A:66:THR:OG1	1:A:89:THR:HG22	0.45	2.11	6	1
1:A:91:ILE:HG12	1:A:92:ASP:N	0.45	2.26	8	1
1:A:65:MET:CE	1:A:86:ILE:HG23	0.45	2.41	10	1
1:A:85:TYR:CE1	1:A:87:GLU:HB2	0.45	2.47	8	1
1:A:22:GLU:H	1:A:23:PRO:HD3	0.45	1.72	5	1
1:A:27:ILE:HG23	1:A:88:ILE:HB	0.45	1.89	4	1
1:A:5:LEU:HD11	1:A:52:VAL:HG13	0.44	1.89	9	1
1:A:14:MET:SD	1:A:14:MET:N	0.44	2.90	3	1
1:A:25:ARG:HB2	1:A:90:GLU:HB3	0.44	1.89	5	1
1:A:48:LYS:HA	1:A:65:MET:CG	0.44	2.38	5	1
1:A:33:VAL:O	1:A:82:SER:HB2	0.44	2.13	9	1
1:A:25:ARG:O	1:A:90:GLU:HB2	0.43	2.13	9	1
1:A:22:GLU:H	1:A:23:PRO:CD	0.43	2.26	5	1
1:A:65:MET:HE2	1:A:86:ILE:HG23	0.43	1.90	10	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:17:TYR:HA	1:A:29:PHE:HB3	0.43	1.90	6	1
1:A:5:LEU:HD23	1:A:12:LEU:HD23	0.43	1.90	7	1
1:A:46:TYR:HB2	1:A:65:MET:HE1	0.43	1.90	7	1
1:A:46:TYR:CD1	1:A:65:MET:HB3	0.42	2.49	9	1
1:A:22:GLU:HB2	1:A:23:PRO:HD3	0.42	1.90	3	1
1:A:22:GLU:C	1:A:24:LYS:H	0.42	2.21	7	1
1:A:22:GLU:O	1:A:24:LYS:N	0.42	2.45	1	2
1:A:16:ALA:HB3	1:A:30:THR:HB	0.42	1.92	2	2
1:A:28:ALA:HA	1:A:87:GLU:HA	0.42	1.91	3	2
1:A:45:LEU:HA	1:A:84:PHE:CE2	0.42	2.49	8	1
1:A:37:THR:HG23	1:A:41:ILE:HD11	0.42	1.92	9	1
1:A:1:MET:HE3	1:A:17:TYR:O	0.42	2.15	10	1
1:A:27:ILE:HG13	1:A:88:ILE:HB	0.42	1.92	6	1
1:A:22:GLU:N	1:A:23:PRO:CD	0.41	2.83	5	2
1:A:12:LEU:HD21	1:A:29:PHE:CZ	0.41	2.50	10	1
1:A:46:TYR:HD1	1:A:65:MET:SD	0.41	2.39	9	1
1:A:37:THR:HB	1:A:41:ILE:HD12	0.41	1.93	3	1
1:A:65:MET:HG3	1:A:88:ILE:HD13	0.41	1.93	4	1
1:A:65:MET:HE2	1:A:87:GLU:O	0.41	2.16	2	1
1:A:45:LEU:O	1:A:47:GLU:N	0.41	2.54	3	1
1:A:8:ASN:HA	1:A:51:ASN:HB3	0.40	1.93	10	1
1:A:5:LEU:HD11	1:A:52:VAL:CG1	0.40	2.47	9	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	81/105 (77%)	69±3 (85±4%)	9±3 (12±4%)	2±1 (3±2%)	5	38
All	All	810/1050 (77%)	692 (85%)	94 (12%)	24 (3%)	5	38

All 11 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	2	VAL	5
1	A	23	PRO	4
1	A	46	TYR	4
1	A	47	GLU	3
1	A	48	LYS	2
1	A	24	LYS	1
1	A	22	GLU	1
1	A	69	SER	1
1	A	8	ASN	1
1	A	67	ASN	1
1	A	9	ASN	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	76/96 (79%)	71±2 (93±2%)	5±2 (7±2%)	17	66
All	All	760/960 (79%)	710 (93%)	50 (7%)	17	66

All 28 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	59	LEU	7
1	A	91	ILE	4
1	A	89	THR	3
1	A	84	PHE	3
1	A	3	LYS	3
1	A	32	ASN	2
1	A	65	MET	2
1	A	35	SER	2
1	A	22	GLU	2
1	A	30	THR	2
1	A	66	THR	2
1	A	57	ARG	2
1	A	14	MET	1
1	A	34	THR	1
1	A	47	GLU	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	45	LEU	1
1	A	56	GLU	1
1	A	27	ILE	1
1	A	46	TYR	1
1	A	2	VAL	1
1	A	38	TYR	1
1	A	87	GLU	1
1	A	18	GLN	1
1	A	25	ARG	1
1	A	64	GLU	1
1	A	19	GLU	1
1	A	26	LYS	1
1	A	37	THR	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided