



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 26, 2026 – 03:26 AM EDT

PDB ID : 7JNH / pdb_00007jnh
Title : Crystal structure of a double-ENE RNA stability element in complex with a 28-mer poly(A) RNA
Authors : Torabi, S.F.; Vaidya, A.T.; Tycowski, K.T.; DeGregorio, S.J.; Wang, J.; Shu, M.D.; Steitz, T.A.; Steitz, J.A.
Deposited on : 2020-08-04
Resolution : 2.89 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)
RNA backbone	3983	1120 (3.10-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	28	
2	B	86	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 28-mer poly(A) RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	25	Total	C	N	O	P	0	0	1
			508	230	115	140	23			

- Molecule 2 is a RNA chain called Core double ENE RNA (Xtal construct) from Oryza sativa transposon,Core double ENE RNA (Xtal construct) from Oryza sativa transposon.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	86	Total	C	N	O	P	0	0	0
			1816	812	304	614	86			


There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	G	-	insertion	GB AL442105.2
B	2	G	-	insertion	GB AL442105.2
B	3	G	-	insertion	GB AL442105.2
B	4	C	-	insertion	GB AL442105.2
B	41	G	-	linker	GB AL442105.2
B	42	G	-	linker	GB AL442105.2
B	43	C	-	linker	GB AL442105.2
B	44	G	-	linker	GB AL442105.2
B	45	A	-	linker	GB AL442105.2
B	46	A	-	linker	GB AL442105.2
B	47	A	-	linker	GB AL442105.2
B	48	G	-	linker	GB AL442105.2
B	49	U	-	linker	GB AL442105.2
B	50	C	-	linker	GB AL442105.2
B	83	G	-	insertion	GB AL442105.2
B	84	C	-	insertion	GB AL442105.2
B	85	C	-	insertion	GB AL442105.2
B	86	CCC	-	insertion	GB AL442105.2

- Molecule 3 is STRONTIUM ION (CCD ID: SR) (formula: Sr).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total 3	Sr 3	0	0
3	B	3	Total 3	Sr 3	0	0

- Molecule 4 is COBALT HEXAMMINE(III) (CCD ID: NCO) (formula: CoH₁₈N₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 7	Co 1	N 6	0	0
4	B	1	Total 7	Co 1	N 6	0	0
4	B	1	Total 7	Co 1	N 6	0	0
4	B	1	Total 7	Co 1	N 6	0	0
4	B	1	Total 7	Co 1	N 6	0	0
4	B	1	Total 7	Co 1	N 6	0	0
4	B	1	Total 7	Co 1	N 6	0	0
4	B	1	Total 7	Co 1	N 6	0	0
4	B	1	Total 7	Co 1	N 6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	Co	N	0	0
			7	1	6		
4	B	1	Total	Co	N	0	0
			7	1	6		
4	B	1	Total	Co	N	0	0
			7	1	6		

- Molecule 5 is SPERMIDINE (CCD ID: SPD) (formula: $C_7H_{19}N_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	N	0	0
			10	7	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	3	Total	Mg	0	0
			3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	5	Total	O	0	0
			5	5		
7	B	13	Total	O	0	0
			13	13		

4 Data and refinement statistics

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants a, b, c, α , β , γ	93.97Å 93.97Å 202.17Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.03 – 2.89 47.03 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.1 (47.03-2.89) 95.1 (47.03-2.90)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.20 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.184 , 0.225 0.183 , 0.232	Depositor DCC
R_{free} test set	581 reflections (4.66%)	wwPDB-VP
Wilson B-factor (Å ²)	114.6	Xtriage
Anisotropy	0.283	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 95.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2445	wwPDB-VP
Average B, all atoms (Å ²)	115.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SPD, SR, NCO, CCC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/575	1.19	6/895 (0.7%)
2	B	0.59	0/2000	1.15	19/3111 (0.6%)
All	All	0.57	0/2575	1.16	25/4006 (0.6%)

There are no bond length outliers.

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	G	C2'-C3'-O3'	11.23	130.55	113.70
1	A	6	A	C4'-C3'-O3'	-9.78	98.32	113.00
1	A	10	A	C1'-C2'-O2'	-7.78	100.14	111.80
2	B	58	U	C3'-C2'-O2'	7.64	122.15	110.70
1	A	23	A	C3'-C2'-O2'	6.56	120.55	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	508	0	255	21	0
2	B	1816	0	916	34	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	7	0	0	1	0
4	B	77	0	0	4	0
5	B	10	0	19	0	0
6	B	3	0	0	0	0
7	A	5	0	0	0	0
7	B	13	0	0	0	0
All	All	2445	0	1190	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:A:OP1	1:A:10:A:H4'	1.67	0.95
1:A:15:A:OP2	2:B:14:A:O2'	1.90	0.89
1:A:24:A:O2'	1:A:25:A:O4'	1.97	0.81
2:B:69:U:H2'	2:B:70:C:O4'	1.88	0.73
1:A:21:A:OP1	4:A:104:NCO:N1	2.23	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	21/28 (75%)	16 (76%)	5 (23%)
2	B	85/86 (98%)	17 (20%)	1 (1%)
All	All	106/114 (92%)	33 (31%)	6 (5%)

5 of 33 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	2	A
1	A	3	A
1	A	4	A
1	A	5	A
1	A	6	A

5 of 6 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	15	A
1	A	24	A
2	B	3	G
1	A	6	A
1	A	5	A

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CCC	B	86	2	20,25,26	1.85	4 (20%)	27,38,41	1.57	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CCC	B	86	2	-	1/7/35/36	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	86	CCC	PC-O1C	5.05	1.68	1.50
2	B	86	CCC	C2'-C3'	-4.05	1.44	1.53
2	B	86	CCC	PC-O2C	2.73	1.68	1.55
2	B	86	CCC	C6-C5	2.44	1.40	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	86	CCC	O4'-C1'-N1	3.33	115.91	108.36
2	B	86	CCC	O4'-C1'-C2'	-3.30	100.91	106.59
2	B	86	CCC	O2-C2-N3	-3.28	117.15	122.33

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	86	CCC	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 9 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NCO	B	108	-	6,6,6	0.69	0	-		
4	NCO	B	111	-	6,6,6	0.79	0	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NCO	B	110	-	6,6,6	0.82	0	-		
4	NCO	B	112	-	6,6,6	0.69	0	-		
4	NCO	B	109	-	6,6,6	0.68	0	-		
5	SPD	B	115	-	9,9,9	0.20	0	8,8,8	0.21	0
4	NCO	B	114	-	6,6,6	0.67	0	-		
4	NCO	B	113	-	6,6,6	0.77	0	-		
4	NCO	B	107	-	6,6,6	0.71	0	-		
4	NCO	B	104	-	6,6,6	0.70	0	-		
4	NCO	B	106	-	6,6,6	0.85	0	-		
4	NCO	A	104	-	6,6,6	0.51	0	-		
4	NCO	B	105	-	6,6,6	0.64	0	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	SPD	B	115	-	-	3/7/7/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	115	SPD	C8-C7-N6-C5
5	B	115	SPD	C3-C4-C5-N6
5	B	115	SPD	C2-C3-C4-C5

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	111	NCO	1	0
4	B	109	NCO	1	0
4	B	114	NCO	1	0
4	B	104	NCO	1	0
4	A	104	NCO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	25/28 (89%)	-0.64	0 100 100	95, 125, 157, 178	0
2	B	85/86 (98%)	-0.78	0 100 100	92, 107, 134, 151	0
All	All	110/114 (96%)	-0.75	0 100 100	92, 110, 147, 178	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CCC	B	86	23/24	0.90	0.06	128,163,178,189	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MG	B	118	1/1	0.80	0.10	169,169,169,169	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SR	B	102	1/1	0.81	0.08	213,213,213,213	0
6	MG	B	116	1/1	0.84	0.14	149,149,149,149	0
6	MG	B	117	1/1	0.84	0.20	160,160,160,160	0
3	SR	B	103	1/1	0.84	0.05	228,228,228,228	0
4	NCO	B	105	7/7	0.85	0.07	188,218,224,224	0
3	SR	A	101	1/1	0.89	0.07	207,207,207,207	0
4	NCO	A	104	7/7	0.91	0.11	193,207,220,226	0
3	SR	A	102	1/1	0.93	0.12	215,215,215,215	0
3	SR	A	103	1/1	0.93	0.17	201,201,201,201	0
5	SPD	B	115	10/10	0.93	0.23	93,109,113,113	10
4	NCO	B	106	7/7	0.94	0.14	97,131,150,155	0
4	NCO	B	109	7/7	0.95	0.08	103,119,134,137	0
4	NCO	B	113	7/7	0.95	0.07	140,146,162,166	0
3	SR	B	101	1/1	0.95	0.05	177,177,177,177	0
4	NCO	B	111	7/7	0.96	0.06	120,128,140,154	0
4	NCO	B	114	7/7	0.96	0.08	124,137,152,162	0
4	NCO	B	107	7/7	0.97	0.08	118,123,130,133	0
4	NCO	B	108	7/7	0.97	0.07	120,127,129,133	0
4	NCO	B	112	7/7	0.97	0.07	120,132,144,155	0
4	NCO	B	104	7/7	0.98	0.07	96,111,123,128	0
4	NCO	B	110	7/7	0.98	0.05	117,132,139,145	0

6.5 Other polymers [i](#)

There are no such residues in this entry.