



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:39 PM UTC

PDB ID : 5JOC / pdb_00005joc
Title : Crystal structure of the S61A mutant of AmpC BER
Authors : Na, J.H.; An, Y.J.; Cha, S.S.
Deposited on : 2016-05-02
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	366	
1	B	366	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5790 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	349	Total	C	N	O	S	0	0	0
			2732	1762	468	495	7			
1	B	350	Total	C	N	O	S	0	0	0
			2737	1765	469	496	7			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	expression tag	UNP A7TUE6
A	-4	HIS	-	expression tag	UNP A7TUE6
A	-3	HIS	-	expression tag	UNP A7TUE6
A	-2	HIS	-	expression tag	UNP A7TUE6
A	-1	HIS	-	expression tag	UNP A7TUE6
A	0	HIS	-	expression tag	UNP A7TUE6
A	1	HIS	-	expression tag	UNP A7TUE6
A	61	ALA	SER	engineered mutation	UNP A7TUE6
B	-5	MET	-	expression tag	UNP A7TUE6
B	-4	HIS	-	expression tag	UNP A7TUE6
B	-3	HIS	-	expression tag	UNP A7TUE6
B	-2	HIS	-	expression tag	UNP A7TUE6
B	-1	HIS	-	expression tag	UNP A7TUE6
B	0	HIS	-	expression tag	UNP A7TUE6
B	1	HIS	-	expression tag	UNP A7TUE6
B	61	ALA	SER	engineered mutation	UNP A7TUE6

- Molecule 2 is CITRIC ACID (CCD ID: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		
2	A	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		
2	B	1	Total	C	O	0	0
			13	6	7		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	134	Total	O	0	0
			134	134		
3	B	135	Total	O	0	0
			135	135		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-lactamase



• Molecule 1: Beta-lactamase



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	207.21 Å 207.21 Å 45.63 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.75 50.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	96.5 (50.00-1.75) 96.5 (50.00-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 1.75 Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.177 , 0.200 0.186 , 0.204	Depositor DCC
R_{free} test set	5514 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.250	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 20.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.028 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5790	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.47	15/2813 (0.5%)	1.37	13/3843 (0.3%)
1	B	1.44	9/2818 (0.3%)	1.22	6/3850 (0.2%)
All	All	1.45	24/5631 (0.4%)	1.29	19/7693 (0.2%)

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	202	GLU	N-CA	10.64	1.62	1.46
1	A	201	ARG	CA-C	-6.76	1.46	1.52
1	A	214	ASP	N-CA	-6.37	1.38	1.46
1	A	343	TYR	N-CA	6.35	1.51	1.45
1	A	236	ASN	CA-C	-6.08	1.45	1.52
1	A	75	ILE	N-CA	5.83	1.53	1.46
1	A	57	PHE	N-CA	-5.82	1.38	1.45
1	B	220	VAL	CA-C	5.82	1.59	1.52
1	A	202	GLU	CA-C	-5.76	1.45	1.53
1	A	220	VAL	CA-C	5.70	1.59	1.52
1	A	29	VAL	C-O	5.63	1.30	1.24
1	A	201	ARG	C-N	-5.56	1.26	1.33
1	A	60	GLY	CA-C	5.54	1.59	1.51
1	B	256	TYR	N-CA	-5.51	1.39	1.46
1	A	324	TYR	CA-C	-5.45	1.45	1.52
1	B	343	TYR	CA-C	5.43	1.57	1.52
1	B	159	ALA	CA-C	-5.36	1.45	1.52
1	B	189	PRO	CA-C	5.36	1.57	1.52
1	B	300	ILE	C-O	5.30	1.29	1.24
1	B	188	VAL	C-O	-5.22	1.17	1.24
1	A	276	ASN	N-CA	5.18	1.52	1.46
1	A	331	LYS	CA-C	-5.12	1.45	1.52
1	B	127	ASP	CA-C	5.08	1.59	1.52
1	B	229	ARG	N-CA	-5.03	1.40	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	201	ARG	CA-C-N	-23.34	91.00	123.20
1	A	201	ARG	C-N-CA	-23.34	91.00	123.20
1	A	201	ARG	O-C-N	-11.83	108.11	121.95
1	A	276	ASN	CB-CA-C	-8.03	96.26	110.47
1	A	241	ASN	N-CA-C	8.02	120.94	111.71
1	B	188	VAL	CA-C-N	-7.90	112.25	120.38
1	B	188	VAL	C-N-CA	-7.90	112.25	120.38
1	B	241	ASN	N-CA-C	6.88	118.43	111.07
1	A	188	VAL	CA-C-N	-6.88	113.30	120.38
1	A	188	VAL	C-N-CA	-6.88	113.30	120.38
1	A	250	GLN	CB-CA-C	-6.53	100.63	110.88
1	B	12	THR	OG1-CB-CG2	6.18	121.66	109.30
1	B	240	ILE	N-CA-C	6.13	117.41	108.58
1	A	190	PRO	N-CA-C	5.49	117.39	110.70
1	A	360	GLN	CA-C-O	-5.44	111.56	120.80
1	A	254	SER	N-CA-C	-5.28	103.06	110.50
1	A	243	LYS	N-CA-C	5.09	116.83	111.28
1	B	221	LYS	CB-CG-CD	5.07	122.97	111.30
1	A	297	VAL	O-C-N	5.07	128.25	122.98

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2732	0	2700	16	0
1	B	2737	0	2705	17	0
2	A	26	0	10	0	0
2	B	26	0	10	1	0
3	A	134	0	0	0	0
3	B	135	0	0	0	0
All	All	5790	0	5425	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:8:ILE:O	1:B:12:THR:HG23	1.84	0.76
1:A:8:ILE:O	1:A:12:THR:HG23	1.95	0.67
1:A:180:LYS:O	1:A:229:ARG:NH1	2.27	0.67
1:B:235:MET:HG2	1:B:327:PHE:CD1	2.30	0.66
1:A:237:PRO:HA	1:A:240:ILE:HD12	1.82	0.62
2:B:401:CIT:O4	2:B:401:CIT:O7	2.10	0.58
1:A:12:THR:HB	1:A:354:GLN:HE21	1.70	0.57
1:A:235:MET:HG2	1:A:327:PHE:CD2	2.40	0.56
1:A:12:THR:HB	1:A:354:GLN:NE2	2.21	0.55
1:B:236:ASN:HB3	1:B:239:ASP:OD2	2.09	0.52
1:B:259:THR:CG2	1:B:262:MET:CE	2.89	0.50
1:B:161:LYS:HB2	1:B:162:PRO:HD3	1.93	0.50
1:B:259:THR:HG22	1:B:262:MET:CE	2.43	0.49
1:A:161:LYS:HB2	1:A:162:PRO:HD3	1.96	0.48
1:A:275:VAL:CG2	1:A:279:SER:HB2	2.43	0.48
1:B:12:THR:HB	1:B:354:GLN:HE21	1.79	0.47
1:A:3:GLN:NE2	1:A:3:GLN:HA	2.30	0.46
1:B:237:PRO:HA	1:B:240:ILE:HD12	1.98	0.46
1:B:298:LYS:HE2	1:B:298:LYS:HB3	1.77	0.45
1:A:8:ILE:O	1:A:12:THR:CG2	2.63	0.45
1:B:273:TRP:CD2	1:B:274:PRO:HA	2.52	0.45
1:A:305:PRO:O	1:A:306:ALA:C	2.60	0.44
1:A:105:HIS:CE1	1:A:141:PRO:HB2	2.53	0.44
1:B:259:THR:HG22	1:B:262:MET:HE1	1.99	0.44
1:A:280:ILE:HD13	1:A:311:TRP:CE2	2.53	0.44
1:B:12:THR:HB	1:B:354:GLN:NE2	2.34	0.42
1:A:357:ASN:HD22	1:A:360:GLN:HE21	1.68	0.42
1:B:277:PRO:O	1:B:281:ILE:HD12	2.19	0.42
1:B:315:THR:HG22	1:B:324:TYR:CD1	2.54	0.42
1:B:259:THR:CG2	1:B:262:MET:HE1	2.49	0.41
1:B:262:MET:HE3	1:B:262:MET:HB3	1.64	0.41
1:A:273:TRP:CD2	1:A:274:PRO:HA	2.55	0.41
1:A:14:THR:HB	1:A:15:PRO:HD3	2.03	0.41
1:B:305:PRO:O	1:B:306:ALA:C	2.63	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/366 (94%)	335 (97%)	10 (3%)	0	100	100
1	B	346/366 (94%)	339 (98%)	6 (2%)	1 (0%)	36	21
All	All	691/732 (94%)	674 (98%)	16 (2%)	1 (0%)	48	32

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	3	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	283/296 (96%)	274 (97%)	9 (3%)	34	14
1	B	283/296 (96%)	277 (98%)	6 (2%)	47	27
All	All	566/592 (96%)	551 (97%)	15 (3%)	39	19

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	THR
1	A	93	LEU
1	A	121	GLU
1	A	125	SER
1	A	238	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	243	LYS
1	A	250	GLN
1	A	262	MET
1	A	278	ASP
1	B	12	THR
1	B	96	LYS
1	B	128	LEU
1	B	243	LYS
1	B	262	MET
1	B	282	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	19	GLN
1	A	20	GLN
1	A	177	GLN
1	A	187	ASN
1	A	236	ASN
1	A	250	GLN
1	A	276	ASN
1	A	354	GLN
1	A	357	ASN
1	B	3	GLN
1	B	49	GLN
1	B	169	GLN
1	B	172	GLN
1	B	282	ASN
1	B	354	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CIT	B	401	-	12,12,12	1.70	3 (25%)	17,17,17	2.30	8 (47%)
2	CIT	B	400	-	12,12,12	1.51	3 (25%)	17,17,17	1.82	5 (29%)
2	CIT	A	401	-	12,12,12	2.11	2 (16%)	17,17,17	2.18	4 (23%)
2	CIT	A	400	-	12,12,12	1.20	1 (8%)	17,17,17	1.53	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CIT	B	401	-	-	6/16/16/16	-
2	CIT	B	400	-	-	1/16/16/16	-
2	CIT	A	401	-	-	8/16/16/16	-
2	CIT	A	400	-	-	5/16/16/16	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	CIT	C3-C6	5.04	1.58	1.53
2	B	401	CIT	C3-C6	3.83	1.57	1.53
2	A	401	CIT	C2-C3	3.31	1.58	1.54
2	B	400	CIT	O3-C5	3.04	1.32	1.22
2	B	400	CIT	C4-C3	2.77	1.57	1.54
2	B	401	CIT	O1-C1	2.25	1.29	1.22

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	CIT	O3-C5	2.17	1.29	1.22
2	B	400	CIT	O5-C6	2.16	1.28	1.22
2	A	400	CIT	O3-C5	2.02	1.28	1.22

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	CIT	O6-C6-C3	6.85	126.29	113.14
2	A	400	CIT	C4-C3-C2	4.18	120.03	109.31
2	B	401	CIT	C4-C3-C6	3.98	118.84	110.03
2	B	401	CIT	C3-C4-C5	-3.67	103.90	113.92
2	B	400	CIT	C4-C3-C2	3.59	118.53	109.31
2	B	401	CIT	O1-C1-C2	-3.59	112.79	122.95
2	A	401	CIT	O6-C6-O5	-3.40	112.98	123.86
2	B	401	CIT	O6-C6-C3	3.16	119.21	113.14
2	B	401	CIT	O6-C6-O5	-2.91	114.54	123.86
2	B	400	CIT	O1-C1-C2	-2.86	114.86	122.95
2	A	400	CIT	C4-C3-C6	-2.79	103.87	110.03
2	B	400	CIT	O7-C3-C2	-2.38	103.95	109.38
2	A	401	CIT	C3-C4-C5	-2.34	107.54	113.92
2	A	401	CIT	O4-C5-O3	-2.29	117.45	123.33
2	B	401	CIT	O2-C1-C2	2.28	121.58	114.35
2	B	400	CIT	O4-C5-O3	2.18	128.94	123.33
2	B	401	CIT	O7-C3-C2	-2.18	104.41	109.38
2	B	401	CIT	O5-C6-C3	2.13	126.21	122.09
2	A	400	CIT	O3-C5-C4	-2.02	117.24	122.95
2	B	400	CIT	O5-C6-C3	-2.00	118.21	122.09

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	CIT	O7-C3-C6-O5
2	A	401	CIT	O7-C3-C6-O6
2	A	401	CIT	C4-C3-C6-O5
2	A	401	CIT	C4-C3-C6-O6
2	B	401	CIT	O7-C3-C6-O5
2	B	401	CIT	O7-C3-C6-O6
2	B	401	CIT	C4-C3-C6-O5
2	B	401	CIT	C4-C3-C6-O6
2	A	400	CIT	C4-C3-C6-O6
2	A	400	CIT	O7-C3-C4-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	400	CIT	O7-C3-C4-C5
2	A	401	CIT	C1-C2-C3-O7
2	A	400	CIT	C4-C3-C6-O5
2	A	401	CIT	C2-C3-C6-O6
2	B	401	CIT	C1-C2-C3-O7
2	A	400	CIT	C2-C3-C6-O5
2	A	401	CIT	C2-C3-C6-O5
2	A	400	CIT	C2-C3-C6-O6
2	A	401	CIT	O1-C1-C2-C3
2	B	401	CIT	C3-C4-C5-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	CIT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	349/366 (95%)	0.03	12 (3%) 48 54	17, 23, 44, 71	0
1	B	350/366 (95%)	0.09	15 (4%) 40 46	17, 24, 47, 81	0
All	All	699/732 (95%)	0.06	27 (3%) 43 49	17, 24, 46, 81	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	292	ALA	8.4
1	B	2	PRO	5.5
1	A	2	PRO	5.3
1	B	238	ARG	4.5
1	B	293	ALA	4.2
1	B	282	ASN	4.1
1	B	294	ALA	4.0
1	A	281	ILE	4.0
1	B	33	GLY	3.7
1	A	282	ASN	3.6
1	B	281	ILE	3.3
1	A	33	GLY	3.2
1	B	239	ASP	3.2
1	B	3	GLN	3.1
1	A	293	ALA	3.1
1	A	4	GLN	2.6
1	B	240	ILE	2.5
1	A	238	ARG	2.5
1	B	237	PRO	2.4
1	B	278	ASP	2.3
1	A	34	LYS	2.3
1	A	360	GLN	2.3
1	A	240	ILE	2.3
1	A	229	ARG	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	11	ARG	2.1
1	A	241	ASN	2.0
1	B	120	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CIT	B	401	13/13	0.88	0.13	27,47,59,73	0
2	CIT	A	401	13/13	0.90	0.14	24,41,63,69	0
2	CIT	A	400	13/13	0.94	0.08	23,28,31,32	0
2	CIT	B	400	13/13	0.96	0.07	21,31,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.