



wwPDB EM Validation Summary Report ⓘ

Mar 10, 2026 – 09:08 AM UTC

PDB ID : 6JP5 / pdb_00006jp5
EMDB ID : EMD-9866
Title : Rabbit Cav1.1-Nifedipine Complex
Authors : Zhao, Y.; Huang, G.; Wu, J.; Yan, N.
Deposited on : 2019-03-25
Resolution : 2.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY




The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Q-score	-	25397	13054 (2.40 - 3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1873	
2	E	222	
3	F	1073	
4	B	450	
4	C	450	
5	D	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	G	2	
5	J	2	
5	K	2	
5	L	2	
6	H	3	
6	I	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	D	1	-	-	X	-
5	NAG	L	1	-	-	X	-
7	NAG	F	1125	-	-	X	-

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 22068 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Voltage-dependent L-type calcium channel subunit alpha-1S.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1274	Total	C	N	O	S	0	0
			10219	6728	1668	1753	70		

- Molecule 2 is a protein called Voltage-dependent calcium channel gamma-1 subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	E	169	Total	C	N	O	S	0	0
			1326	872	216	220	18		

- Molecule 3 is a protein called Voltage-dependent calcium channel subunit alpha-2/delta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	F	973	Total	C	N	O	S	1	0
			7804	4942	1320	1510	32		

- Molecule 4 is a protein called Voltage-dependent L-type calcium channel subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	B	100	Total	C	N	O	S	0	0
			710	455	125	129	1		
4	C	178	Total	C	N	O	S	0	0
			1367	876	232	254	5		

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	75	GLN	-	expression tag	UNP P19517
B	76	GLY	-	expression tag	UNP P19517
B	77	PRO	-	expression tag	UNP P19517
B	78	HIS	-	expression tag	UNP P19517
B	79	MET	-	expression tag	UNP P19517
C	75	GLN	-	expression tag	UNP P19517
C	76	GLY	-	expression tag	UNP P19517

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	77	PRO	-	expression tag	UNP P19517
C	78	HIS	-	expression tag	UNP P19517
C	79	MET	-	expression tag	UNP P19517

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
5	D	2	Total	C	N	O	0	0
			28	16	2	10		
5	G	2	Total	C	N	O	0	0
			28	16	2	10		
5	J	2	Total	C	N	O	0	0
			28	16	2	10		
5	K	2	Total	C	N	O	0	0
			28	16	2	10		
5	L	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
6	H	3	Total	C	N	O	0	0
			39	22	2	15		
6	I	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				AltConf
7	A	1	Total	C	N	O	0
			14	8	1	5	
7	F	1	Total	C	N	O	0
			14	8	1	5	
7	F	1	Total	C	N	O	0
			14	8	1	5	
7	F	1	Total	C	N	O	0
			14	8	1	5	
7	F	1	Total	C	N	O	0
			14	8	1	5	
7	F	1	Total	C	N	O	0
			14	8	1	5	
7	F	1	Total	C	N	O	0
			14	8	1	5	
7	F	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 8 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		AltConf
8	A	3	Total	Ca	0
			3	3	

- Molecule 9 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (CCD ID: 3PE) (formula: C₄₁H₈₂NO₈P).




Mol	Chain	Residues	Atoms					AltConf
9	A	1	Total	C	N	O	P	0
			33	23	1	8	1	
9	A	1	Total	C	N	O	P	0
			30	20	1	8	1	
9	A	1	Total	C	N	O	P	0
			32	22	1	8	1	
9	A	1	Total	C	N	O	P	0
			19	11	1	6	1	
9	A	1	Total	C	N	O	P	0
			30	20	1	8	1	
9	A	1	Total	C	N	O	P	0
			30	20	1	8	1	
9	A	1	Total	C	N	O	P	0
			21	13	1	6	1	
9	A	1	Total	C	N	O	P	0
			32	22	1	8	1	

- Molecule 10 is dimethyl 2,6-dimethyl-4-(2-nitrophenyl)-1,4-dihydropyridine-3,5-dicarboxylate (CCD ID: C5U) (formula: C₁₇H₁₈N₂O₆).




Mol	Chain	Residues	Atoms				AltConf
10	A	1	Total	C	N	O	0
			25	17	2	6	

- Molecule 11 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PC1) (formula: $C_{44}H_{88}NO_8P$).



Mol	Chain	Residues	Atoms					AltConf
11	A	1	Total	C	N	O	P	0
			39	29	1	8	1	

- Molecule 12 is ETHANOLAMINE (CCD ID: ETA) (formula: C_2H_7NO).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
12	F	1	4	2	1	1	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Voltage-dependent L-type calcium channel subunit alpha-1S





[illegible]

- Molecule 2: Voltage-dependent calcium channel gamma-1 subunit




- Molecule 3: Voltage-dependent calcium channel subunit alpha-2/delta-1






• Molecule 4: Voltage-dependent L-type calcium channel subunit beta-1



- Molecule 4: Voltage-dependent L-type calcium channel subunit beta-1



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-3)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	245171	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	48	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.201	Depositor
Minimum map value	-0.107	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.023	Depositor
Map size ($\text{\AA}$)	349.12, 349.12, 349.12	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.091, 1.091, 1.091	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ETA, NAG, PC1, C5U, CA, BMA, 3PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.73	0/10463	0.96	37/14191 (0.3%)
2	E	0.49	0/1358	0.81	2/1832 (0.1%)
3	F	0.81	1/7974 (0.0%)	1.01	22/10816 (0.2%)
4	B	0.18	0/723	0.37	0/979
4	C	0.26	0/1394	0.49	1/1892 (0.1%)
All	All	0.72	1/21912 (0.0%)	0.93	62/29710 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	665	ALA	C-N	-5.45	1.21	1.33

The worst 5 of 62 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	478	LYS	N-CA-C	9.73	121.89	111.28
1	A	493	MET	N-CA-C	9.07	121.17	111.28
3	F	869	ASP	N-CA-C	-8.10	103.40	113.28
3	F	991	LYS	N-CA-C	8.02	120.02	111.28
3	F	325	ASN	N-CA-C	7.95	120.85	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10219	0	10303	577	0
2	E	1326	0	1345	150	0
3	F	7804	0	7608	530	0
4	B	710	0	633	7	0
4	C	1367	0	1343	65	0
5	D	28	0	25	8	0
5	G	28	0	25	1	0
5	J	28	0	25	0	0
5	K	28	0	25	7	0
5	L	28	0	25	11	0
6	H	39	0	34	5	0
6	I	39	0	34	1	0
7	A	14	0	13	0	0
7	F	112	0	104	24	0
8	A	3	0	0	0	0
9	A	227	0	260	46	0
10	A	25	0	0	8	0
11	A	39	0	52	8	0
12	F	4	0	0	2	0
All	All	22068	0	21854	1256	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 1256 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:F:784:ASN:ND2	7:F:1120:NAG:C1	1.68	1.55
3:F:678:ASN:HD21	5:K:1:NAG:C1	1.19	1.52
3:F:326:ASN:ND2	7:F:1104:NAG:C1	1.72	1.50
3:F:988:ASN:HD21	7:F:1122:NAG:C1	1.18	1.50
3:F:1001:ASN:HD21	7:F:1121:NAG:C1	1.27	1.44

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

16 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	D	1	5	14,14,15	0.28	0	17,19,21	0.99	1 (5%)
5	NAG	D	2	5	14,14,15	0.31	0	17,19,21	0.50	0
5	NAG	G	1	3,5	14,14,15	0.30	0	17,19,21	0.56	0
5	NAG	G	2	5	14,14,15	0.28	0	17,19,21	0.55	0
6	NAG	H	1	3,6	14,14,15	0.69	1 (7%)	17,19,21	0.70	0
6	NAG	H	2	6	14,14,15	0.37	0	17,19,21	0.80	0
6	BMA	H	3	6	11,11,12	0.73	0	15,15,17	0.94	1 (6%)
6	NAG	I	1	3,6	14,14,15	0.80	1 (7%)	17,19,21	0.84	0
6	NAG	I	2	6	14,14,15	1.54	1 (7%)	17,19,21	0.93	1 (5%)
6	BMA	I	3	6	11,11,12	0.90	0	15,15,17	0.92	0
5	NAG	J	1	3,5	14,14,15	0.52	0	17,19,21	0.64	0
5	NAG	J	2	5	14,14,15	0.31	0	17,19,21	0.40	0
5	NAG	K	1	5	14,14,15	0.28	0	17,19,21	0.62	0
5	NAG	K	2	5	14,14,15	0.29	0	17,19,21	0.51	0
5	NAG	L	1	5	14,14,15	0.31	0	17,19,21	0.65	0
5	NAG	L	2	5	14,14,15	0.27	0	17,19,21	1.32	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	D	1	5	-	5/6/23/26	0/1/1/1
5	NAG	D	2	5	-	2/6/23/26	0/1/1/1
5	NAG	G	1	3,5	-	0/6/23/26	0/1/1/1
5	NAG	G	2	5	-	2/6/23/26	0/1/1/1
6	NAG	H	1	3,6	-	0/6/23/26	0/1/1/1
6	NAG	H	2	6	-	2/6/23/26	0/1/1/1
6	BMA	H	3	6	-	0/2/19/22	0/1/1/1
6	NAG	I	1	3,6	-	2/6/23/26	0/1/1/1
6	NAG	I	2	6	-	2/6/23/26	0/1/1/1
6	BMA	I	3	6	-	2/2/19/22	0/1/1/1
5	NAG	J	1	3,5	-	2/6/23/26	0/1/1/1
5	NAG	J	2	5	-	0/6/23/26	0/1/1/1
5	NAG	K	1	5	-	4/6/23/26	0/1/1/1
5	NAG	K	2	5	-	6/6/23/26	0/1/1/1
5	NAG	L	1	5	-	5/6/23/26	0/1/1/1
5	NAG	L	2	5	-	4/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	I	2	NAG	O5-C1	-5.58	1.34	1.43
6	I	1	NAG	O5-C1	-2.50	1.39	1.43
6	H	1	NAG	O5-C1	-2.23	1.40	1.43

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	L	2	NAG	C1-O5-C5	4.37	118.04	112.19
5	D	1	NAG	C3-C4-C5	2.50	114.76	110.23
6	H	3	BMA	C1-O5-C5	2.48	115.51	112.19
6	I	2	NAG	C4-C3-C2	2.29	114.38	111.02

There are no chirality outliers.

5 of 38 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	D	1	NAG	C1-C2-N2-C7
5	D	1	NAG	C8-C7-N2-C2
5	D	1	NAG	O7-C7-N2-C2
5	K	1	NAG	C8-C7-N2-C2
5	K	1	NAG	O7-C7-N2-C2


There are no ring outliers.


10 monomers are involved in 33 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	L	1	NAG	7	0
5	G	2	NAG	1	0
5	K	2	NAG	5	0
5	K	1	NAG	4	0
5	D	2	NAG	4	0
5	D	1	NAG	8	0
5	L	2	NAG	4	0
5	G	1	NAG	1	0
6	I	1	NAG	1	0
6	H	1	NAG	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.















5.6 Ligand geometry

Of 23 ligands modelled in this entry, 3 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
7	NAG	F	1115	3	14,14,15	0.85	1 (7%)	17,19,21	1.02	1 (5%)
7	NAG	F	1104	-	14,14,15	0.30	0	17,19,21	0.64	0
7	NAG	A	1901	1	14,14,15	0.31	0	17,19,21	0.49	0
7	NAG	F	1125	3	14,14,15	0.28	0	17,19,21	0.59	0
9	3PE	A	1906	-	31,31,50	1.16	2 (6%)	34,36,55	1.04	2 (5%)
9	3PE	A	1904	-	32,32,50	1.14	2 (6%)	35,37,55	1.30	4 (11%)
9	3PE	A	1905	-	29,29,50	1.20	2 (6%)	32,34,55	1.14	2 (6%)
9	3PE	A	1909	-	29,29,50	1.18	2 (6%)	32,34,55	1.10	2 (6%)
10	C5U	A	1911	-	26,26,26	4.61	11 (42%)	34,37,37	3.13	13 (38%)
9	3PE	A	1913	-	31,31,50	1.18	2 (6%)	34,36,55	1.12	2 (5%)
7	NAG	F	1121	-	14,14,15	0.47	0	17,19,21	0.53	0
12	ETA	F	1101	-	3,3,3	0.41	0	2,2,2	0.49	0
7	NAG	F	1120	-	14,14,15	0.29	0	17,19,21	0.55	0
7	NAG	F	1122	-	14,14,15	0.28	0	17,19,21	0.57	0
11	PC1	A	1912	-	38,38,53	1.08	2 (5%)	44,46,61	1.06	3 (6%)
7	NAG	F	1107	3	14,14,15	0.30	0	17,19,21	0.57	0
7	NAG	F	1114	3	14,14,15	0.25	0	17,19,21	0.67	1 (5%)
9	3PE	A	1910	-	20,20,50	1.03	1 (5%)	22,23,55	0.99	1 (4%)
9	3PE	A	1907	-	18,18,50	1.11	1 (5%)	20,21,55	0.98	1 (5%)
9	3PE	A	1908	-	29,29,50	1.19	2 (6%)	32,34,55	1.13	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	F	1115	3	-	0/6/23/26	0/1/1/1
7	NAG	F	1104	-	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	1901	1	-	1/6/23/26	0/1/1/1
7	NAG	F	1125	3	-	2/6/23/26	0/1/1/1
9	3PE	A	1906	-	-	7/35/35/54	-
9	3PE	A	1904	-	-	7/36/36/54	-
9	3PE	A	1905	-	-	2/33/33/54	-
9	3PE	A	1909	-	-	3/33/33/54	-
10	C5U	A	1911	-	-	8/18/40/40	0/2/2/2
9	3PE	A	1913	-	-	9/35/35/54	-
7	NAG	F	1121	-	-	0/6/23/26	0/1/1/1
12	ETA	F	1101	-	-	1/1/1/1	-
7	NAG	F	1120	-	-	2/6/23/26	0/1/1/1
7	NAG	F	1122	-	-	2/6/23/26	0/1/1/1
11	PC1	A	1912	-	-	12/42/42/57	-
7	NAG	F	1107	3	-	2/6/23/26	0/1/1/1
7	NAG	F	1114	3	-	0/6/23/26	0/1/1/1
9	3PE	A	1910	-	-	2/21/21/54	-
9	3PE	A	1907	-	-	1/19/19/54	-
9	3PE	A	1908	-	-	8/33/33/54	-

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	A	1911	C5U	C06-C05	14.63	1.52	1.35
10	A	1911	C5U	C08-C09	14.53	1.52	1.35
10	A	1911	C5U	C06-N07	4.86	1.44	1.38
10	A	1911	C5U	C08-N07	4.44	1.44	1.38
9	A	1907	3PE	O31-C31	4.28	1.45	1.33

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	1911	C5U	C25-C06-C05	-9.20	118.46	127.61
10	A	1911	C5U	C24-C08-C09	-7.71	119.95	127.61
10	A	1911	C5U	C25-C06-N07	5.17	119.44	113.42
10	A	1911	C5U	C24-C08-N07	5.17	119.43	113.42
10	A	1911	C5U	C08-N07-C06	-4.66	119.10	123.44

There are no chirality outliers.

5 of 71 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
7	A	1901	NAG	C3-C2-N2-C7
7	F	1104	NAG	C8-C7-N2-C2
7	F	1104	NAG	O7-C7-N2-C2
7	F	1125	NAG	C8-C7-N2-C2
7	F	1125	NAG	O7-C7-N2-C2



There are no ring outliers.


18 monomers are involved in 87 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	F	1115	NAG	2	0
7	F	1104	NAG	3	0
7	F	1125	NAG	9	0
9	A	1906	3PE	3	0
9	A	1904	3PE	9	0
9	A	1905	3PE	7	0
9	A	1909	3PE	4	0
10	A	1911	C5U	8	0
9	A	1913	3PE	7	0
7	F	1121	NAG	3	0
12	F	1101	ETA	2	0
7	F	1120	NAG	2	0
7	F	1122	NAG	4	0
11	A	1912	PC1	8	0
7	F	1107	NAG	1	0
9	A	1910	3PE	3	0
9	A	1907	3PE	1	0
9	A	1908	3PE	12	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-9866. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 163



Y Index: 151




Z Index: 207


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views

This section was not generated.


6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 109 nm^3 ; this corresponds to an approximate mass of 98 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.345 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-9866 and PDB model 6JP5. Per-residue inclusion information can be found in section [3](#) on page [10](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.023 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.023).

9.4 Atom inclusion [i](#)



At the recommended contour level, 63% of all backbone atoms, 58% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.023) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5780	0.3900
A	0.5600	0.3990
B	0.0000	0.0150
C	0.0000	0.0150
D	0.4290	0.3220
E	0.1140	0.1500
F	0.8320	0.5180
G	0.5710	0.3830
H	0.4870	0.3010
I	0.9490	0.5390
J	0.6430	0.4070
K	0.5360	0.3380
L	0.3210	0.1600

