



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:24 PM UTC

PDB ID : 7JPH / pdb_00007jph
Title : Crystal structure of EBOV glycoprotein with modified HR1c and HR2 stalk at 3.2 Å resolution
Authors : Chaudhary, A.; Stanfield, R.L.; Wilson, I.A.; Zhu, J.
Deposited on : 2020-08-08
Resolution : 3.19 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	284	10% 65% 25% 8%
2	B	173	10% 60% 16% 21%
3	E	2	100%
4	C	6	100%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3256 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	262	Total	C	N	O	S	0	0	0
			2050	1307	346	392	5			

- Molecule 2 is a protein called Envelope glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	136	Total	C	N	O	S	0	0	0
			1073	684	184	199	6			

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	577	PRO	THR	engineered mutation	UNP Q05320
B	615	LEU	TRP	engineered mutation	UNP Q05320
B	638	ALA	-	expression tag	UNP Q05320
B	639	SER	-	expression tag	UNP Q05320
B	640	GLY	-	expression tag	UNP Q05320
B	641	TYR	-	expression tag	UNP Q05320
B	642	ILE	-	expression tag	UNP Q05320
B	643	PRO	-	expression tag	UNP Q05320
B	644	GLU	-	expression tag	UNP Q05320
B	645	ALA	-	expression tag	UNP Q05320
B	646	PRO	-	expression tag	UNP Q05320
B	647	ARG	-	expression tag	UNP Q05320
B	648	ASP	-	expression tag	UNP Q05320
B	649	GLY	-	expression tag	UNP Q05320
B	650	GLN	-	expression tag	UNP Q05320
B	651	ALA	-	expression tag	UNP Q05320
B	652	TYR	-	expression tag	UNP Q05320
B	653	VAL	-	expression tag	UNP Q05320
B	654	ARG	-	expression tag	UNP Q05320
B	655	LYS	-	expression tag	UNP Q05320

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	656	ASP	-	expression tag	UNP Q05320
B	657	GLY	-	expression tag	UNP Q05320
B	658	GLU	-	expression tag	UNP Q05320
B	659	TRP	-	expression tag	UNP Q05320
B	660	VAL	-	expression tag	UNP Q05320
B	661	LEU	-	expression tag	UNP Q05320
B	662	LEU	-	expression tag	UNP Q05320
B	663	SER	-	expression tag	UNP Q05320
B	664	THR	-	expression tag	UNP Q05320
B	665	PHE	-	expression tag	UNP Q05320
B	666	LEU	-	expression tag	UNP Q05320
B	667	GLY	-	expression tag	UNP Q05320
B	668	SER	-	expression tag	UNP Q05320
B	669	HIS	-	expression tag	UNP Q05320
B	670	HIS	-	expression tag	UNP Q05320
B	671	HIS	-	expression tag	UNP Q05320
B	672	HIS	-	expression tag	UNP Q05320
B	673	HIS	-	expression tag	UNP Q05320
B	674	HIS	-	expression tag	UNP Q05320

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	P	0	0
			5	4	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Envelope glycoprotein



• Molecule 2: Envelope glycoprotein



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 4: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
BMA3
MAN4
MAN5
MAN6

4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	114.06Å 114.06Å 136.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.44 – 3.19 49.44 – 3.19	Depositor EDS
% Data completeness (in resolution range)	97.7 (49.44-3.19) 97.7 (49.44-3.19)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.22 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.277 , 0.322 0.278 , 0.324	Depositor DCC
R_{free} test set	852 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	82.8	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 71.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.024 for -h,-k,l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	3256	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, PO4, MAN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	6/2069 (0.3%)	1.34	8/2807 (0.3%)
2	B	1.03	1/1099 (0.1%)	1.35	2/1498 (0.1%)
All	All	1.04	7/3168 (0.2%)	1.34	10/4305 (0.2%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	560	GLN	C-O	6.55	1.31	1.24
1	A	69	ASN	C-O	6.07	1.31	1.23
1	A	64	ARG	C-O	-5.78	1.16	1.23
1	A	189	ALA	C-O	5.74	1.30	1.24
1	A	152	ALA	C-O	-5.28	1.17	1.24
1	A	177	ALA	C-O	5.18	1.30	1.23
1	A	68	LEU	C-O	-5.18	1.17	1.24

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	187	PRO	CA-C-O	-6.30	114.01	122.08
1	A	98	ASN	CA-CB-CG	5.94	118.54	112.60
1	A	251	GLN	O-C-N	5.52	127.83	122.09
2	B	522	ASP	CA-CB-CG	5.50	118.10	112.60
2	B	516	HIS	CA-C-O	-5.42	114.58	120.81
1	A	147	CYS	CB-CA-C	-5.33	103.88	112.09
1	A	250	PRO	O-C-N	5.23	127.82	122.18
1	A	266	ARG	CA-C-O	-5.21	114.75	120.69
1	A	254	LEU	CA-C-O	-5.08	115.03	120.42
1	A	73	ASN	CB-CA-C	5.03	118.50	110.09

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2050	0	1955	48	0
2	B	1073	0	1045	27	0
3	E	28	0	25	3	0
4	C	72	0	61	0	0
5	A	14	0	13	0	0
5	B	14	0	13	0	0
6	B	5	0	0	0	0
All	All	3256	0	3112	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:180:VAL:HG12	2:B:562:ALA:HB1	1.54	0.89
1:A:278:ASN:HD22	1:A:476:UNK:C	2.04	0.70
1:A:51:LEU:HB3	2:B:595:GLN:HE22	1.57	0.70
2:B:603:ILE:HG22	2:B:604:LEU:HG	1.77	0.66
1:A:132:PHE:CD1	1:A:133:PRO:HD2	2.33	0.64
1:A:180:VAL:HG13	2:B:566:THR:CG2	2.30	0.61
2:B:529:LEU:HG	2:B:532:ILE:HD12	1.83	0.60
1:A:180:VAL:HG13	2:B:566:THR:HG23	1.86	0.58
1:A:79:VAL:N	1:A:80:PRO:HD2	2.20	0.56
1:A:113:ILE:O	1:A:121:CYS:HB2	2.05	0.56
2:B:584:ILE:O	2:B:588:LYS:HG2	2.07	0.54
2:B:515:LEU:HB2	2:B:555:ILE:HD12	1.90	0.54
1:A:256:LEU:HD12	1:A:256:LEU:O	2.08	0.53
1:A:63:LEU:CD2	1:A:185:ILE:HG12	2.38	0.53
1:A:70:LEU:HD12	1:A:178:GLU:HB3	1.91	0.53
1:A:218:ILE:HG22	1:A:220:TYR:CE1	2.46	0.51
1:A:180:VAL:HG11	2:B:562:ALA:C	2.35	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:PRO:HD3	1:A:171:TYR:CD1	2.44	0.50
1:A:180:VAL:CG1	2:B:562:ALA:C	2.85	0.49
1:A:96:VAL:HG22	1:A:167:SER:HB2	1.94	0.49
1:A:130:ARG:O	1:A:162:TYR:HB3	2.12	0.49
2:B:592:PHE:O	2:B:596:ARG:HD2	2.12	0.48
2:B:560:GLN:HE21	2:B:564:GLU:HG3	1.79	0.48
1:A:237:ASP:OD1	1:A:237:ASP:C	2.57	0.47
1:A:96:VAL:O	2:B:580:ARG:HA	2.14	0.47
1:A:278:ASN:ND2	1:A:476:UNK:C	2.74	0.47
1:A:223:THR:O	1:A:230:THR:HG23	2.15	0.47
2:B:506:ASN:ND2	2:B:511:CYS:HB2	2.30	0.46
2:B:572:PHE:C	2:B:572:PHE:CD1	2.93	0.46
1:A:33:ILE:HG21	1:A:183:PHE:CE2	2.50	0.46
1:A:258:GLU:HG3	1:A:258:GLU:O	2.16	0.46
1:A:267:SER:HB3	1:A:273:LEU:HB3	1.96	0.46
2:B:585:LEU:HD23	2:B:585:LEU:HA	1.83	0.46
1:A:180:VAL:HG22	2:B:566:THR:HG21	1.98	0.46
1:A:129:ILE:HD13	1:A:165:LEU:HD13	1.97	0.45
1:A:114:LYS:O	1:A:144:THR:HA	2.17	0.44
1:A:131:GLY:HA2	1:A:175:THR:O	2.17	0.44
2:B:626:ILE:O	2:B:631:VAL:HG23	2.18	0.44
1:A:261:TYR:CD2	3:E:1:NAG:H5	2.53	0.44
1:A:35:LEU:HD22	2:B:588:LYS:HE3	2.00	0.44
1:A:261:TYR:HA	1:A:266:ARG:HD2	2.00	0.44
1:A:307:SER:HA	1:A:476:UNK:HA	2.00	0.44
3:E:1:NAG:H61	3:E:2:NAG:C1	2.48	0.44
1:A:109:TYR:CD1	1:A:109:TYR:N	2.87	0.43
1:A:120:GLU:HG2	1:A:172:ARG:CZ	2.48	0.43
2:B:540:GLU:OE2	2:B:540:GLU:N	2.52	0.43
1:A:79:VAL:N	1:A:80:PRO:CD	2.81	0.42
1:A:234:PHE:CD1	1:A:234:PHE:C	2.96	0.42
1:A:104:TRP:CG	2:B:516:HIS:HD1	2.37	0.42
1:A:104:TRP:HE1	2:B:518:TRP:HB3	1.83	0.42
1:A:291:TRP:CZ3	2:B:512:ASN:HB2	2.55	0.42
1:A:291:TRP:CE3	2:B:512:ASN:HB2	2.55	0.42
1:A:180:VAL:HG12	2:B:562:ALA:CB	2.37	0.42
1:A:63:LEU:HB3	2:B:585:LEU:HD22	2.02	0.42
1:A:52:VAL:O	1:A:55:ASP:HB2	2.20	0.41
2:B:504:ILE:HD13	2:B:561:LEU:HB2	2.02	0.41
1:A:188:GLN:HA	1:A:191:LYS:HB3	2.03	0.41
1:A:265:LYS:HB3	1:A:275:TRP:CZ2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:270:THR:HG22	1:A:270:THR:O	2.22	0.40
1:A:273:LEU:HD21	1:A:275:TRP:NE1	2.36	0.40
1:A:222:ALA:HA	1:A:231:GLU:O	2.21	0.40
3:E:1:NAG:H62	3:E:2:NAG:C7	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/284 (87%)	234 (94%)	14 (6%)	0	100	100
2	B	134/173 (78%)	122 (91%)	11 (8%)	1 (1%)	18	52
All	All	382/457 (84%)	356 (93%)	25 (6%)	1 (0%)	36	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	523	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/239 (92%)	208 (95%)	11 (5%)	22	55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	115/145 (79%)	109 (95%)	6 (5%)	21	54
All	All	334/384 (87%)	317 (95%)	17 (5%)	21	54

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	41	SER
1	A	65	SER
1	A	69	ASN
1	A	98	ASN
1	A	112	GLU
1	A	129	ILE
1	A	167	SER
1	A	174	THR
1	A	184	LEU
1	A	259	THR
1	A	281	ILE
2	B	519	THR
2	B	529	LEU
2	B	581	THR
2	B	596	ARG
2	B	600	THR
2	B	626	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	HIS
1	A	44	GLN
1	A	107	ASN
1	A	188	GLN
1	A	278	ASN
2	B	508	GLN
2	B	521	GLN
2	B	595	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	1	4,2	14,14,15	1.85	4 (28%)	17,19,21	1.90	4 (23%)
4	NAG	C	2	4	14,14,15	0.87	0	17,19,21	1.76	4 (23%)
4	BMA	C	3	4	11,11,12	1.37	2 (18%)	15,15,17	1.57	4 (26%)
4	MAN	C	4	4	11,11,12	1.62	2 (18%)	15,15,17	1.30	2 (13%)
4	MAN	C	5	4	11,11,12	1.72	3 (27%)	15,15,17	1.49	3 (20%)
4	MAN	C	6	4	11,11,12	1.51	2 (18%)	15,15,17	0.98	0
3	NAG	E	1	3,1	14,14,15	0.65	0	17,19,21	1.69	4 (23%)
3	NAG	E	2	3	14,14,15	3.98	13 (92%)	17,19,21	2.23	9 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	4,2	-	0/6/23/26	0/1/1/1
4	NAG	C	2	4	-	2/6/23/26	0/1/1/1
4	BMA	C	3	4	-	2/2/19/22	0/1/1/1
4	MAN	C	4	4	-	2/2/19/22	0/1/1/1
4	MAN	C	5	4	-	0/2/19/22	0/1/1/1
4	MAN	C	6	4	-	2/2/19/22	0/1/1/1
3	NAG	E	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	E	2	3	-	0/6/23/26	0/1/1/1

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	NAG	C4-C5	6.54	1.67	1.53
3	E	2	NAG	O7-C7	5.93	1.36	1.23
3	E	2	NAG	O5-C5	5.44	1.54	1.43
3	E	2	NAG	C1-C2	4.75	1.58	1.52
3	E	2	NAG	C8-C7	4.60	1.60	1.50
3	E	2	NAG	O4-C4	4.31	1.53	1.43
4	C	3	BMA	O5-C1	3.49	1.49	1.43
4	C	1	NAG	O3-C3	3.20	1.50	1.43
3	E	2	NAG	C4-C3	3.13	1.60	1.52
3	E	2	NAG	C6-C5	2.92	1.61	1.51
3	E	2	NAG	C7-N2	2.84	1.43	1.34
4	C	5	MAN	O5-C1	2.82	1.48	1.43
4	C	1	NAG	O4-C4	2.76	1.49	1.43
4	C	4	MAN	C2-C3	2.73	1.56	1.52
4	C	1	NAG	C1-C2	2.73	1.56	1.52
3	E	2	NAG	C3-C2	2.70	1.58	1.52
4	C	1	NAG	C4-C3	2.70	1.59	1.52
4	C	6	MAN	C2-C3	2.67	1.56	1.52
3	E	2	NAG	C2-N2	2.63	1.50	1.46
3	E	2	NAG	O3-C3	2.63	1.49	1.43
4	C	5	MAN	C2-C3	2.53	1.56	1.52
4	C	3	BMA	O2-C2	2.14	1.47	1.43
3	E	2	NAG	O6-C6	2.10	1.51	1.42
4	C	4	MAN	O2-C2	2.07	1.47	1.43
4	C	6	MAN	O2-C2	2.03	1.47	1.43
4	C	5	MAN	O5-C5	2.03	1.47	1.43

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	2	NAG	C2-N2-C7	-4.24	117.22	122.90
3	E	1	NAG	C3-C4-C5	3.78	117.09	110.23
4	C	1	NAG	O3-C3-C4	3.74	119.19	110.38
4	C	3	BMA	O5-C1-C2	3.65	119.49	110.79
3	E	2	NAG	O5-C1-C2	3.57	116.81	111.29
3	E	2	NAG	O4-C4-C5	3.56	118.09	109.32
3	E	2	NAG	C2-N2-C7	-3.41	118.33	122.90
3	E	2	NAG	C8-C7-N2	-3.40	110.48	116.12
4	C	1	NAG	O5-C1-C2	-3.22	106.30	111.29
4	C	4	MAN	O5-C1-C2	3.04	118.04	110.79
4	C	1	NAG	O4-C4-C3	3.02	117.50	110.38
4	C	2	NAG	C3-C4-C5	2.95	115.58	110.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	5	MAN	O5-C1-C2	2.91	117.73	110.79
4	C	5	MAN	C1-O5-C5	2.80	115.94	112.19
3	E	2	NAG	C1-O5-C5	-2.77	108.48	112.19
4	C	3	BMA	C1-O5-C5	2.76	115.88	112.19
4	C	2	NAG	O3-C3-C2	-2.64	103.92	109.40
3	E	2	NAG	C1-C2-N2	2.58	114.50	110.43
3	E	1	NAG	O4-C4-C3	-2.56	104.34	110.38
4	C	3	BMA	O3-C3-C2	-2.44	105.08	110.05
4	C	5	MAN	C1-C2-C3	-2.38	106.18	109.64
3	E	2	NAG	O5-C5-C6	2.36	112.25	107.66
3	E	1	NAG	O3-C3-C2	2.27	114.12	109.40
3	E	2	NAG	O3-C3-C4	2.27	115.73	110.38
3	E	1	NAG	C4-C3-C2	-2.26	107.70	111.02
4	C	3	BMA	C2-C3-C4	2.17	114.68	110.86
3	E	2	NAG	O7-C7-C8	2.17	125.92	122.05
4	C	4	MAN	C3-C4-C5	2.16	114.16	110.23
4	C	2	NAG	O3-C3-C4	2.05	115.21	110.38
4	C	1	NAG	C6-C5-C4	-2.02	108.07	113.02

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	C	3	BMA	O5-C5-C6-O6
3	E	1	NAG	O5-C5-C6-O6
4	C	3	BMA	C4-C5-C6-O6
3	E	1	NAG	C4-C5-C6-O6
4	C	6	MAN	C4-C5-C6-O6
4	C	4	MAN	O5-C5-C6-O6
4	C	4	MAN	C4-C5-C6-O6
4	C	2	NAG	C4-C5-C6-O6
4	C	2	NAG	O5-C5-C6-O6
4	C	6	MAN	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	1	NAG	3	0
3	E	2	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	701	2	14,14,15	1.01	1 (7%)	17,19,21	1.37	3 (17%)
5	NAG	A	501	1	14,14,15	0.66	0	17,19,21	1.22	2 (11%)
6	PO4	B	702	-	4,4,4	0.87	0	6,6,6	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	701	2	-	2/6/23/26	0/1/1/1
5	NAG	A	501	1	-	1/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	701	NAG	C1-C2	2.92	1.56	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	501	NAG	C2-N2-C7	3.29	127.31	122.90
5	B	701	NAG	C1-O5-C5	3.29	116.59	112.19
5	B	701	NAG	C1-C2-N2	2.56	114.47	110.43
5	B	701	NAG	O4-C4-C5	2.15	114.61	109.32
5	A	501	NAG	O5-C5-C6	2.04	111.63	107.66

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	701	NAG	O5-C5-C6-O6
5	B	701	NAG	C4-C5-C6-O6
5	A	501	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	310:VAL	C	471:UNK	N	8.80

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/284 (90%)	0.92	28 (10%) 10 7	55, 94, 172, 201	0
2	B	136/173 (78%)	0.79	18 (13%) 7 6	52, 82, 204, 234	0
All	All	392/457 (85%)	0.88	46 (11%) 9 6	52, 89, 183, 234	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	309	THR	7.4
1	A	308	PHE	5.7
2	B	627	ILE	5.7
1	A	281	ILE	4.5
2	B	630	PHE	4.5
1	A	283	THR	4.3
1	A	51	LEU	4.3
2	B	623	ILE	4.1
1	A	307	SER	4.0
1	A	310	VAL	3.9
1	A	293	THR	3.5
2	B	631	VAL	3.5
1	A	306	LEU	3.5
1	A	280	GLU	3.3
2	B	636	PRO	3.3
1	A	270	THR	3.2
1	A	214	TYR	3.2
2	B	635	LEU	3.2
2	B	624	ASP	3.2
1	A	278	ASN	3.2
2	B	637	ASP	3.1
1	A	279	PRO	3.0
2	B	525	ALA	2.9
1	A	197	HIS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	527	ILE	2.8
1	A	284	THR	2.7
1	A	48	VAL	2.7
2	B	524	GLY	2.6
1	A	305	GLU	2.6
1	A	288	TRP	2.5
1	A	47	ASP	2.4
2	B	626	ILE	2.4
1	A	247	ARG	2.4
1	A	236	VAL	2.3
1	A	277	VAL	2.3
1	A	239	LEU	2.3
2	B	620	THR	2.3
2	B	629	ASP	2.2
1	A	68	LEU	2.2
2	B	539	ALA	2.1
1	A	282	ASP	2.1
2	B	622	LYS	2.1
1	A	188	GLN	2.1
2	B	526	ALA	2.1
2	B	628	HIS	2.1
1	A	191	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	E	1	14/15	-	-	103,118,126,133	0
3	NAG	E	2	14/15	-	-	50,50,50,50	0
4	NAG	C	2	14/15	0.69	0.22	67,85,95,97	0
4	NAG	C	1	14/15	0.72	0.15	57,64,70,70	0
4	BMA	C	3	11/12	-	-	93,106,112,117	0
4	MAN	C	4	11/12	-	-	107,114,123,128	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MAN	C	5	11/12	-	-	111,129,138,141	0
4	MAN	C	6	11/12	-	-	103,115,125,126	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	501	14/15	0.34	0.19	164,195,221,222	0
5	NAG	B	701	14/15	0.48	0.19	125,170,181,183	0
6	PO4	B	702	5/5	0.76	0.18	160,176,179,197	0

6.5 Other polymers [i](#)

There are no such residues in this entry.