



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 03:18 AM UTC

PDB ID : 8JPZ / pdb_00008jpz
Title : The thermostability mutant Gox_M8 from *Aspergillus niger*
Authors : Tu, T.; Yan, Y.R.
Deposited on : 2023-06-13
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8172 (2.10-2.06)
Clashscore	190562	8714 (2.10-2.06)
Ramachandran outliers	187476	8641 (2.10-2.06)
Sidechain outliers	187428	8642 (2.10-2.06)
RSRZ outliers	180081	8177 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	581	 94% 6%
1	B	581	 95% 5%
2	C	6	 33% 33% 33%
2	D	6	 67% 33%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 10304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucose oxidase (Fragment).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	581	Total	C	N	O	S	0	6	0
			4504	2837	780	872	15			
1	B	581	Total	C	N	O	S	7	5	0
			4492	2833	776	868	15			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	31	VAL	THR	conflict	UNP Q5Q041
A	32	VAL	THR	conflict	UNP Q5Q041
A	68	LYS	ASP	conflict	UNP Q5Q041
A	82	CYS	GLU	conflict	UNP Q5Q041
A	88	ARG	GLN	conflict	UNP Q5Q041
A	94	ALA	SER	conflict	UNP Q5Q041
A	274	PHE	THR	conflict	UNP Q5Q041
A	278	THR	TYR	conflict	UNP Q5Q041
A	313	LYS	ASP	conflict	UNP Q5Q041
A	418	GLU	VAL	conflict	UNP Q5Q041
A	508	HIS	ASN	conflict	UNP Q5Q041
B	31	VAL	THR	conflict	UNP Q5Q041
B	32	VAL	THR	conflict	UNP Q5Q041
B	68	LYS	ASP	conflict	UNP Q5Q041
B	82	CYS	GLU	conflict	UNP Q5Q041
B	88	ARG	GLN	conflict	UNP Q5Q041
B	94	ALA	SER	conflict	UNP Q5Q041
B	274	PHE	THR	conflict	UNP Q5Q041
B	278	THR	TYR	conflict	UNP Q5Q041
B	313	LYS	ASP	conflict	UNP Q5Q041
B	418	GLU	VAL	conflict	UNP Q5Q041
B	508	HIS	ASN	conflict	UNP Q5Q041


- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-

beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	0	0	0
			72	40	2	30			
2	D	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		
4	B	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 5 is OXYGEN MOLECULE (CCD ID: OXY) (formula: O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	O	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total O 2 2	0	0

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N O 14 8 1 5	0	0
6	A	1	Total C N O 14 8 1 5	0	0
6	A	1	Total C N O 14 8 1 5	0	0
6	A	1	Total C N O 14 8 1 5	0	0
6	A	1	Total C N O 14 8 1 5	0	0
6	B	1	Total C N O 14 8 1 5	0	0
6	B	1	Total C N O 14 8 1 5	0	0
6	B	1	Total C N O 14 8 1 5	0	0
6	B	1	Total C N O 14 8 1 5	0	0
6	B	1	Total C N O 14 8 1 5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			8	6	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	430	Total	O	0	0
			430	430		
8	B	432	Total	O	0	0
			432	432		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucose oxidase (Fragment)




- Molecule 1: Glucose oxidase (Fragment)



- Molecule 2: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	126.45Å 126.45Å 193.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	21.18 – 2.08 21.18 – 2.08	Depositor EDS
% Data completeness (in resolution range)	100.0 (21.18-2.08) 79.6 (21.18-2.08)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 2.08Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.166 , 0.204 0.166 , 0.203	Depositor DCC
R_{free} test set	4709 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	36.3	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10304	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, MAN, BMA, NAG, EPE, MPD, OXY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/4609	0.52	0/6280
1	B	0.39	0/4597	0.53	0/6265
All	All	0.38	0/9206	0.53	0/12545

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4504	0	4332	25	1
1	B	4492	0	4328	18	0
2	C	72	0	61	4	0
2	D	72	0	61	2	0
3	A	53	0	31	2	0
3	B	53	0	31	3	0
4	A	15	0	17	1	0
4	B	15	0	17	1	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	70	0	65	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	84	0	78	0	0
7	A	8	0	14	1	0
8	A	430	0	0	3	1
8	B	432	0	0	4	0
All	All	10304	0	9035	47	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:82:CYS:SG	8:B:1074:HOH:O	2.20	0.98
1:A:82:CYS:SG	8:A:1069:HOH:O	2.22	0.97
1:A:131:TRP:HE1	7:A:609:MPD:HM2	1.46	0.80
1:B:105:ASN:HB2	3:B:601:FAD:C5X	2.13	0.79
1:A:105:ASN:HB2	3:A:601:FAD:C5X	2.13	0.78
1:A:9:ASP:O	1:A:12:GLU:HG2	1.90	0.70
1:B:351:THR:O	1:B:355:THR:HG23	1.94	0.67
1:B:105:ASN:HB2	3:B:601:FAD:N5	2.11	0.65
1:A:105:ASN:HB2	3:A:601:FAD:N5	2.12	0.64
2:C:2:NAG:H3	2:C:2:NAG:H83	1.78	0.64
2:C:1:NAG:H62	2:C:2:NAG:H82	1.82	0.61
8:B:914:HOH:O	2:C:2:NAG:H81	2.02	0.60
1:A:491:ASN:HA	4:B:602:EPE:H82	1.86	0.56
1:A:129:TRP:CE2	1:A:568:LYS:HE3	2.41	0.56
8:A:703:HOH:O	2:C:1:NAG:H81	2.06	0.55
4:A:602:EPE:H72	1:B:491:ASN:HD22	1.70	0.55
1:B:449:ASP:OD2	8:B:701:HOH:O	2.18	0.55
1:A:415:THR:HA	1:A:418:GLU:O	2.07	0.54
1:A:13:VAL:C	1:A:16:ARG:HE	2.17	0.52
1:A:255:GLN:H	1:A:255:GLN:CD	2.18	0.51
1:B:304:LYS:NZ	1:B:308:GLU:OE2	2.43	0.51
1:A:312:ILE:HD11	1:A:439:LYS:HA	1.92	0.51
1:A:252:LEU:HD21	1:A:266[B]:GLU:HG3	1.92	0.50
1:A:365:GLU:OE1	8:A:701:HOH:O	2.20	0.48
1:B:191:VAL:HB	1:B:196:VAL:HB	1.94	0.48
1:A:496:ALA:HB1	1:A:500:ALA:HB3	1.97	0.46
1:A:11:LYS:O	1:A:16:ARG:NH2	2.49	0.46
1:B:194:ARG:NH2	1:B:471:ASN:OD1	2.49	0.46
1:B:437:LEU:HD11	1:B:447:ALA:HB2	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:VAL:CA	1:A:16:ARG:HE	2.30	0.45
8:B:743:HOH:O	2:D:1:NAG:H81	2.16	0.45
1:A:452:TYR:OH	1:A:511:PRO:HB3	2.18	0.44
1:A:13:VAL:HA	1:A:16:ARG:HE	1.83	0.44
1:A:270:HIS:CD2	1:B:51[A]:SER:HB2	2.53	0.44
1:A:335[A]:ARG:NH2	2:D:2:NAG:O3	2.44	0.43
1:A:412:PHE:O	1:A:421:PHE:HA	2.18	0.43
1:A:292:SER:N	1:A:293:PRO:HD2	2.33	0.43
1:B:35:ARG:HD2	1:B:35:ARG:HA	1.88	0.42
1:B:342:GLY:O	1:B:416:ALA:HA	2.19	0.42
1:A:112:PRO:HB3	1:A:554:MET:HB2	2.01	0.42
1:B:200:LYS:HD3	1:B:208:HIS:O	2.19	0.42
1:B:124:PHE:O	1:B:523:PRO:HD3	2.19	0.42
1:A:102:THR:O	1:A:223:ARG:HD3	2.20	0.41
1:B:399:ASP:OD1	1:B:403:LYS:HE3	2.20	0.41
1:B:308:GLU:HB3	1:B:309:PRO:HD3	2.02	0.41
1:B:557:HIS:HB3	3:B:601:FAD:C2	2.51	0.41
1:A:14:ALA:O	1:A:16:ARG:NH2	2.54	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:701:HOH:O	8:A:902:HOH:O[6_554]	1.93	0.27
1:A:16:ARG:NH1	1:A:365:GLU:OE2[6_454]	1.98	0.22

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	585/581 (101%)	565 (97%)	19 (3%)	1 (0%)	43 44

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	584/581 (100%)	564 (97%)	20 (3%)	0	100	100
All	All	1169/1162 (101%)	1129 (97%)	39 (3%)	1 (0%)	48	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	256	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	469/463 (101%)	469 (100%)	0	100	100
1	B	468/463 (101%)	466 (100%)	2 (0%)	84	89
All	All	937/926 (101%)	935 (100%)	2 (0%)	88	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	13[A]	VAL
1	B	13[B]	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	234	ASN
1	A	396	ASN
1	B	96	ASN
1	B	140	GLN
1	B	234	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	C	1	2,1	14,14,15	0.69	1 (7%)	17,19,21	0.69	1 (5%)
2	NAG	C	2	2	14,14,15	0.42	0	17,19,21	1.33	2 (11%)
2	BMA	C	3	2	11,11,12	0.96	0	15,15,17	1.01	1 (6%)
2	MAN	C	4	2	11,11,12	0.85	0	15,15,17	1.45	3 (20%)
2	NAG	D	1	2,1	14,14,15	0.28	0	17,19,21	0.57	0
2	NAG	D	2	2	14,14,15	0.22	0	17,19,21	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	4/6/23/26	0/1/1/1
2	BMA	C	3	2	-	2/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	NAG	D	1	2,1	-	3/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1	NAG	O5-C1	-2.33	1.39	1.43

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	MAN	C1-O5-C5	4.43	118.13	112.19
2	C	2	NAG	C2-N2-C7	4.34	128.72	122.90
2	C	4	MAN	O2-C2-C3	-2.26	105.48	110.15
2	C	3	BMA	C3-C4-C5	2.22	114.26	110.23
2	C	4	MAN	C1-C2-C3	2.15	112.77	109.64
2	C	2	NAG	C1-C2-N2	2.09	113.73	110.43
2	C	1	NAG	C1-O5-C5	2.08	114.97	112.19

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	3	BMA	O5-C5-C6-O6
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2
2	D	1	NAG	O7-C7-N2-C2
2	C	3	BMA	C4-C5-C6-O6
2	D	1	NAG	O5-C5-C6-O6
2	C	2	NAG	C3-C2-N2-C7
2	D	2	NAG	C3-C2-N2-C7
2	C	2	NAG	C1-C2-N2-C7
2	D	2	NAG	C1-C2-N2-C7


There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	2	0
2	C	2	NAG	3	0
2	D	2	NAG	1	0
2	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EPE	B	602	-	15,15,15	0.83	1 (6%)	19,20,20	1.75	2 (10%)
5	OXY	B	603	-	1,1,1	0.17	0	-		
6	NAG	B	606	1	14,14,15	0.41	0	17,19,21	0.42	0
7	MPD	A	609	-	7,7,7	0.33	0	9,10,10	0.75	0
6	NAG	B	608	1	14,14,15	1.21	1 (7%)	17,19,21	1.46	2 (11%)
6	NAG	B	605	1	14,14,15	0.34	0	17,19,21	0.99	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	B	609	1	14,14,15	0.51	0	17,19,21	1.07	2 (11%)
6	NAG	B	604	1	14,14,15	0.22	0	17,19,21	1.19	2 (11%)
3	FAD	B	601	-	58,58,58	0.39	0	85,89,89	0.64	2 (2%)
6	NAG	B	607	1	14,14,15	0.54	0	17,19,21	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EPE	B	602	-	-	2/9/19/19	0/1/1/1
6	NAG	B	606	1	-	0/6/23/26	0/1/1/1
7	MPD	A	609	-	-	3/5/5/5	-
6	NAG	B	608	1	-	4/6/23/26	0/1/1/1
6	NAG	B	605	1	-	2/6/23/26	0/1/1/1
6	NAG	B	609	1	-	1/6/23/26	0/1/1/1
6	NAG	B	604	1	-	2/6/23/26	0/1/1/1
3	FAD	B	601	-	-	6/34/50/50	0/6/6/6
6	NAG	B	607	1	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	608	NAG	O5-C1	-4.30	1.36	1.43
4	B	602	EPE	C10-S	2.90	1.81	1.77

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	602	EPE	C5-N4-C3	4.57	118.69	108.84
6	B	608	NAG	C3-C4-C5	4.51	118.41	110.23
4	B	602	EPE	O1S-S-C10	3.24	111.62	106.73
6	B	604	NAG	C1-O5-C5	3.17	116.44	112.19
3	B	601	FAD	C4'-C3'-C2'	3.01	118.58	113.57
6	B	608	NAG	C4-C3-C2	2.98	115.39	111.02
6	B	609	NAG	C1-O5-C5	2.85	116.01	112.19
3	B	601	FAD	C5'-C4'-C3'	-2.73	107.07	112.22
6	B	604	NAG	C1-C2-N2	2.20	113.90	110.43
6	B	609	NAG	C1-C2-N2	2.07	113.70	110.43

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	601	FAD	N10-C1'-C2'-O2'
3	B	601	FAD	PA-O3P-P-O5'
6	B	609	NAG	C1-C2-N2-C7
7	A	609	MPD	O2-C2-C3-C4
7	A	609	MPD	CM-C2-C3-C4
6	B	608	NAG	O5-C5-C6-O6
6	B	608	NAG	C4-C5-C6-O6
4	B	602	EPE	C8-C7-N4-C5
3	B	601	FAD	P-O3P-PA-O1A
6	B	604	NAG	C1-C2-N2-C7
6	B	605	NAG	C1-C2-N2-C7
4	B	602	EPE	N4-C7-C8-O8
6	B	608	NAG	C3-C2-N2-C7
3	B	601	FAD	O4B-C4B-C5B-O5B
6	B	608	NAG	C1-C2-N2-C7
3	B	601	FAD	C2B-C1B-N9A-C8A
6	B	604	NAG	C3-C2-N2-C7
6	B	605	NAG	C3-C2-N2-C7
3	B	601	FAD	P-O3P-PA-O2A
7	A	609	MPD	C1-C2-C3-C4
6	B	607	NAG	C4-C5-C6-O6



There are no ring outliers.



3 monomers are involved in 5 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	602	EPE	1	0
7	A	609	MPD	1	0
3	B	601	FAD	3	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	581/581 (100%)	-0.47	1 (0%) 91 92	16, 36, 51, 88	6 (1%)
1	B	580/581 (99%)	-0.50	1 (0%) 91 92	14, 35, 48, 72	4 (0%)
All	All	1161/1162 (99%)	-0.49	2 (0%) 91 92	14, 36, 49, 88	10 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	580	MET	2.5
1	A	16	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MAN	D	6	11/12	0.58	0.14	65,89,95,100	0
2	MAN	C	6	11/12	0.69	0.13	57,82,89,99	0
2	MAN	D	4	11/12	0.73	0.10	67,73,78,82	0
2	MAN	D	5	11/12	0.80	0.09	76,88,94,95	0
2	MAN	C	4	11/12	0.81	0.11	67,76,81,85	0
2	BMA	D	3	11/12	0.83	0.10	67,75,85,91	0
2	MAN	C	5	11/12	0.83	0.09	66,79,85,90	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BMA	C	3	11/12	0.89	0.08	63,69,80,84	0
2	NAG	D	1	14/15	0.89	0.10	36,42,50,52	0
2	NAG	C	2	14/15	0.90	0.09	40,48,57,63	0
2	NAG	C	1	14/15	0.95	0.07	29,40,46,46	0
2	NAG	D	2	14/15	0.95	0.07	47,51,58,65	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	B	608	14/15	0.71	0.13	71,83,94,102	0
6	NAG	A	607	14/15	0.75	0.12	73,83,99,100	0
6	NAG	B	609	14/15	0.77	0.11	58,72,81,83	0
6	NAG	A	608	14/15	0.83	0.10	66,71,84,87	0
7	MPD	A	609	8/8	0.84	0.18	48,57,63,67	8
6	NAG	B	605	14/15	0.86	0.09	52,61,70,73	0
6	NAG	A	606	14/15	0.86	0.10	49,59,62,63	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NAG	B	607	14/15	0.87	0.08	49,64,69,73	0
6	NAG	B	604	14/15	0.88	0.09	51,55,64,69	0
6	NAG	A	605	14/15	0.88	0.08	47,56,58,67	0
6	NAG	A	604	14/15	0.92	0.07	42,50,59,60	0
6	NAG	B	606	14/15	0.92	0.07	34,48,60,60	0
5	OXY	B	603	2/2	0.95	0.07	37,37,37,43	0
5	OXY	A	603	2/2	0.96	0.06	40,40,40,41	0
4	EPE	B	602	15/15	0.96	0.07	39,46,49,53	0
3	FAD	B	601	53/53	0.97	0.06	24,32,44,47	0
3	FAD	A	601	53/53	0.97	0.06	25,32,43,46	0
4	EPE	A	602	15/15	0.98	0.05	41,44,51,52	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NAG B 608:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAG A 607:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAG B 609:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAG A 608:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MPD A 609:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAG B 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAG A 606:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAG B 607:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAG B 604:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAG A 605:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAG A 604:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAG B 606:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OXY B 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OXY A 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.