



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:41 AM UTC

PDB ID : 5JQS / pdb_00005jqs
Title : Crystal structure of deubiquitinase MINDY-1 in complex with Ubiquitin
Authors : Abdul Rehman, S.A.; Kulathu, Y.
Deposited on : 2016-05-05
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	8% 76% 10% 13%
2	D	76	96% 75% 18% 5%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3157 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein FAM63A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	0	2	0
			1968	1268	320	368	12			


There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	96	GLY	-	expression tag	UNP Q8N5J2
A	97	PRO	-	expression tag	UNP Q8N5J2
A	98	LEU	-	expression tag	UNP Q8N5J2
A	99	GLY	-	expression tag	UNP Q8N5J2
A	100	SER	-	expression tag	UNP Q8N5J2
A	101	PRO	-	expression tag	UNP Q8N5J2
A	102	GLU	-	expression tag	UNP Q8N5J2
A	103	PHE	-	expression tag	UNP Q8N5J2
A	104	PRO	-	expression tag	UNP Q8N5J2
A	105	GLY	-	expression tag	UNP Q8N5J2
A	106	ARG	-	expression tag	UNP Q8N5J2
A	107	LEU	-	expression tag	UNP Q8N5J2
A	108	GLU	-	expression tag	UNP Q8N5J2
A	109	MET	-	expression tag	UNP Q8N5J2

- Molecule 2 is a protein called Ubiquitin-40S ribosomal protein S27a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	75	Total	C	N	O	S	0	74	0
			1110	702	189	217	2			

- Molecule 3 is prop-2-en-1-amine (CCD ID: AYE) (formula: C₃H₇N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			4	3	1		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0

- Molecule 6 is 1,4-DIETHYLENE DIOXIDE (CCD ID: DIO) (formula: C₄H₈O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 4 2	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	51	Total O 51 51	0	0
7	D	2	Total O 2 2	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protein FAM63A



• Molecule 2: Ubiquitin-40S ribosomal protein S27a



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	82.34Å 82.34Å 332.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.63 – 2.65 48.63 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.63-2.65) 99.8 (48.63-2.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 2.65Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.205 , 0.231 0.212 , 0.235	Depositor DCC
R_{free} test set	1076 reflections (5.26%)	wwPDB-VP
Wilson B-factor (Å ²)	51.5	Xtriage
Anisotropy	0.644	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 72.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3157	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AYE, DIO, CL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/2021	0.91	7/2755 (0.3%)
2	D	0.24	0/1123	0.55	0/1526
All	All	0.49	0/3144	0.80	7/4281 (0.2%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	137	CYS	CA-C-N	-7.46	110.52	119.84
1	A	137	CYS	C-N-CA	-7.46	110.52	119.84
1	A	155	PRO	CA-C-N	7.01	128.61	119.84
1	A	155	PRO	C-N-CA	7.01	128.61	119.84
1	A	246	SER	CA-C-N	5.45	125.27	119.28
1	A	246	SER	C-N-CA	5.45	125.27	119.28
1	A	228	VAL	CB-CA-C	-5.05	105.51	111.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1968	0	1896	16	0
2	D	1110	0	1094	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	4	0	5	2	0
4	A	10	0	0	0	0
4	D	5	0	0	0	0
5	A	1	0	0	0	0
6	A	6	0	8	0	0
7	A	51	0	0	0	0
7	D	2	0	0	0	0
All	All	3157	0	3003	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:42[A]:ARG:HD2	2:D:72[A]:ARG:HD3	1.62	0.81
1:A:318:ASN:HD21	2:D:71[B]:LEU:HD11	1.52	0.72
2:D:42[A]:ARG:HH21	2:D:70[A]:VAL:HG11	1.57	0.69
1:A:128:ILE:HD13	1:A:333:LEU:HD11	1.74	0.69
1:A:197:ALA:O	1:A:200:VAL:HG12	1.97	0.65
1:A:193:ASN:HB3	1:A:231:LEU:HD11	1.80	0.64
1:A:131[A]:GLN:HE22	1:A:137:CYS:HB3	1.68	0.58
2:D:42[A]:ARG:HB2	2:D:70[A]:VAL:O	2.06	0.56
2:D:23[A]:ILE:HB	2:D:52[A]:ASP:HA	1.87	0.55
1:A:114:TYR:CZ	1:A:136:PRO:HG3	2.44	0.51
2:D:5[A]:VAL:HB	2:D:13[A]:ILE:HG12	1.92	0.50
1:A:299:LEU:HD22	1:A:352:ASN:CB	2.42	0.50
1:A:200:VAL:HG21	1:A:225:GLU:HA	1.94	0.49
2:D:42[A]:ARG:HD2	2:D:72[A]:ARG:CD	2.38	0.48
1:A:131[A]:GLN:HE22	3:A:401:AYE:H3A	1.78	0.48
2:D:36[A]:ILE:HG21	2:D:71[A]:LEU:HD11	1.95	0.48
1:A:307:LYS:HG3	1:A:310:GLU:HB2	1.95	0.47
2:D:15[A]:LEU:HD23	2:D:26[A]:VAL:HG13	1.97	0.47
2:D:23[B]:ILE:HB	2:D:52[B]:ASP:HA	1.96	0.46
1:A:131[A]:GLN:HG3	1:A:132:SER:N	2.31	0.46
2:D:61[A]:ILE:HD13	2:D:67[A]:LEU:HD21	1.99	0.45
2:D:23[B]:ILE:HG23	2:D:43[B]:LEU:HD12	1.98	0.45
2:D:44[B]:ILE:HD13	2:D:49[B]:GLN:HA	1.99	0.44
2:D:42[B]:ARG:O	2:D:69[B]:LEU:HD12	2.18	0.44
1:A:111:PRO:HB2	1:A:112:ASP:H	1.61	0.42
1:A:138:PRO:HG3	1:A:320:PHE:CD2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:241:LEU:HD22	1:A:298:GLY:HA2	2.02	0.41
1:A:137:CYS:HB3	3:A:401:AYE:H3A	1.76	0.41
1:A:213:ARG:HB2	1:A:219:ASP:HB3	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/289 (86%)	243 (98%)	6 (2%)	0	100	100
2	D	146/76 (192%)	140 (96%)	6 (4%)	0	100	100
All	All	395/365 (108%)	383 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/253 (85%)	209 (97%)	7 (3%)	34	55
2	D	116/68 (171%)	106 (91%)	10 (9%)	10	16
All	All	332/321 (103%)	315 (95%)	17 (5%)	25	36

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	124	GLU
1	A	132	SER
1	A	140	LEU
1	A	157	GLN
1	A	228	VAL
1	A	333	LEU
1	A	337	GLN
2	D	13[A]	ILE
2	D	13[B]	ILE
2	D	15[A]	LEU
2	D	15[B]	LEU
2	D	71[A]	LEU
2	D	71[B]	LEU
2	D	72[A]	ARG
2	D	72[B]	ARG
2	D	74[A]	ARG
2	D	74[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	D	101	-	4,4,4	0.43	0	6,6,6	0.08	0
4	SO4	A	403	-	4,4,4	0.26	0	6,6,6	0.13	0
6	DIO	A	405	-	6,6,6	0.46	0	6,6,6	0.62	0
4	SO4	A	402	-	4,4,4	0.32	0	6,6,6	0.16	0
3	AYE	A	401	2,1	3,3,3	0.47	0	2,2,2	1.70	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	DIO	A	405	-	-	-	0/1/1/1
3	AYE	A	401	2,1	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	AYE	C1-C2-C3	-2.33	120.00	126.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	AYE	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	251/289 (86%)	0.24	23 (9%)	14 11	26, 44, 106, 174	2 (0%)
2	D	75/76 (98%)	5.61	73 (97%)	0 0	24, 47, 51, 62	74 (98%)
All	All	326/365 (89%)	1.47	96 (29%)	1 1	24, 45, 94, 174	76 (23%)

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	13[A]	ILE	25.3
2	D	12[A]	THR	10.8
2	D	17[A]	VAL	10.1
2	D	38[A]	PRO	10.1
2	D	8[A]	LEU	9.9
2	D	2[A]	GLN	9.9
2	D	56[A]	LEU	9.8
2	D	70[A]	VAL	9.0
2	D	24[A]	GLU	8.9
2	D	61[A]	ILE	8.4
2	D	20[A]	SER	8.3
2	D	47[A]	GLY	7.9
2	D	23[A]	ILE	7.5
1	A	356	ASP	7.3
2	D	52[A]	ASP	7.2
2	D	4[A]	PHE	7.1
2	D	5[A]	VAL	7.0
2	D	22[A]	THR	6.8
2	D	64[A]	GLU	6.7
2	D	41[A]	GLN	6.6
2	D	25[A]	ASN	6.6
2	D	32[A]	ASP	6.6
2	D	21[A]	ASP	6.3
2	D	36[A]	ILE	6.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	53[A]	GLY	6.2
2	D	27[A]	LYS	6.2
2	D	18[A]	GLU	6.0
2	D	1[A]	MET	5.8
2	D	59[A]	TYR	5.8
2	D	26[A]	VAL	5.7
2	D	50[A]	LEU	5.6
1	A	353	VAL	5.5
2	D	63[A]	LYS	5.3
2	D	39[A]	ASP	5.3
2	D	43[A]	LEU	5.2
2	D	46[A]	ALA	5.2
2	D	19[A]	PRO	5.2
2	D	69[A]	LEU	5.2
2	D	34[A]	GLU	5.1
2	D	55[A]	THR	4.9
1	A	352	ASN	4.9
2	D	16[A]	GLU	4.9
1	A	187	GLN	4.8
2	D	44[A]	ILE	4.8
2	D	40[A]	GLN	4.8
2	D	45[A]	PHE	4.7
1	A	133	THR	4.7
2	D	67[A]	LEU	4.7
2	D	54[A]	ARG	4.6
2	D	37[A]	PRO	4.6
2	D	66[A]	THR	4.6
2	D	7[A]	THR	4.6
1	A	111	PRO	4.5
1	A	351	HIS	4.5
2	D	42[A]	ARG	4.5
1	A	357	SER	4.4
2	D	60[A]	ASN	4.4
2	D	71[A]	LEU	4.4
2	D	51[A]	GLU	4.3
2	D	3[A]	ILE	4.3
2	D	68[A]	HIS	4.2
2	D	48[A]	LYS	4.2
2	D	62[A]	GLN	4.1
2	D	35[A]	GLY	3.8
2	D	49[A]	GLN	3.8
2	D	28[A]	ALA	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	355	GLY	3.7
2	D	31[A]	GLN	3.7
2	D	6[A]	LYS	3.6
1	A	180	GLN	3.5
2	D	15[A]	LEU	3.5
2	D	65[A]	SER	3.3
2	D	30[A]	ILE	3.3
2	D	58[A]	ASP	3.3
2	D	75	GLY	3.1
1	A	134	ASN	3.1
1	A	132	SER	3.0
2	D	57[A]	SER	3.0
2	D	9[A]	THR	3.0
2	D	14[A]	THR	2.9
2	D	72[A]	ARG	2.8
1	A	113	PHE	2.7
1	A	354	ASP	2.5
1	A	136	PRO	2.4
2	D	11[A]	LYS	2.4
1	A	131[A]	GLN	2.3
2	D	73[A]	LEU	2.3
1	A	227	SER	2.2
1	A	358	CYS	2.2
2	D	74[A]	ARG	2.2
1	A	112	ASP	2.1
1	A	224	PRO	2.1
1	A	135	GLY	2.1
1	A	225	GLU	2.1
1	A	275	ASN	2.1
2	D	10[A]	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	D	101	5/5	0.43	0.15	123,130,149,169	0
6	DIO	A	405	6/6	0.86	0.25	63,68,75,76	6
3	AYE	A	401	4/4	0.89	0.24	69,76,78,81	0
4	SO4	A	403	5/5	0.89	0.12	65,71,85,89	5
4	SO4	A	402	5/5	0.91	0.26	70,76,95,126	0
5	CL	A	404	1/1	0.98	0.12	62,62,62,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.