



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 08:44 PM UTC

PDB ID : 9JQ6 / pdb_00009jq6
EMDB ID : EMD-61724
Title : Cryo-EM structure of BTN2A1 in complex with antagonist antibody TH002
Authors : Zhang, M.; Wang, Y.Q.; Xiao, J.Y.
Deposited on : 2024-09-27
Resolution : 3.34 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14446 (2.84 - 3.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	532	
1	F	532	
2	H	231	
2	J	231	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	I	218	
3	K	218	
4	A	2	
4	B	2	
4	C	2	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10236 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Butyrophilin subfamily 2 member A1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	E	215	Total	C	N	O	S	0	0
			1691	1067	296	317	11		
1	F	215	Total	C	N	O	S	0	0
			1691	1067	296	317	11		

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	500	GLY	-	expression tag	UNP Q7KYR7
E	501	GLY	-	expression tag	UNP Q7KYR7
E	502	GLY	-	expression tag	UNP Q7KYR7
E	503	GLY	-	expression tag	UNP Q7KYR7
E	504	SER	-	expression tag	UNP Q7KYR7
E	505	TRP	-	expression tag	UNP Q7KYR7
E	506	SER	-	expression tag	UNP Q7KYR7
E	507	HIS	-	expression tag	UNP Q7KYR7
E	508	PRO	-	expression tag	UNP Q7KYR7
E	509	GLN	-	expression tag	UNP Q7KYR7
E	510	PHE	-	expression tag	UNP Q7KYR7
E	511	GLU	-	expression tag	UNP Q7KYR7
E	512	LYS	-	expression tag	UNP Q7KYR7
E	513	GLY	-	expression tag	UNP Q7KYR7
E	514	GLY	-	expression tag	UNP Q7KYR7
E	515	GLY	-	expression tag	UNP Q7KYR7
E	516	SER	-	expression tag	UNP Q7KYR7
E	517	GLY	-	expression tag	UNP Q7KYR7
E	518	GLY	-	expression tag	UNP Q7KYR7
E	519	GLY	-	expression tag	UNP Q7KYR7
E	520	SER	-	expression tag	UNP Q7KYR7
E	521	GLY	-	expression tag	UNP Q7KYR7
E	522	GLY	-	expression tag	UNP Q7KYR7
E	523	SER	-	expression tag	UNP Q7KYR7
E	524	ALA	-	expression tag	UNP Q7KYR7
E	525	TRP	-	expression tag	UNP Q7KYR7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	526	SER	-	expression tag	UNP Q7KYR7
E	527	HIS	-	expression tag	UNP Q7KYR7
E	528	PRO	-	expression tag	UNP Q7KYR7
E	529	GLN	-	expression tag	UNP Q7KYR7
E	530	PHE	-	expression tag	UNP Q7KYR7
E	531	GLU	-	expression tag	UNP Q7KYR7
E	532	LYS	-	expression tag	UNP Q7KYR7
F	500	GLY	-	expression tag	UNP Q7KYR7
F	501	GLY	-	expression tag	UNP Q7KYR7
F	502	GLY	-	expression tag	UNP Q7KYR7
F	503	GLY	-	expression tag	UNP Q7KYR7
F	504	SER	-	expression tag	UNP Q7KYR7
F	505	TRP	-	expression tag	UNP Q7KYR7
F	506	SER	-	expression tag	UNP Q7KYR7
F	507	HIS	-	expression tag	UNP Q7KYR7
F	508	PRO	-	expression tag	UNP Q7KYR7
F	509	GLN	-	expression tag	UNP Q7KYR7
F	510	PHE	-	expression tag	UNP Q7KYR7
F	511	GLU	-	expression tag	UNP Q7KYR7
F	512	LYS	-	expression tag	UNP Q7KYR7
F	513	GLY	-	expression tag	UNP Q7KYR7
F	514	GLY	-	expression tag	UNP Q7KYR7
F	515	GLY	-	expression tag	UNP Q7KYR7
F	516	SER	-	expression tag	UNP Q7KYR7
F	517	GLY	-	expression tag	UNP Q7KYR7
F	518	GLY	-	expression tag	UNP Q7KYR7
F	519	GLY	-	expression tag	UNP Q7KYR7
F	520	SER	-	expression tag	UNP Q7KYR7
F	521	GLY	-	expression tag	UNP Q7KYR7
F	522	GLY	-	expression tag	UNP Q7KYR7
F	523	SER	-	expression tag	UNP Q7KYR7
F	524	ALA	-	expression tag	UNP Q7KYR7
F	525	TRP	-	expression tag	UNP Q7KYR7
F	526	SER	-	expression tag	UNP Q7KYR7
F	527	HIS	-	expression tag	UNP Q7KYR7
F	528	PRO	-	expression tag	UNP Q7KYR7
F	529	GLN	-	expression tag	UNP Q7KYR7
F	530	PHE	-	expression tag	UNP Q7KYR7
F	531	GLU	-	expression tag	UNP Q7KYR7
F	532	LYS	-	expression tag	UNP Q7KYR7

- Molecule 2 is a protein called BTN2A1 antagonist antibody TH002-Fab heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	H	219	Total	C	N	O	S	0	0
			1670	1059	280	323	8		
2	J	221	Total	C	N	O	S	0	0
			1684	1067	282	327	8		

- Molecule 3 is a protein called BTN2A1 antagonist antibody TH002-Fab light chain.


Mol	Chain	Residues	Atoms					AltConf	Trace
3	I	216	Total	C	N	O	S	0	0
			1659	1032	279	343	5		
3	K	216	Total	C	N	O	S	0	0
			1659	1032	279	343	5		

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
4	A	2	Total	C	N	O	0	0
			28	16	2	10		
4	B	2	Total	C	N	O	0	0
			28	16	2	10		
4	C	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				AltConf
5	E	1	Total	C	N	O	0
			14	8	1	5	
5	E	1	Total	C	N	O	0
			14	8	1	5	
5	E	1	Total	C	N	O	0
			14	8	1	5	
5	F	1	Total	C	N	O	0
			14	8	1	5	
5	F	1	Total	C	N	O	0
			14	8	1	5	
5	F	1	Total	C	N	O	0
			14	8	1	5	
5	F	1	Total	C	N	O	0
			14	8	1	5	

SER	TRP	SER	HIS	PRO	GLN	PHE	GLU	LYS	GLY	GLY	SER	VAL	ARG	PRO	PHE	GLY	GLY	GLY	GLY	ALA	TRP	GLU	ASP	SER	HIS	PRO	GLN	PHE	GLY	GLY	ALA	LEU	THR	GLY	ALA	SER	ASN	GLY	VAL	THR	VAL	VAL	PRO	GLU	PHE	LEU	GLY	LEU	THR	GLU	ALA	LEU	HIS	GLY	ARG	VAL	SER	PHE	TYR	GLN	ASN	MET	LEU	ARG	ASP	GLY	SER	GLY	HIS
ILE	TYR	THR	CYS	PRO	ARG	SER	ALA	PHE	SER	VAL	PRO	VAL	ARG	PRO	PHE	PHE	ARG	LEU	GLY	CYS	GLU	ASP	SER	SER	PRO	ILE	PHE	ILE	CYS	PRO	ALA	LEU	THR	THR	GLY	ALA	SER	ASN	GLY	VAL	THR	VAL	VAL	PRO	GLU	GLY	LEU	THR	GLU	ALA	LEU	HIS	ARG	VAL	SER	PHE	TYR	GLN	ASN	MET	LEU	ARG	ASP	GLY	SER	GLY	HIS		


• Molecule 2: BTN2A1 antagonist antibody TH002-Fab heavy chain

Chain H:  76% 18% 5%




• Molecule 2: BTN2A1 antagonist antibody TH002-Fab heavy chain

Chain J:  68% 26% 6%




• Molecule 3: BTN2A1 antagonist antibody TH002-Fab light chain


Chain I:  5% 68% 29%



• Molecule 3: BTN2A1 antagonist antibody TH002-Fab light chain

Chain K:  6% 72% 26%





• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	247369	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60.0	Depositor
Minimum defocus (nm)	1300	Depositor
Maximum defocus (nm)	1700	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	17.000	Depositor
Minimum map value	-0.132	Depositor
Average map value	-0.033	Depositor
Map value standard deviation	0.279	Depositor
Recommended contour level	4	Depositor
Map size (Å)	380.0, 380.0, 380.0	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.95, 0.95, 0.95	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	E	0.27	0/1728	0.51	4/2338 (0.2%)
1	F	0.18	0/1728	0.38	0/2338
2	H	0.27	0/1712	0.51	2/2330 (0.1%)
2	J	0.30	0/1727	0.56	0/2353
3	I	0.24	0/1699	0.47	1/2313 (0.0%)
3	K	0.19	0/1699	0.43	1/2313 (0.0%)
All	All	0.25	0/10293	0.48	8/13985 (0.1%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	K	130	THR	N-CA-C	-9.41	100.98	111.14
1	E	155	GLY	N-CA-C	7.03	121.10	113.58
2	H	195	THR	N-CA-C	-6.04	105.77	113.02
2	H	196	TRP	N-CA-C	-5.59	105.14	112.75
1	E	156	GLY	N-CA-C	5.53	119.61	112.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1691	0	1670	34	0
1	F	1691	0	1670	24	0
2	H	1670	0	1646	34	0
2	J	1684	0	1661	58	0
3	I	1659	0	1569	77	0
3	K	1659	0	1569	42	0
4	A	28	0	25	0	0
4	B	28	0	25	1	0
4	C	28	0	25	0	0
5	E	42	0	39	0	0
5	F	56	0	52	2	0
All	All	10236	0	9951	258	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 258 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:195:THR:HA	2:H:198:SER:HB2	1.47	0.96
3:I:157:SER:HB3	3:I:159:ARG:HH11	1.31	0.92
3:I:159:ARG:HA	3:I:159:ARG:CZ	2.04	0.87
2:H:196:TRP:CZ2	2:H:220:PRO:HA	2.11	0.86
2:H:196:TRP:HZ2	2:H:220:PRO:HA	1.44	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	213/532 (40%)	201 (94%)	12 (6%)	0	100	100
1	F	213/532 (40%)	202 (95%)	11 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	215/231 (93%)	201 (94%)	14 (6%)	0	100	100
2	J	219/231 (95%)	200 (91%)	19 (9%)	0	100	100
3	I	214/218 (98%)	196 (92%)	18 (8%)	0	100	100
3	K	214/218 (98%)	195 (91%)	19 (9%)	0	100	100
All	All	1288/1962 (66%)	1195 (93%)	93 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	186/463 (40%)	182 (98%)	4 (2%)	45	65
1	F	186/463 (40%)	183 (98%)	3 (2%)	55	71
2	H	188/199 (94%)	186 (99%)	2 (1%)	65	75
2	J	190/199 (96%)	185 (97%)	5 (3%)	40	63
3	I	187/189 (99%)	183 (98%)	4 (2%)	47	66
3	K	187/189 (99%)	186 (100%)	1 (0%)	81	82
All	All	1124/1702 (66%)	1105 (98%)	19 (2%)	52	70

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	204	ASN
2	J	216	LYS
3	K	154	ILE
2	J	215	ASP
2	H	196	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
3	K	57	ASN
3	K	78	ASN
3	I	128	GLN
3	I	161	ASN
2	J	99	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1	1,4	14,14,15	0.18	0	17,19,21	0.50	0
4	NAG	A	2	4	14,14,15	0.24	0	17,19,21	0.43	0
4	NAG	B	1	1,4	14,14,15	0.82	1 (7%)	17,19,21	1.01	1 (5%)
4	NAG	B	2	4	14,14,15	0.20	0	17,19,21	0.52	0
4	NAG	C	1	1,4	14,14,15	0.24	0	17,19,21	0.50	0
4	NAG	C	2	4	14,14,15	0.20	0	17,19,21	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	1,4	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	2	4	-	2/6/23/26	0/1/1/1
4	NAG	B	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	B	2	4	-	3/6/23/26	0/1/1/1
4	NAG	C	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	C	2	4	-	0/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1	NAG	C1-C2	2.78	1.56	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1	NAG	C1-O5-C5	3.43	116.79	112.19

There are no chirality outliers.

5 of 9 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	2	NAG	O5-C5-C6-O6
4	A	2	NAG	C4-C5-C6-O6
4	C	1	NAG	C8-C7-N2-C2
4	C	1	NAG	O7-C7-N2-C2
4	B	2	NAG	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	F	603	1	14,14,15	0.28	0	17,19,21	0.40	0
5	NAG	E	602	1	14,14,15	0.30	0	17,19,21	0.56	0
5	NAG	F	602	1	14,14,15	0.25	0	17,19,21	0.37	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	F	604	1	14,14,15	0.28	0	17,19,21	0.92	0
5	NAG	F	601	1	14,14,15	0.51	0	17,19,21	1.35	2 (11%)
5	NAG	E	603	1	14,14,15	0.29	0	17,19,21	0.53	0
5	NAG	E	601	1	14,14,15	0.27	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	F	603	1	-	4/6/23/26	0/1/1/1
5	NAG	E	602	1	-	0/6/23/26	0/1/1/1
5	NAG	F	602	1	-	2/6/23/26	0/1/1/1
5	NAG	F	604	1	-	3/6/23/26	0/1/1/1
5	NAG	F	601	1	-	4/6/23/26	0/1/1/1
5	NAG	E	603	1	-	3/6/23/26	0/1/1/1
5	NAG	E	601	1	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	601	NAG	C2-N2-C7	4.56	129.02	122.90
5	F	601	NAG	C1-C2-N2	2.21	113.91	110.43

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	E	601	NAG	C1-C2-N2-C7
5	E	601	NAG	C8-C7-N2-C2
5	E	601	NAG	O7-C7-N2-C2
5	F	604	NAG	C3-C2-N2-C7
5	F	604	NAG	C8-C7-N2-C2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	604	NAG	1	0
5	F	601	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-61724. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200

6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 199




Y Index: 168




Z Index: 136


6.3.2 Raw map



X Index: 0



Y Index: 0



Z Index: 399


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 4.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 61 nm^3 ; this corresponds to an approximate mass of 55 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.299 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.299 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.34	-	-
Author-provided FSC curve	3.34	3.74	3.39
Unmasked-calculated*	4.18	7.88	4.26

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.18 differs from the reported value 3.34 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-61724 and PDB model 9JQ6. Per-residue inclusion information can be found in section [3](#) on page [8](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 4.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (4).

9.4 Atom inclusion [i](#)



At the recommended contour level, 94% of all backbone atoms, 84% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (4) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8440	0.5590
A	0.8210	0.5280
B	0.5360	0.3120
C	0.8570	0.5070
E	0.8780	0.5970
F	0.8760	0.5920
H	0.8630	0.5710
I	0.8240	0.5360
J	0.8350	0.5490
K	0.7900	0.5110

