



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 26, 2026 – 03:38 AM EDT

PDB ID : 7JR1 / pdb_00007jr1
Title : Crystal structure of the R64F mutant of Bauhinia Bauhinioides Kallikrein Inhibitor complexed with Bovine Trypsin
Authors : Li, M.; Wlodawer, A.; Gustchina, A.
Deposited on : 2020-08-11
Resolution : 2.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	223	
1	B	223	
1	C	223	
1	D	223	
1	E	223	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	223	
2	G	164	
2	H	164	
2	I	164	
2	J	164	
2	K	164	
2	L	164	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 18035 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cationic trypsin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	0	1	0
			1631	1013	279	325	14			
1	B	223	Total	C	N	O	S	0	2	0
			1640	1020	280	326	14			
1	C	223	Total	C	N	O	S	0	0	0
			1629	1012	279	324	14			
1	D	223	Total	C	N	O	S	0	0	0
			1629	1012	279	324	14			
1	E	223	Total	C	N	O	S	0	1	0
			1631	1013	279	325	14			
1	F	223	Total	C	N	O	S	0	2	0
			1637	1019	279	325	14			

- Molecule 2 is a protein called Kunitz-type inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	163	Total	C	N	O	S	0	1	0
			1265	813	214	237	1			
2	H	163	Total	C	N	O	S	0	1	0
			1265	813	214	237	1			
2	I	163	Total	C	N	O	S	0	1	0
			1265	813	214	237	1			
2	J	163	Total	C	N	O	S	0	1	0
			1265	813	214	237	1			
2	K	163	Total	C	N	O	S	0	0	0
			1260	810	214	235	1			
2	L	163	Total	C	N	O	S	0	1	0
			1265	813	214	237	1			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
G	0	GLY	-	expression tag	UNP Q6VEQ7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	64	PHE	ARG	engineered mutation	UNP Q6VEQ7
H	0	GLY	-	expression tag	UNP Q6VEQ7
H	64	PHE	ARG	engineered mutation	UNP Q6VEQ7
I	0	GLY	-	expression tag	UNP Q6VEQ7
I	64	PHE	ARG	engineered mutation	UNP Q6VEQ7
J	0	GLY	-	expression tag	UNP Q6VEQ7
J	64	PHE	ARG	engineered mutation	UNP Q6VEQ7
K	0	GLY	-	expression tag	UNP Q6VEQ7
K	64	PHE	ARG	engineered mutation	UNP Q6VEQ7
L	0	GLY	-	expression tag	UNP Q6VEQ7
L	64	PHE	ARG	engineered mutation	UNP Q6VEQ7

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0
3	F	1	Total O S 5 4 1	0	0
3	F	1	Total O S 5 4 1	0	0
3	I	1	Total O S 5 4 1	0	0
3	J	1	Total O S 5 4 1	0	0
3	K	1	Total O S 5 4 1	0	0
3	L	1	Total O S 5 4 1	0	0
3	L	1	Total O S 5 4 1	0	0

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Na 1 1	0	0
4	E	1	Total Na 1 1	0	0
4	I	1	Total Na 1 1	0	0
4	L	1	Total Na 1 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	60	Total O 60 60	0	0
5	G	36	Total O 36 36	0	0
5	B	41	Total O 41 41	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	47	Total 47	O 47	0	0
5	D	51	Total 51	O 51	0	0
5	E	82	Total 82	O 82	0	0
5	F	74	Total 74	O 74	0	0
5	H	27	Total 27	O 27	0	0
5	I	28	Total 28	O 28	0	0
5	J	30	Total 30	O 30	0	0
5	K	51	Total 51	O 51	0	0
5	L	42	Total 42	O 42	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cationic trypsin




• Molecule 1: Cationic trypsin



• Molecule 1: Cationic trypsin



• Molecule 1: Cationic trypsin






- Molecule 1: Cationic trypsin

Chain E: 77% 22%



- Molecule 1: Cationic trypsin

Chain F: 81% 17%



- Molecule 2: Kunitz-type inhibitor

Chain G: 76% 19%




- Molecule 2: Kunitz-type inhibitor

Chain H: 75% 21%




- Molecule 2: Kunitz-type inhibitor

Chain I: 76% 21%




- Molecule 2: Kunitz-type inhibitor

Chain J: 77% 20% ..




- Molecule 2: Kunitz-type inhibitor

Chain K: 76% 21% ..



- Molecule 2: Kunitz-type inhibitor

Chain L: 78% 18% . .



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	207.06Å 207.06Å 107.23Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	89.66 – 2.05 89.66 – 2.05	Depositor EDS
% Data completeness (in resolution range)	100.0 (89.66-2.05) 100.0 (89.66-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.172 , 0.220 0.172 , 0.210	Depositor DCC
R_{free} test set	1522 reflections (0.93%)	wwPDB-VP
Wilson B-factor (Å ²)	41.5	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 36.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.488 for h,-h-k,-l	Xtriage
Reported twinning fraction	0.500 for H, K, L 0.500 for -K, -H, -L	Depositor
Outliers	9 of 163637 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	18035	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 71.68 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5200e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.08	0/1666	1.38	6/2258 (0.3%)
1	B	1.05	0/1677	1.33	0/2272
1	C	1.05	0/1660	1.34	0/2250
1	D	1.06	0/1660	1.38	5/2250 (0.2%)
1	E	1.11	0/1666	1.35	3/2258 (0.1%)
1	F	1.07	0/1674	1.41	5/2269 (0.2%)
2	G	1.02	0/1300	1.29	2/1767 (0.1%)
2	H	0.99	0/1300	1.32	4/1767 (0.2%)
2	I	1.03	0/1300	1.27	0/1767
2	J	1.06	0/1300	1.29	1/1767 (0.1%)
2	K	1.07	1/1292 (0.1%)	1.39	3/1756 (0.2%)
2	L	1.08	0/1300	1.32	2/1767 (0.1%)
All	All	1.06	1/17795 (0.0%)	1.34	31/24148 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1
2	H	0	2
2	K	0	2
All	All	0	5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	K	123	ILE	C-O	5.55	1.30	1.24

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	162	ALA	CA-C-N	7.25	134.74	121.70
2	L	162	ALA	C-N-CA	7.25	134.74	121.70
1	A	79	ASN	CB-CA-C	6.92	120.68	111.82
1	D	79	ASN	CB-CA-C	6.42	121.66	111.51
2	H	94	ASP	CB-CA-C	6.17	119.20	109.27

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	77	GLU	Peptide
2	H	157	PHE	Peptide
2	H	74	LEU	Peptide
2	K	47	LEU	Peptide
2	K	74	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1631	0	1589	25	0
1	B	1640	0	1605	39	0
1	C	1629	0	1587	23	0
1	D	1629	0	1588	42	0
1	E	1631	0	1589	27	0
1	F	1637	0	1604	24	0
2	G	1265	0	1265	24	0
2	H	1265	0	1265	25	0
2	I	1265	0	1265	23	1
2	J	1265	0	1265	28	0
2	K	1260	0	1261	22	0
2	L	1265	0	1265	29	0
3	A	5	0	0	0	0
3	B	5	0	0	1	0
3	C	5	0	0	0	0
3	D	10	0	0	0	0
3	E	15	0	0	0	0
3	F	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	5	0	0	0	0
3	I	5	0	0	1	0
3	J	5	0	0	0	0
3	K	5	0	0	0	0
3	L	10	0	0	1	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
4	I	1	0	0	0	0
4	L	1	0	0	0	0
5	A	60	0	0	3	0
5	B	41	0	0	4	0
5	C	47	0	0	3	0
5	D	51	0	0	19	0
5	E	82	0	0	5	0
5	F	74	0	0	9	0
5	G	36	0	0	6	0
5	H	27	0	0	2	0
5	I	28	0	0	1	0
5	J	30	0	0	5	0
5	K	51	0	0	6	0
5	L	42	0	0	6	0
All	All	18035	0	17148	318	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 318 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:61:SER:HB2	5:E:423:HOH:O	1.57	1.04
1:D:77:GLU:HA	5:D:406:HOH:O	1.61	0.99
1:D:72:ASN:CG	1:D:75:VAL:HG13	1.88	0.97
1:D:61:SER:HB2	5:D:410:HOH:O	1.65	0.93
1:D:76:VAL:HG13	5:D:411:HOH:O	1.68	0.92

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:96:ILE:O	2:I:131:GLN:NE2[4_655]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/223 (100%)	214 (96%)	7 (3%)	1 (0%)	24	16
1	B	223/223 (100%)	212 (95%)	9 (4%)	2 (1%)	14	7
1	C	221/223 (99%)	209 (95%)	11 (5%)	1 (0%)	24	16
1	D	221/223 (99%)	208 (94%)	12 (5%)	1 (0%)	24	16
1	E	222/223 (100%)	213 (96%)	8 (4%)	1 (0%)	24	16
1	F	223/223 (100%)	212 (95%)	9 (4%)	2 (1%)	14	7
2	G	162/164 (99%)	150 (93%)	10 (6%)	2 (1%)	10	4
2	H	162/164 (99%)	155 (96%)	7 (4%)	0	100	100
2	I	162/164 (99%)	151 (93%)	9 (6%)	2 (1%)	10	4
2	J	162/164 (99%)	151 (93%)	10 (6%)	1 (1%)	21	13
2	K	161/164 (98%)	150 (93%)	10 (6%)	1 (1%)	21	13
2	L	162/164 (99%)	155 (96%)	6 (4%)	1 (1%)	21	13
All	All	2303/2322 (99%)	2180 (95%)	108 (5%)	15 (1%)	21	10

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	165[A]	ASP
1	B	165[B]	ASP
1	A	78	GLY
2	G	118	GLY
2	I	118	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/184 (100%)	181 (98%)	4 (2%)	45	44
1	B	186/184 (101%)	171 (92%)	15 (8%)	11	5
1	C	184/184 (100%)	168 (91%)	16 (9%)	9	4
1	D	184/184 (100%)	173 (94%)	11 (6%)	17	10
1	E	185/184 (100%)	176 (95%)	9 (5%)	22	15
1	F	186/184 (101%)	176 (95%)	10 (5%)	20	12
2	G	138/137 (101%)	129 (94%)	9 (6%)	15	9
2	H	138/137 (101%)	128 (93%)	10 (7%)	13	7
2	I	138/137 (101%)	130 (94%)	8 (6%)	18	11
2	J	138/137 (101%)	133 (96%)	5 (4%)	31	26
2	K	137/137 (100%)	129 (94%)	8 (6%)	18	11
2	L	138/137 (101%)	128 (93%)	10 (7%)	13	7
All	All	1937/1926 (101%)	1822 (94%)	115 (6%)	18	10

5 of 115 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	114	LEU
2	L	103	THR
1	F	167	SER
2	L	83	SER
2	K	36	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	240	GLN
2	H	28	HIS
2	J	10	GLN
1	C	64	GLN
1	C	50	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 4 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	G	201	-	4,4,4	0.32	0	6,6,6	0.10	0
3	SO4	L	202	-	4,4,4	0.30	0	6,6,6	0.12	0
3	SO4	K	201	-	4,4,4	0.35	0	6,6,6	0.10	0
3	SO4	E	302	-	4,4,4	0.28	0	6,6,6	0.12	0
3	SO4	F	302	-	4,4,4	0.37	0	6,6,6	0.12	0
3	SO4	I	202	-	4,4,4	0.34	0	6,6,6	0.09	0
3	SO4	F	301	-	4,4,4	0.36	0	6,6,6	0.08	0
3	SO4	A	301	-	4,4,4	0.36	0	6,6,6	0.07	0
3	SO4	J	201	-	4,4,4	0.32	0	6,6,6	0.10	0
3	SO4	D	302	-	4,4,4	0.31	0	6,6,6	0.05	0
3	SO4	C	302	-	4,4,4	0.30	0	6,6,6	0.12	0
3	SO4	E	304	-	4,4,4	0.33	0	6,6,6	0.07	0
3	SO4	B	301	-	4,4,4	0.32	0	6,6,6	0.08	0
3	SO4	E	303	-	4,4,4	0.36	0	6,6,6	0.08	0
3	SO4	D	301	-	4,4,4	0.33	0	6,6,6	0.09	0
3	SO4	L	203	-	4,4,4	0.35	0	6,6,6	0.10	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	I	202	SO4	1	0
3	B	301	SO4	1	0
3	L	203	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/223 (100%)	-1.39	0 100 100	31, 45, 62, 98	1 (0%)
1	B	223/223 (100%)	-1.36	0 100 100	31, 51, 72, 91	2 (0%)
1	C	223/223 (100%)	-1.40	0 100 100	35, 50, 67, 94	0
1	D	223/223 (100%)	-1.40	0 100 100	30, 47, 61, 106	0
1	E	223/223 (100%)	-1.34	0 100 100	29, 42, 57, 86	1 (0%)
1	F	223/223 (100%)	-1.31	0 100 100	30, 48, 68, 85	2 (0%)
2	G	163/164 (99%)	-1.38	0 100 100	25, 50, 82, 103	1 (0%)
2	H	163/164 (99%)	-1.41	0 100 100	27, 55, 83, 98	1 (0%)
2	I	163/164 (99%)	-1.38	0 100 100	27, 52, 84, 102	1 (0%)
2	J	163/164 (99%)	-1.43	0 100 100	25, 48, 76, 100	1 (0%)
2	K	163/164 (99%)	-1.39	0 100 100	28, 46, 76, 92	0
2	L	163/164 (99%)	-1.36	0 100 100	24, 48, 75, 87	1 (0%)
All	All	2316/2322 (99%)	-1.38	0 100 100	24, 48, 75, 106	11 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	K	201	5/5	0.98	0.05	64,64,67,68	5
3	SO4	B	301	5/5	0.99	0.03	61,62,64,65	5
3	SO4	C	302	5/5	0.99	0.03	49,49,56,59	0
3	SO4	D	301	5/5	0.99	0.05	44,45,50,50	5
3	SO4	D	302	5/5	0.99	0.04	44,45,48,48	5
3	SO4	E	302	5/5	0.99	0.05	28,28,32,33	5
3	SO4	F	301	5/5	0.99	0.03	40,42,44,45	5
3	SO4	J	201	5/5	0.99	0.04	46,47,50,53	5
3	SO4	A	301	5/5	0.99	0.09	33,34,36,37	5
4	NA	I	201	1/1	0.99	0.05	46,46,46,46	0
3	SO4	F	302	5/5	1.00	0.03	38,38,41,44	5
3	SO4	I	202	5/5	1.00	0.01	76,78,78,84	0
3	SO4	E	303	5/5	1.00	0.02	56,57,60,67	0
3	SO4	E	304	5/5	1.00	0.04	58,59,61,63	5
3	SO4	L	202	5/5	1.00	0.03	67,71,79,84	0
3	SO4	L	203	5/5	1.00	0.02	66,70,71,72	0
4	NA	C	301	1/1	1.00	0.05	38,38,38,38	0
4	NA	E	301	1/1	1.00	0.07	40,40,40,40	0
3	SO4	G	201	5/5	1.00	0.01	65,66,69,76	0
4	NA	L	201	1/1	1.00	0.02	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.