



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 01:12 PM UTC

PDB ID : 7JRX / pdb_00007jrx
Title : Crystal structure of the R64F mutant of Bauhinia Bauhinioides complexed with Bovine Chymotrypsin
Authors : Li, M.; Wlodawer, A.; Gustchina, A.
Deposited on : 2020-08-13
Resolution : 1.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	13	8% 77% 23%
1	a	13	8% 77% 15% 8%
2	B	131	% 95% 5%
2	b	131	% 92% 7%
3	C	97	2% 86% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	c	97	85% 15%
4	I	166	16% 78% 20% ..
4	i	166	19% 80% 16% ...

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6902 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chymotrypsin A chain A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	13	Total	C	N	O	S	0	0	0
			86	56	14	15	1			
1	a	12	Total	C	N	O	S	0	0	0
			78	50	13	14	1			

- Molecule 2 is a protein called Chymotrypsin A chain B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	131	Total	C	N	O	S	0	3	0
			1004	635	166	199	4			
2	b	131	Total	C	N	O	S	0	4	0
			1002	633	165	200	4			

- Molecule 3 is a protein called Chymotrypsin A chain C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	97	Total	C	N	O	S	0	2	0
			710	441	124	138	7			
3	c	97	Total	C	N	O	S	0	4	0
			716	445	125	139	7			

- Molecule 4 is a protein called Kunitz-type inhibitor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I	163	Total	C	N	O	S	0	3	0
			1264	813	212	238	1			
4	i	163	Total	C	N	O	S	0	2	0
			1269	817	214	237	1			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	0	GLY	-	expression tag	UNP Q6VEQ7
I	64	PHE	ARG	engineered mutation	UNP Q6VEQ7
i	0	GLY	-	expression tag	UNP Q6VEQ7
i	64	PHE	ARG	engineered mutation	UNP Q6VEQ7



- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	21	Total O 21 21	0	0
5	B	173	Total O 173 173	0	0
5	C	88	Total O 88 88	0	0
5	I	99	Total O 99 99	0	0
5	a	21	Total O 21 21	0	0
5	b	184	Total O 184 184	0	0
5	c	90	Total O 90 90	0	0
5	i	97	Total O 97 97	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chymotrypsin A chain A





- Molecule 1: Chymotrypsin A chain A




- Molecule 2: Chymotrypsin A chain B



- Molecule 2: Chymotrypsin A chain B



- Molecule 3: Chymotrypsin A chain C



- Molecule 3: Chymotrypsin A chain C

Chain c:

85%

15%



● Molecule 4: Kunitz-type inhibitor


Chain I:

16%

78%

20%

..



● Molecule 4: Kunitz-type inhibitor



Chain i:

19%

80%

16%

...



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	60.02Å 60.69Å 425.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.02 – 1.77 71.02 – 1.77	Depositor EDS
% Data completeness (in resolution range)	94.9 (71.02-1.77) 92.9 (71.02-1.77)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.175 , 0.208 (Not available) , 0.228	Depositor DCC
R_{free} test set	1451 reflections (1.98%)	wwPDB-VP
Wilson B-factor (Å ²)	16.3	Xtriage
Anisotropy	0.118	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.076 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6902	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	0/87	1.32	0/119
1	a	0.96	0/79	1.23	0/108
2	B	1.07	0/1028	1.12	0/1400
2	b	1.10	0/1031	1.14	0/1404
3	C	1.13	1/730 (0.1%)	1.25	3/993 (0.3%)
3	c	1.11	0/745	1.16	0/1014
4	I	1.03	0/1308	1.29	2/1778 (0.1%)
4	i	1.03	0/1309	1.23	1/1779 (0.1%)
All	All	1.07	1/6317 (0.0%)	1.21	6/8595 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	190	SER	C-O	5.71	1.30	1.23

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	166	THR	CB-CA-C	-11.63	91.48	110.79
3	C	166	THR	N-CA-CB	7.75	121.51	110.12
4	I	107[A]	SER	CA-CB-OG	-7.31	96.48	111.10
4	I	107[B]	SER	CA-CB-OG	-7.31	96.48	111.10
3	C	166	THR	CA-CB-OG1	5.33	117.59	109.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	86	0	95	3	0
1	a	78	0	84	3	0
2	B	1004	0	972	18	0
2	b	1002	0	966	19	1
3	C	710	0	707	12	0
3	c	716	0	717	17	0
4	I	1264	0	1268	27	0
4	i	1269	0	1274	30	2
5	A	21	0	0	1	0
5	B	173	0	0	9	0
5	C	88	0	0	7	0
5	I	99	0	0	15	0
5	a	21	0	0	3	0
5	b	184	0	0	10	0
5	c	90	0	0	6	0
5	i	97	0	0	10	1
All	All	6902	0	6083	114	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 114 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:163:LEU:HD22	5:C:384:HOH:O	1.12	1.30
2:B:35:ASP:HB2	5:B:206:HOH:O	1.38	1.17
2:b:35:ASP:OD2	2:b:39[A]:PHE:CE2	1.98	1.16
2:B:49:GLU:HG2	5:B:336:HOH:O	1.44	1.15
2:B:35:ASP:OD2	2:B:39[A]:PHE:CE2	1.99	1.14

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:i:92:GLN:OE1	4:i:92:GLN:OE1[4_556]	1.72	0.48
2:b:18[A]:ASN:OD1	4:i:132:THR:OG1[5_555]	1.94	0.26
5:i:217:HOH:O	5:i:217:HOH:O[4_566]	2.00	0.20

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	11/13 (85%)	11 (100%)	0	0	100	100
1	a	10/13 (77%)	10 (100%)	0	0	100	100
2	B	132/131 (101%)	129 (98%)	3 (2%)	0	100	100
2	b	133/131 (102%)	130 (98%)	3 (2%)	0	100	100
3	C	97/97 (100%)	96 (99%)	1 (1%)	0	100	100
3	c	99/97 (102%)	97 (98%)	2 (2%)	0	100	100
4	I	164/166 (99%)	157 (96%)	6 (4%)	1 (1%)	21	9
4	i	163/166 (98%)	157 (96%)	4 (2%)	2 (1%)	10	2
All	All	809/814 (99%)	787 (97%)	19 (2%)	3 (0%)	30	16

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	i	27	GLY
4	I	27	GLY
4	i	129	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	10/10 (100%)	9 (90%)	1 (10%)	7	1
1	a	9/10 (90%)	9 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	112/109 (103%)	111 (99%)	1 (1%)	70	59
2	b	113/109 (104%)	112 (99%)	1 (1%)	70	59
3	C	79/77 (103%)	75 (95%)	4 (5%)	21	5
3	c	81/77 (105%)	79 (98%)	2 (2%)	42	22
4	I	139/139 (100%)	127 (91%)	12 (9%)	10	1
4	i	139/139 (100%)	134 (96%)	5 (4%)	31	10
All	All	682/670 (102%)	656 (96%)	26 (4%)	31	9

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	I	132	THR
4	I	144	LYS
4	i	119	GLU
4	I	142	ASN
2	b	99	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
4	I	142	ASN
2	b	34	GLN
4	i	92	GLN
3	c	204	ASN
4	i	75	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	13/13 (100%)	0.31	1 (7%) 19 23	14, 16, 36, 54	0
1	a	12/13 (92%)	0.31	1 (8%) 17 20	13, 17, 38, 51	0
2	B	131/131 (100%)	-0.33	1 (0%) 82 87	7, 13, 23, 41	3 (2%)
2	b	131/131 (100%)	-0.36	1 (0%) 82 87	7, 13, 23, 30	4 (3%)
3	C	97/97 (100%)	-0.20	2 (2%) 63 72	8, 13, 28, 48	2 (2%)
3	c	97/97 (100%)	-0.25	0 100 100	8, 14, 24, 38	4 (4%)
4	I	163/166 (98%)	0.84	26 (15%) 5 4	11, 27, 64, 84	3 (1%)
4	i	163/166 (98%)	0.90	32 (19%) 3 3	12, 26, 59, 76	2 (1%)
All	All	807/814 (99%)	0.19	64 (7%) 18 22	7, 17, 54, 84	18 (2%)

The worst 5 of 64 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	i	163	THR	6.5
4	I	36	ILE	5.5
1	A	13	LEU	4.6
4	I	37	GLY	4.5
1	a	12	GLY	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.