



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:56 PM UTC

PDB ID : 4JS8 / pdb_00004js8
Title : Crystal structure of TTK kinase domain with an inhibitor: 401348
Authors : Qiu, W.; Plotnikov, A.N.; Plotnikova, O.; Feher, M.; Awrey, D.E.; Chirgadze, N.Y.
Deposited on : 2013-03-22
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	281	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	804	-	-	X	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 2276 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual specificity protein kinase TTK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	261	2136	1367	354	400	2	13	0	0	0

- Molecule 2 is 4-(cyclohexylmethoxy)-3-{4-[(1-methylpiperidin-4-yl)oxy]phenyl}-2H-indazole (CCD ID: 1PF) (formula: C₂₆H₃₃N₃O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	31	26	3	2	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	73	Total	O	0	0
			73	73		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity protein kinase TTK



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	70.70Å 107.94Å 113.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.06 – 1.94 39.06 – 1.94	Depositor EDS
% Data completeness (in resolution range)	98.9 (39.06-1.94) 99.5 (39.06-1.94)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 1.95Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.231 , 0.253 0.246 , 0.282	Depositor DCC
R_{free} test set	1074 reflections (3.35%)	wwPDB-VP
Wilson B-factor (Å ²)	48.7	Xtriage
Anisotropy	0.408	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.011 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2276	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, SO4, 1PF, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/2159	1.16	0/2914

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2136	0	2140	21	0
2	A	31	0	33	2	0
3	A	10	0	0	0	0
4	A	14	0	20	7	0
5	A	12	0	16	1	0
6	A	73	0	0	0	0
All	All	2276	0	2209	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:571:GLU:HG2	4:A:804:PEG:H42	1.66	0.76
1:A:571:GLU:CG	4:A:804:PEG:H42	2.16	0.75
1:A:600:MET:HE1	4:A:804:PEG:H41	1.76	0.68
1:A:605:GLY:H	2:A:801:1PF:H32	1.44	0.65
1:A:786:LEU:HD22	4:A:805:PEG:H32	1.81	0.62
1:A:686:TPO:HG23	1:A:687:VAL:H	1.66	0.59
1:A:774:ARG:HB2	5:A:806:GOL:H2	1.84	0.59
1:A:789:PRO:HB3	1:A:793:ILE:HD12	1.85	0.58
1:A:653:PHE:HB3	1:A:660:LEU:HD22	1.85	0.57
1:A:571:GLU:HG3	4:A:804:PEG:H42	1.88	0.55
1:A:686:TPO:HG23	1:A:688:ASN:H	1.74	0.53
1:A:686:TPO:HG23	1:A:687:VAL:N	2.25	0.52
1:A:686:TPO:HG21	1:A:688:ASN:HD22	1.74	0.50
1:A:654:LEU:HD11	2:A:801:1PF:N12	2.26	0.50
1:A:555:VAL:HB	1:A:598:ILE:HB	1.96	0.47
1:A:571:GLU:HG2	4:A:804:PEG:C4	2.39	0.47
1:A:539:VAL:HG23	1:A:671:MET:HE1	1.97	0.46
1:A:764:LEU:HD13	1:A:790:TYR:CZ	2.51	0.45
1:A:613:LEU:HD21	1:A:727:MET:HG2	1.98	0.45
1:A:786:LEU:CD2	4:A:805:PEG:H32	2.45	0.44
1:A:728:THR:HG22	1:A:764:LEU:HD21	2.01	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/281 (90%)	249 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	239/258 (93%)	225 (94%)	14 (6%)	18 6

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	521	LYS
1	A	565	LEU
1	A	566	ASP
1	A	575	LEU
1	A	586	ILE
1	A	587	ARG
1	A	616	LYS
1	A	617	LYS
1	A	643	ILE
1	A	654	LEU
1	A	669	ASN
1	A	711	ILE
1	A	738	ILE
1	A	767	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	556	ASN
1	A	606	ASN
1	A	641	HIS
1	A	688	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	686	1	8,10,11	1.00	0	10,14,16	1.62	1 (10%)
1	TPO	A	675	1	8,10,11	1.30	1 (12%)	10,14,16	1.62	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	686	1	-	6/9/11/13	-
1	TPO	A	675	1	-	2/9/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	675	TPO	P-OG1	-3.03	1.54	1.59

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	686	TPO	CG2-CB-CA	-3.77	105.92	113.26
1	A	675	TPO	P-OG1-CB	-3.77	113.09	123.33

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	675	TPO	N-CA-CB-OG1
1	A	686	TPO	N-CA-CB-OG1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	686	TPO	CG2-CB-OG1-P
1	A	675	TPO	CB-OG1-P-O2P
1	A	686	TPO	CB-OG1-P-O3P
1	A	686	TPO	CB-OG1-P-O1P
1	A	686	TPO	O-C-CA-CB
1	A	686	TPO	N-CA-CB-CG2

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	686	TPO	4	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	802	-	4,4,4	0.24	0	6,6,6	0.08	0
2	1PF	A	801	-	35,35,35	1.68	5 (14%)	42,48,48	1.86	13 (30%)
5	GOL	A	806	-	5,5,5	0.04	0	5,5,5	0.11	0
4	PEG	A	804	-	6,6,6	0.10	0	5,5,5	0.11	0
3	SO4	A	803	-	4,4,4	0.26	0	6,6,6	0.08	0
5	GOL	A	807	-	5,5,5	0.11	0	5,5,5	0.25	0
4	PEG	A	805	-	6,6,6	0.33	0	5,5,5	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1PF	A	801	-	-	4/13/31/31	0/5/5/5
5	GOL	A	806	-	-	0/4/4/4	-
4	PEG	A	804	-	-	3/4/4/4	-
5	GOL	A	807	-	-	0/4/4/4	-
4	PEG	A	805	-	-	2/4/4/4	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	1PF	C10-C11	-4.33	1.41	1.47
2	A	801	1PF	C11-C27	4.28	1.45	1.40
2	A	801	1PF	C15-C14	4.23	1.45	1.40
2	A	801	1PF	C14-N13	2.23	1.36	1.33
2	A	801	1PF	C14-C27	-2.14	1.39	1.43

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	1PF	C01-N02-C31	3.80	117.88	110.63
2	A	801	1PF	C03-N02-C31	3.75	115.51	109.54
2	A	801	1PF	C04-C03-N02	3.42	115.61	111.20
2	A	801	1PF	O19-C20-C21	3.30	118.82	108.73
2	A	801	1PF	C03-C04-C05	3.16	113.75	110.32
2	A	801	1PF	C31-C30-C05	3.02	113.59	110.32
2	A	801	1PF	C01-N02-C03	2.97	116.28	110.63
2	A	801	1PF	C20-O19-C18	2.73	124.27	118.21
2	A	801	1PF	C28-C10-C11	-2.67	117.59	120.76
2	A	801	1PF	C09-C10-C11	2.60	123.85	120.76
2	A	801	1PF	O06-C05-C30	2.29	113.66	108.31
2	A	801	1PF	C30-C31-N02	2.26	114.11	111.20
2	A	801	1PF	C11-N12-N13	-2.17	109.21	112.88

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	1PF	O19-C20-C21-C26
4	A	805	PEG	O1-C1-C2-O2
4	A	805	PEG	O2-C3-C4-O4

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	801	1PF	O19-C20-C21-C22
4	A	804	PEG	O1-C1-C2-O2
2	A	801	1PF	C08-C07-O06-C05
2	A	801	1PF	C29-C07-O06-C05
4	A	804	PEG	C4-C3-O2-C2
4	A	804	PEG	C1-C2-O2-C3

There are no ring outliers.

4 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	1PF	2	0
5	A	806	GOL	1	0
4	A	804	PEG	5	0
4	A	805	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/281 (92%)	1.68	85 (32%) 1 0	38, 60, 91, 117	0

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	711	ILE	7.6
1	A	622	TRP	6.2
1	A	684	VAL	5.8
1	A	748	ILE	5.5
1	A	747	ILE	5.2
1	A	698	MET	5.2
1	A	741	ILE	4.9
1	A	793	ILE	4.9
1	A	737	ILE	4.8
1	A	695	ILE	4.7
1	A	710	LYS	4.6
1	A	738	ILE	4.5
1	A	754	ILE	4.3
1	A	742	SER	4.1
1	A	746	ALA	3.9
1	A	687	VAL	3.7
1	A	674	ASP	3.7
1	A	522	GLY	3.7
1	A	536	SER	3.5
1	A	617	LYS	3.5
1	A	643	ILE	3.4
1	A	534	GLY	3.3
1	A	735	GLN	3.2
1	A	736	GLN	3.2
1	A	566	ASP	3.2
1	A	745	HIS	3.2
1	A	694	ALA	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	685	GLY	3.2
1	A	750	PRO	3.1
1	A	744	LEU	3.1
1	A	619	ILE	3.1
1	A	740	GLN	3.1
1	A	615	LYS	3.0
1	A	613	LEU	3.0
1	A	521	LYS	3.0
1	A	743	LYS	3.0
1	A	578	LEU	2.9
1	A	763	ASP	2.9
1	A	576	ASN	2.9
1	A	518	ILE	2.9
1	A	667	ILE	2.9
1	A	665	PHE	2.8
1	A	577	LYS	2.8
1	A	764	LEU	2.8
1	A	729	TYR	2.7
1	A	535	GLY	2.7
1	A	728	THR	2.7
1	A	734	PHE	2.7
1	A	557	LEU	2.7
1	A	759	ILE	2.7
1	A	657	ASP	2.6
1	A	730	GLY	2.6
1	A	568	TYR	2.6
1	A	689	TYR	2.6
1	A	760	PRO	2.6
1	A	756	PHE	2.6
1	A	531	ILE	2.5
1	A	656	VAL	2.5
1	A	559	GLU	2.5
1	A	749	ASP	2.5
1	A	515	ASN	2.4
1	A	524	ILE	2.4
1	A	612	TRP	2.4
1	A	589	TYR	2.4
1	A	766	ASP	2.4
1	A	726	TYR	2.3
1	A	772	LEU	2.3
1	A	739	ASN	2.3
1	A	658	GLY	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	696	LYS	2.3
1	A	527	ILE	2.2
1	A	791	VAL	2.2
1	A	571	GLU	2.2
1	A	565	LEU	2.2
1	A	776	PRO	2.2
1	A	520	VAL	2.1
1	A	671	MET	2.1
1	A	755	GLU	2.1
1	A	709	SER	2.1
1	A	528	LEU	2.1
1	A	663	ILE	2.1
1	A	564	THR	2.1
1	A	621	PRO	2.1
1	A	765	GLN	2.1
1	A	773	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	A	686	11/12	0.65	0.19	81,82,84,84	0
1	TPO	A	675	11/12	0.66	0.20	77,116,116,148	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	803	5/5	0.69	0.14	129,129,129,129	0
4	PEG	A	805	7/7	0.73	0.26	93,93,94,94	0
5	GOL	A	806	6/6	0.74	0.25	90,91,91,91	0
4	PEG	A	804	7/7	0.75	0.35	98,98,99,99	0
5	GOL	A	807	6/6	0.81	0.25	111,112,113,114	0
3	SO4	A	802	5/5	0.84	0.17	135,135,135,135	0
2	1PF	A	801	31/31	0.93	0.11	39,44,51,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.