



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 02:24 PM UTC

PDB ID : 4JSP / pdb_00004jssp
Title : structure of mTORDeltaN-mLST8-ATPgammaS-Mg complex
Authors : Pavletich, N.P.; Yang, H.
Deposited on : 2013-03-22
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1169 (3.32-3.28)
Clashscore	190562	1209 (3.32-3.28)
Ramachandran outliers	187476	1188 (3.32-3.28)
Sidechain outliers	187428	1187 (3.32-3.28)
RSRZ outliers	180081	1169 (3.32-3.28)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1174	
1	B	1174	
2	C	326	
2	D	326	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 22194 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serine/threonine-protein kinase mTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	1058	Total 8608	C 5472	N 1521	O 1552	S 63	0	0	0
1	A	1058	Total 8608	C 5472	N 1521	O 1552	S 63	0	0	0

- Molecule 2 is a protein called Target of rapamycin complex subunit LST8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	317	Total 2456	C 1526	N 436	O 476	S 18	0	0	0
2	C	317	Total 2456	C 1526	N 436	O 476	S 18	0	0	0


- Molecule 3 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (CCD ID: AGS) (formula: C₁₀H₁₆N₅O₁₂P₃S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
3	B	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		
3	A	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		


- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Mg	0	0
			2	2		
4	A	2	Total	Mg	0	0
			2	2		




● Molecule 1: Serine/threonine-protein kinase mTOR






• Molecule 2: Target of rapamycin complex subunit LST8



• Molecule 2: Target of rapamycin complex subunit LST8



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	139.40Å 163.20Å 207.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 3.30 100.00 – 3.30	Depositor EDS
% Data completeness (in resolution range)	89.9 (100.00-3.30) 89.9 (100.00-3.30)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 3.19Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.233 , 0.268 0.234 , 0.271	Depositor DCC
R_{free} test set	1919 reflections (2.42%)	wwPDB-VP
Wilson B-factor (Å ²)	51.6	Xtrriage
Anisotropy	0.559	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 31.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	22194	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5835e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AGS, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	3/8805 (0.0%)	0.96	15/11920 (0.1%)
1	B	0.48	1/8805 (0.0%)	0.95	15/11920 (0.1%)
2	C	0.51	0/2514	0.90	4/3426 (0.1%)
2	D	0.57	0/2514	0.95	8/3426 (0.2%)
All	All	0.50	4/22638 (0.0%)	0.95	42/30692 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
2	C	0	1
2	D	0	1
All	All	0	4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1607	LEU	CA-C	9.29	1.61	1.52
1	B	1968	HIS	N-CA	5.95	1.50	1.46
1	A	1610	GLU	CA-C	5.52	1.58	1.53
1	A	1968	HIS	N-CA	5.13	1.50	1.46

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1915	HIS	N-CA-C	12.46	127.09	109.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1915	HIS	N-CA-C	12.32	126.89	109.15
1	A	1610	GLU	N-CA-C	12.30	126.58	109.11
2	D	170	GLU	N-CA-C	-9.78	99.70	112.41
2	C	170	GLU	N-CA-C	-9.00	100.71	112.41

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1914	GLY	Peptide
1	B	1914	GLY	Peptide
2	C	169	PRO	Peptide
2	D	169	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8608	0	8593	256	6
1	B	8608	0	8593	249	0
2	C	2456	0	2341	114	0
2	D	2456	0	2341	123	0
3	A	31	0	12	1	0
3	B	31	0	12	2	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
All	All	22194	0	21892	724	6

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 724 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2503:ARG:HH11	1:A:2507:LYS:HE2	1.11	1.13

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2503:ARG:HH11	1:B:2507:LYS:HE2	1.10	1.07
2:C:76:ASN:HB3	2:C:77:PRO:HD2	1.38	1.06
2:C:231:ASP:HB3	2:C:233:THR:OG1	1.59	1.02
2:D:231:ASP:HB3	2:D:233:THR:OG1	1.61	1.00

The worst 5 of 6 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1607:LEU:N	1:A:1609:PRO:O[2_554]	1.39	0.81
1:A:1607:LEU:CA	1:A:1609:PRO:O[2_554]	1.77	0.43
1:A:1608:VAL:O	1:A:1608:VAL:O[2_554]	1.84	0.36
1:A:1606:LYS:C	1:A:1609:PRO:O[2_554]	1.92	0.28
1:A:1607:LEU:N	1:A:1609:PRO:C[2_554]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1052/1174 (90%)	938 (89%)	85 (8%)	29 (3%)	4	21
1	B	1052/1174 (90%)	940 (89%)	84 (8%)	28 (3%)	4	22
2	C	315/326 (97%)	270 (86%)	31 (10%)	14 (4%)	2	13
2	D	315/326 (97%)	273 (87%)	27 (9%)	15 (5%)	2	12
All	All	2734/3000 (91%)	2421 (89%)	227 (8%)	86 (3%)	3	20

5 of 86 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1630	VAL
1	B	1650	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1937	GLN
1	B	1970	GLN
1	B	2094	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	931/1024 (91%)	846 (91%)	85 (9%)	9 31
1	B	931/1024 (91%)	846 (91%)	85 (9%)	9 31
2	C	269/276 (98%)	225 (84%)	44 (16%)	2 11
2	D	269/276 (98%)	227 (84%)	42 (16%)	2 12
All	All	2400/2600 (92%)	2144 (89%)	256 (11%)	6 24

5 of 256 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	165	LEU
2	C	174	THR
2	D	91	VAL
2	D	85	ASN
2	C	215	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 103 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1719	HIS
1	A	2082	GLN
2	C	242	GLN
1	A	1760	ASN
1	A	1941	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	AGS	B	3000	4	32,33,33	2.27	10 (31%)	45,52,52	1.68	9 (20%)
3	AGS	A	3000	4	32,33,33	2.19	7 (21%)	45,52,52	1.94	11 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AGS	B	3000	4	-	5/21/38/38	0/3/3/3
3	AGS	A	3000	4	-	5/21/38/38	0/3/3/3

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	3000	AGS	PG-S1G	8.89	2.09	1.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	3000	AGS	PG-S1G	8.86	2.09	1.90
3	B	3000	AGS	C5-C4	4.61	1.47	1.39
3	A	3000	AGS	C5-C4	4.54	1.47	1.39
3	B	3000	AGS	PB-O3B	3.41	1.63	1.59

The worst 5 of 20 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	3000	AGS	C5-C4-N3	-5.63	118.96	126.72
3	B	3000	AGS	C5-C4-N3	-4.99	119.85	126.72
3	A	3000	AGS	N3-C4-N9	4.48	134.78	127.17
3	B	3000	AGS	N3-C4-N9	3.96	133.90	127.17
3	A	3000	AGS	C2-N3-C4	3.95	121.47	111.83

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	3000	AGS	PB-O3B-PG-O2G
3	A	3000	AGS	PB-O3B-PG-O3G
3	B	3000	AGS	C3'-C4'-C5'-O5'
3	B	3000	AGS	O4'-C4'-C5'-O5'
3	B	3000	AGS	PA-O3A-PB-O3B


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	3000	AGS	2	0
3	A	3000	AGS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1058/1174 (90%)	0.49	89 (8%) 17 13	30, 64, 156, 230	0
1	B	1058/1174 (90%)	0.26	58 (5%) 30 21	25, 54, 134, 177	0
2	C	317/326 (97%)	0.26	12 (3%) 44 30	36, 63, 113, 157	0
2	D	317/326 (97%)	0.09	10 (3%) 50 34	27, 43, 94, 127	0
All	All	2750/3000 (91%)	0.33	169 (6%) 27 18	25, 57, 143, 230	0

The worst 5 of 169 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1599	LEU	5.4
1	A	1610	GLU	5.3
1	B	2434	THR	5.2
1	A	1619	TRP	5.2
1	A	1614	ILE	5.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	MG	A	3002	1/1	0.80	0.19	72,72,72,72	0
4	MG	A	3001	1/1	0.82	0.12	66,66,66,66	0
3	AGS	A	3000	31/31	0.90	0.11	51,60,81,85	0
4	MG	B	3002	1/1	0.90	0.13	69,69,69,69	0
4	MG	B	3001	1/1	0.91	0.09	63,63,63,63	0
3	AGS	B	3000	31/31	0.91	0.11	50,56,79,84	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.