



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 08:31 PM UTC

PDB ID : 1JTS / pdb_00001jts
Title : DNA PROTECTION AND BINDING BY E. COLI DPS PROTEIN
Authors : Luo, J.; Liu, D.; White, M.A.; Fox, R.O.
Deposited on : 2001-08-22
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	167	
1	B	167	
1	C	167	
1	D	167	
1	E	167	
1	F	167	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	167	 2% 54% 38% • 7%
1	H	167	 % 54% 37% • 8%
1	I	167	 54% 38% 8%
1	J	167	 % 55% 37% • 5%
1	K	167	 % 53% 38% • 8%
1	L	167	 % 50% 43% • 7%
1	M	167	 59% 35% 7%
1	N	167	 % 53% 40% • 7%
1	O	167	 53% 40% • 7%
1	P	167	 53% 39% • 8%
1	Q	167	 53% 38% • 8%
1	R	167	 54% 39% • 7%
1	S	167	 59% 32% • 7%
1	T	167	 % 52% 41% • 7%
1	U	167	 % 53% 40% • 6%
1	V	167	 54% 37% • 7%
1	W	167	 57% 35% • 7%
1	X	167	 55% 37% • 7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 29850 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA PROTECTION DURING STARVATION PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	155	Total	C	N	O	S	0	0	0
			1224	770	215	235	4			
1	B	155	Total	C	N	O	S	0	0	0
			1224	770	215	235	4			
1	C	154	Total	C	N	O	S	0	0	0
			1216	766	213	233	4			
1	D	154	Total	C	N	O	S	0	0	0
			1216	766	213	233	4			
1	E	156	Total	C	N	O	S	0	0	0
			1231	774	216	237	4			
1	F	155	Total	C	N	O	S	0	0	0
			1224	770	215	235	4			
1	G	155	Total	C	N	O	S	0	0	0
			1224	770	215	235	4			
1	H	154	Total	C	N	O	S	0	0	0
			1216	766	213	233	4			
1	I	154	Total	C	N	O	S	0	0	0
			1216	766	213	233	4			
1	J	158	Total	C	N	O	S	0	0	0
			1245	783	219	239	4			
1	K	154	Total	C	N	O	S	0	0	0
			1216	766	213	233	4			
1	L	155	Total	C	N	O	S	0	0	0
			1224	770	215	235	4			
1	M	156	Total	C	N	O	S	0	0	0
			1231	774	216	237	4			
1	N	156	Total	C	N	O	S	0	0	0
			1231	774	216	237	4			
1	O	156	Total	C	N	O	S	0	0	0
			1231	774	216	237	4			
1	P	154	Total	C	N	O	S	0	0	0
			1216	766	213	233	4			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	154	Total	C	N	O	S	0	0	0
			1216	766	213	233	4			
1	R	156	Total	C	N	O	S	0	0	0
			1231	774	216	237	4			
1	S	156	Total	C	N	O	S	0	0	0
			1231	774	216	237	4			
1	T	156	Total	C	N	O	S	0	0	0
			1231	774	216	237	4			
1	U	157	Total	C	N	O	S	0	0	0
			1236	777	217	238	4			
1	V	155	Total	C	N	O	S	0	0	0
			1224	770	215	235	4			
1	W	156	Total	C	N	O	S	0	0	0
			1231	774	216	237	4			
1	X	156	Total	C	N	O	S	0	0	0
			1231	774	216	237	4			

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	75	CYS	ASP	engineered mutation	UNP P0ABT2
A	78	ALA	ASP	engineered mutation	UNP P0ABT2
B	75	CYS	ASP	engineered mutation	UNP P0ABT2
B	78	ALA	ASP	engineered mutation	UNP P0ABT2
C	75	CYS	ASP	engineered mutation	UNP P0ABT2
C	78	ALA	ASP	engineered mutation	UNP P0ABT2
D	75	CYS	ASP	engineered mutation	UNP P0ABT2
D	78	ALA	ASP	engineered mutation	UNP P0ABT2
E	75	CYS	ASP	engineered mutation	UNP P0ABT2
E	78	ALA	ASP	engineered mutation	UNP P0ABT2
F	75	CYS	ASP	engineered mutation	UNP P0ABT2
F	78	ALA	ASP	engineered mutation	UNP P0ABT2
G	75	CYS	ASP	engineered mutation	UNP P0ABT2
G	78	ALA	ASP	engineered mutation	UNP P0ABT2
H	75	CYS	ASP	engineered mutation	UNP P0ABT2
H	78	ALA	ASP	engineered mutation	UNP P0ABT2
I	75	CYS	ASP	engineered mutation	UNP P0ABT2
I	78	ALA	ASP	engineered mutation	UNP P0ABT2
J	75	CYS	ASP	engineered mutation	UNP P0ABT2
J	78	ALA	ASP	engineered mutation	UNP P0ABT2
K	75	CYS	ASP	engineered mutation	UNP P0ABT2
K	78	ALA	ASP	engineered mutation	UNP P0ABT2
L	75	CYS	ASP	engineered mutation	UNP P0ABT2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	78	ALA	ASP	engineered mutation	UNP P0ABT2
M	75	CYS	ASP	engineered mutation	UNP P0ABT2
M	78	ALA	ASP	engineered mutation	UNP P0ABT2
N	75	CYS	ASP	engineered mutation	UNP P0ABT2
N	78	ALA	ASP	engineered mutation	UNP P0ABT2
O	75	CYS	ASP	engineered mutation	UNP P0ABT2
O	78	ALA	ASP	engineered mutation	UNP P0ABT2
P	75	CYS	ASP	engineered mutation	UNP P0ABT2
P	78	ALA	ASP	engineered mutation	UNP P0ABT2
Q	75	CYS	ASP	engineered mutation	UNP P0ABT2
Q	78	ALA	ASP	engineered mutation	UNP P0ABT2
R	75	CYS	ASP	engineered mutation	UNP P0ABT2
R	78	ALA	ASP	engineered mutation	UNP P0ABT2
S	75	CYS	ASP	engineered mutation	UNP P0ABT2
S	78	ALA	ASP	engineered mutation	UNP P0ABT2
T	75	CYS	ASP	engineered mutation	UNP P0ABT2
T	78	ALA	ASP	engineered mutation	UNP P0ABT2
U	75	CYS	ASP	engineered mutation	UNP P0ABT2
U	78	ALA	ASP	engineered mutation	UNP P0ABT2
V	75	CYS	ASP	engineered mutation	UNP P0ABT2
V	78	ALA	ASP	engineered mutation	UNP P0ABT2
W	75	CYS	ASP	engineered mutation	UNP P0ABT2
W	78	ALA	ASP	engineered mutation	UNP P0ABT2
X	75	CYS	ASP	engineered mutation	UNP P0ABT2
X	78	ALA	ASP	engineered mutation	UNP P0ABT2

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			8	4	1	3		
2	A	1	Total	C	N	O	0	0
			8	4	1	3		
2	C	1	Total	C	N	O	0	0
			8	4	1	3		
2	D	1	Total	C	N	O	0	0
			8	4	1	3		
2	D	1	Total	C	N	O	0	0
			8	4	1	3		
2	D	1	Total	C	N	O	0	0
			8	4	1	3		
2	E	1	Total	C	N	O	0	0
			8	4	1	3		
2	G	1	Total	C	N	O	0	0
			8	4	1	3		
2	G	1	Total	C	N	O	0	0
			8	4	1	3		
2	H	1	Total	C	N	O	0	0
			8	4	1	3		
2	H	1	Total	C	N	O	0	0
			8	4	1	3		
2	L	1	Total	C	N	O	0	0
			8	4	1	3		
2	M	1	Total	C	N	O	0	0
			8	4	1	3		
2	M	1	Total	C	N	O	0	0
			8	4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	P	1	Total	C	N	O	0	0
			8	4	1	3		
2	P	1	Total	C	N	O	0	0
			8	4	1	3		
2	P	1	Total	C	N	O	0	0
			8	4	1	3		
2	Q	1	Total	C	N	O	0	0
			8	4	1	3		
2	R	1	Total	C	N	O	0	0
			8	4	1	3		
2	T	1	Total	C	N	O	0	0
			8	4	1	3		
2	T	1	Total	C	N	O	0	0
			8	4	1	3		
2	W	1	Total	C	N	O	0	0
			8	4	1	3		
2	W	1	Total	C	N	O	0	0
			8	4	1	3		
2	X	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	10	Total	O	0	0
			10	10		
3	B	8	Total	O	0	0
			8	8		
3	C	8	Total	O	0	0
			8	8		
3	D	8	Total	O	0	0
			8	8		
3	E	10	Total	O	0	0
			10	10		
3	F	12	Total	O	0	0
			12	12		
3	G	4	Total	O	0	0
			4	4		
3	H	4	Total	O	0	0
			4	4		
3	I	8	Total	O	0	0
			8	8		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	J	6	Total 6	O 6	0	0
3	K	5	Total 5	O 5	0	0
3	L	8	Total 8	O 8	0	0
3	M	11	Total 11	O 11	0	0
3	N	12	Total 12	O 12	0	0
3	O	11	Total 11	O 11	0	0
3	P	3	Total 3	O 3	0	0
3	Q	7	Total 7	O 7	0	0
3	R	10	Total 10	O 10	0	0
3	S	11	Total 11	O 11	0	0
3	T	16	Total 16	O 16	0	0
3	U	10	Total 10	O 10	0	0
3	V	11	Total 11	O 11	0	0
3	W	27	Total 27	O 27	0	0
3	X	22	Total 22	O 22	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN






• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain E: 60% 32% 7%




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain F: 56% 37% 7%




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain G: 54% 38% 7%




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain H: 54% 37% 8%



• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain I: 54% 38% 8%






• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN






• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain O: 53% 40% 7%




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain P: 53% 39% 8%




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain Q: 53% 38% 8%




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN

Chain R: 54% 39% 7%




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN


Chain S: 59% 32% 7%




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN




• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN



• Molecule 1: DNA PROTECTION DURING STARVATION PROTEIN





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	91.03Å 91.28Å 164.44Å 96.88° 97.18° 119.86°	Depositor
Resolution (Å)	29.76 – 2.60 29.76 – 2.60	Depositor EDS
% Data completeness (in resolution range)	89.0 (29.76-2.60) 88.9 (29.76-2.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 2.57Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.238 , 0.268 0.232 , 0.262	Depositor DCC
R_{free} test set	6574 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	54.5	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for k,h,-h-k-l 0.000 for -k,-h,-l 0.000 for -h,-k,h+k+l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	29850	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 31.98 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0127e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1242	0.92	5/1681 (0.3%)
1	B	0.50	0/1242	0.90	2/1681 (0.1%)
1	C	0.48	0/1234	0.90	4/1670 (0.2%)
1	D	0.50	0/1234	0.91	4/1670 (0.2%)
1	E	0.52	0/1249	0.92	4/1691 (0.2%)
1	F	0.52	0/1242	0.91	3/1681 (0.2%)
1	G	0.41	0/1242	0.87	3/1681 (0.2%)
1	H	0.45	0/1234	0.90	3/1670 (0.2%)
1	I	0.52	0/1234	0.89	3/1670 (0.2%)
1	J	0.49	0/1263	0.93	6/1709 (0.4%)
1	K	0.48	0/1234	0.90	4/1670 (0.2%)
1	L	0.47	0/1242	0.90	3/1681 (0.2%)
1	M	0.53	0/1249	0.92	3/1691 (0.2%)
1	N	0.54	0/1249	0.91	4/1691 (0.2%)
1	O	0.49	0/1249	0.90	4/1691 (0.2%)
1	P	0.50	0/1234	0.90	4/1670 (0.2%)
1	Q	0.49	0/1234	0.87	3/1670 (0.2%)
1	R	0.52	0/1249	0.90	3/1691 (0.2%)
1	S	0.56	0/1249	0.90	5/1691 (0.3%)
1	T	0.62	0/1249	0.97	4/1691 (0.2%)
1	U	0.50	0/1254	0.90	4/1698 (0.2%)
1	V	0.55	0/1242	0.95	5/1681 (0.3%)
1	W	0.59	0/1249	0.95	5/1691 (0.3%)
1	X	0.57	0/1249	0.97	5/1691 (0.3%)
All	All	0.51	0/29848	0.91	93/40402 (0.2%)

There are no bond length outliers.

The worst 5 of 93 bond angle outliers are listed below:

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	W	101	LYS	N-CA-C	8.82	124.09	113.16
1	V	101	LYS	N-CA-C	7.71	122.33	113.15
1	J	15	LEU	N-CA-C	-7.41	100.55	110.55
1	P	53	ASN	N-CA-C	7.40	121.16	112.72
1	V	107	TYR	CA-C-N	7.36	128.52	119.98

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1224	0	1227	70	0
1	B	1224	0	1227	74	0
1	C	1216	0	1221	64	0
1	D	1216	0	1221	69	0
1	E	1231	0	1234	59	0
1	F	1224	0	1227	55	0
1	G	1224	0	1227	69	0
1	H	1216	0	1221	65	0
1	I	1216	0	1221	55	0
1	J	1245	0	1252	65	0
1	K	1216	0	1221	82	0
1	L	1224	0	1227	82	0
1	M	1231	0	1234	54	0
1	N	1231	0	1234	61	0
1	O	1231	0	1234	71	0
1	P	1216	0	1221	71	0
1	Q	1216	0	1221	69	0
1	R	1231	0	1234	71	0
1	S	1231	0	1234	55	0
1	T	1231	0	1234	68	0
1	U	1236	0	1239	69	0
1	V	1224	0	1227	64	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	W	1231	0	1234	57	0
1	X	1231	0	1234	57	0
2	A	16	0	24	0	0
2	C	8	0	12	0	0
2	D	24	0	36	0	0
2	E	8	0	12	1	0
2	G	16	0	24	1	0
2	H	16	0	24	0	0
2	L	8	0	12	0	0
2	M	16	0	24	0	0
2	P	24	0	36	1	0
2	Q	8	0	12	1	0
2	R	8	0	12	0	0
2	T	16	0	24	0	0
2	W	16	0	24	0	0
2	X	8	0	12	0	0
3	A	10	0	0	0	0
3	B	8	0	0	1	0
3	C	8	0	0	0	0
3	D	8	0	0	1	0
3	E	10	0	0	1	0
3	F	12	0	0	2	0
3	G	4	0	0	0	0
3	H	4	0	0	0	0
3	I	8	0	0	2	0
3	J	6	0	0	0	0
3	K	5	0	0	0	0
3	L	8	0	0	1	0
3	M	11	0	0	0	0
3	N	12	0	0	1	0
3	O	11	0	0	0	0
3	P	3	0	0	0	0
3	Q	7	0	0	0	0
3	R	10	0	0	1	0
3	S	11	0	0	0	0
3	T	16	0	0	1	0
3	U	10	0	0	2	0
3	V	11	0	0	2	0
3	W	27	0	0	0	0
3	X	22	0	0	1	0
All	All	29850	0	29794	1484	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 1484 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:R:14:LEU:H	1:R:14:LEU:HD23	1.19	1.02
1:J:12:THR:HA	1:J:27:LYS:NZ	1.75	1.01
1:L:111:ILE:HD11	1:L:116:ASP:HB3	1.45	0.97
1:J:12:THR:HA	1:J:27:LYS:HZ3	1.28	0.96
1:A:109:LEU:HD23	1:B:92:LEU:HD22	1.48	0.94

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	K	83/167 (50%)	81 (98%)	2 (2%)	0	100	100
1	L	9/167 (5%)	9 (100%)	0	0	100	100
1	M	93/167 (56%)	88 (95%)	5 (5%)	0	100	100
1	N	55/167 (33%)	55 (100%)	0	0	100	100
1	O	120/167 (72%)	112 (93%)	8 (7%)	0	100	100
1	P	152/167 (91%)	145 (95%)	7 (5%)	0	100	100
1	Q	152/167 (91%)	140 (92%)	11 (7%)	1 (1%)	18	38
1	R	154/167 (92%)	148 (96%)	6 (4%)	0	100	100
1	S	154/167 (92%)	144 (94%)	10 (6%)	0	100	100
1	T	154/167 (92%)	150 (97%)	4 (3%)	0	100	100
1	U	155/167 (93%)	142 (92%)	12 (8%)	1 (1%)	21	42
1	V	153/167 (92%)	145 (95%)	8 (5%)	0	100	100
1	W	154/167 (92%)	144 (94%)	9 (6%)	1 (1%)	21	42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	X	154/167 (92%)	144 (94%)	10 (6%)	0	100	100
All	All	1742/2338 (74%)	1647 (94%)	92 (5%)	3 (0%)	43	66

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Q	113	ASN
1	U	113	ASN
1	W	20	ASP

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TRS	L	311	-	7,7,7	0.42	0	9,9,9	1.06	0
2	TRS	D	310	-	7,7,7	0.44	0	9,9,9	1.16	0
2	TRS	H	308	-	7,7,7	0.47	0	9,9,9	1.11	1 (11%)
2	TRS	T	320	-	7,7,7	0.48	0	9,9,9	1.08	1 (11%)
2	TRS	E	312	-	7,7,7	0.52	0	9,9,9	1.09	1 (11%)
2	TRS	D	305	-	7,7,7	0.40	0	9,9,9	1.11	1 (11%)
2	TRS	W	315	-	7,7,7	0.34	0	9,9,9	1.04	0
2	TRS	R	319	-	7,7,7	0.31	0	9,9,9	1.10	0
2	TRS	P	317	-	7,7,7	0.42	0	9,9,9	1.07	1 (11%)
2	TRS	W	314	-	7,7,7	0.62	0	9,9,9	1.08	1 (11%)
2	TRS	G	302	-	7,7,7	0.58	0	9,9,9	1.03	0
2	TRS	H	304	-	7,7,7	0.54	0	9,9,9	1.04	0
2	TRS	P	318	-	7,7,7	0.50	0	9,9,9	1.13	1 (11%)
2	TRS	P	322	-	7,7,7	0.41	0	9,9,9	1.21	1 (11%)
2	TRS	X	323	-	7,7,7	0.35	0	9,9,9	1.06	0
2	TRS	G	303	-	7,7,7	0.44	0	9,9,9	1.16	1 (11%)
2	TRS	D	306	-	7,7,7	0.32	0	9,9,9	1.09	1 (11%)
2	TRS	C	307	-	7,7,7	0.43	0	9,9,9	1.05	0
2	TRS	A	309	-	7,7,7	0.34	0	9,9,9	1.14	1 (11%)
2	TRS	M	313	-	7,7,7	0.28	0	9,9,9	1.11	0
2	TRS	M	321	-	7,7,7	0.31	0	9,9,9	0.94	0
2	TRS	Q	324	-	7,7,7	0.33	0	9,9,9	1.08	1 (11%)
2	TRS	A	301	-	7,7,7	0.48	0	9,9,9	1.02	0
2	TRS	T	316	-	7,7,7	0.45	0	9,9,9	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	L	311	-	-	0/9/9/9	-
2	TRS	D	310	-	-	2/9/9/9	-
2	TRS	H	308	-	-	0/9/9/9	-
2	TRS	T	320	-	-	0/9/9/9	-
2	TRS	E	312	-	-	1/9/9/9	-
2	TRS	D	305	-	-	0/9/9/9	-
2	TRS	W	315	-	-	1/9/9/9	-
2	TRS	R	319	-	-	1/9/9/9	-
2	TRS	P	317	-	-	0/9/9/9	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	W	314	-	-	0/9/9/9	-
2	TRS	G	302	-	-	0/9/9/9	-
2	TRS	H	304	-	-	2/9/9/9	-
2	TRS	P	318	-	-	1/9/9/9	-
2	TRS	P	322	-	-	2/9/9/9	-
2	TRS	X	323	-	-	0/9/9/9	-
2	TRS	G	303	-	-	1/9/9/9	-
2	TRS	D	306	-	-	1/9/9/9	-
2	TRS	C	307	-	-	1/9/9/9	-
2	TRS	A	309	-	-	1/9/9/9	-
2	TRS	M	313	-	-	1/9/9/9	-
2	TRS	M	321	-	-	1/9/9/9	-
2	TRS	Q	324	-	-	1/9/9/9	-
2	TRS	A	301	-	-	1/9/9/9	-
2	TRS	T	316	-	-	2/9/9/9	-

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	309	TRS	C1-C-N	-2.29	102.32	108.17
2	G	303	TRS	C1-C-N	-2.20	102.56	108.17
2	D	305	TRS	C1-C-N	-2.15	102.69	108.17
2	P	318	TRS	C1-C-N	-2.15	102.70	108.17
2	D	306	TRS	C1-C-N	-2.13	102.73	108.17

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	310	TRS	N-C-C2-O2
2	H	304	TRS	N-C-C2-O2
2	P	322	TRS	N-C-C2-O2
2	A	309	TRS	C1-C-C2-O2
2	D	306	TRS	C1-C-C2-O2

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	312	TRS	1	0
2	P	318	TRS	1	0
2	G	303	TRS	1	0
2	Q	324	TRS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	155/167 (92%)	0.08	0 100 100	39, 59, 80, 85	0
1	B	155/167 (92%)	0.24	1 (0%) 85 83	39, 63, 78, 86	0
1	C	154/167 (92%)	0.29	0 100 100	42, 65, 84, 95	0
1	D	154/167 (92%)	0.35	5 (3%) 50 44	34, 62, 91, 96	0
1	E	156/167 (93%)	0.03	0 100 100	36, 59, 78, 88	0
1	F	155/167 (92%)	-0.01	0 100 100	35, 57, 86, 98	0
1	G	155/167 (92%)	0.61	3 (1%) 66 61	54, 81, 100, 108	0
1	H	154/167 (92%)	0.44	1 (0%) 85 83	45, 71, 95, 100	0
1	I	154/167 (92%)	-0.03	0 100 100	33, 59, 78, 83	0
1	J	158/167 (94%)	0.37	2 (1%) 75 71	47, 67, 82, 107	0
1	K	154/167 (92%)	0.64	2 (1%) 75 71	53, 75, 90, 94	0
1	L	155/167 (92%)	0.58	1 (0%) 85 83	49, 74, 90, 95	0
1	M	156/167 (93%)	0.08	0 100 100	30, 59, 82, 96	0
1	N	156/167 (93%)	-0.30	1 (0%) 85 83	29, 50, 68, 88	0
1	O	156/167 (93%)	0.18	0 100 100	39, 62, 79, 97	0
1	P	154/167 (92%)	0.12	0 100 100	38, 57, 78, 85	0
1	Q	154/167 (92%)	0.20	0 100 100	37, 65, 85, 89	0
1	R	156/167 (93%)	0.14	0 100 100	40, 59, 83, 89	0
1	S	156/167 (93%)	-0.15	0 100 100	36, 54, 69, 80	0
1	T	156/167 (93%)	-0.26	1 (0%) 85 83	23, 46, 66, 84	0
1	U	157/167 (94%)	0.25	1 (0%) 85 83	38, 61, 78, 89	0
1	V	155/167 (92%)	-0.11	0 100 100	32, 50, 70, 77	0
1	W	156/167 (93%)	-0.24	0 100 100	29, 48, 66, 79	0
1	X	156/167 (93%)	-0.11	0 100 100	30, 49, 78, 105	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	3727/4008 (92%)	0.14	18 (0%) 87 85	23, 61, 87, 108	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	11	ALA	2.7
1	H	92	LEU	2.7
1	T	14	LEU	2.7
1	D	16	TYR	2.4
1	U	11	ALA	2.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TRS	G	302	8/8	0.42	0.22	105,106,107,108	0
2	TRS	P	317	8/8	0.53	0.22	87,88,89,92	0
2	TRS	D	305	8/8	0.56	0.25	83,83,85,86	0
2	TRS	T	320	8/8	0.58	0.25	86,87,88,89	0
2	TRS	X	323	8/8	0.60	0.25	82,83,85,85	0
2	TRS	L	311	8/8	0.61	0.18	74,75,76,77	0
2	TRS	H	308	8/8	0.65	0.18	86,88,89,91	0
2	TRS	W	314	8/8	0.67	0.17	68,69,70,70	0
2	TRS	E	312	8/8	0.82	0.11	73,75,76,76	0
2	TRS	D	306	8/8	0.82	0.14	65,68,69,70	0
2	TRS	W	315	8/8	0.83	0.15	87,88,88,88	0
2	TRS	G	303	8/8	0.83	0.16	101,104,104,105	0
2	TRS	A	309	8/8	0.84	0.12	58,61,61,62	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	TRS	P	318	8/8	0.85	0.12	64,69,70,72	0
2	TRS	A	301	8/8	0.86	0.12	54,56,57,59	0
2	TRS	Q	324	8/8	0.88	0.11	67,69,70,71	0
2	TRS	D	310	8/8	0.89	0.11	64,64,65,66	0
2	TRS	M	321	8/8	0.90	0.11	54,55,56,57	0
2	TRS	C	307	8/8	0.90	0.09	51,52,53,54	0
2	TRS	H	304	8/8	0.91	0.09	54,56,58,59	0
2	TRS	R	319	8/8	0.91	0.11	45,50,53,56	0
2	TRS	P	322	8/8	0.91	0.07	44,46,47,48	0
2	TRS	M	313	8/8	0.93	0.08	48,50,51,52	0
2	TRS	T	316	8/8	0.93	0.11	51,53,56,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.