



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:38 AM UTC

PDB ID : 4JTB / pdb_00004jtb
Title : Crystal structure of Ribosome inactivating protein from Momordica balsamina complexed with phosphate ion at 1.71 Angstrom resolution
Authors : Pandey, S.; Tyagi, T.K.; Singh, A.; Kushwaha, G.S.; Sinha, M.; Bhushan, A.; Kaur, P.; Sharma, S.; Singh, T.P.
Deposited on : 2013-03-23
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	8% 78% 20% .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2094 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called rRNA N-glycosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	246	1918	1220	329	365	4	0	1	0

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	157	Total	O	0	0
			157	157		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: rRNA N-glycosidase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	133.15Å 133.15Å 38.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	66.58 – 1.71 66.58 – 1.71	Depositor EDS
% Data completeness (in resolution range)	100.0 (66.58-1.71) 100.0 (66.58-1.71)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 1.71Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.212 , 0.256 (Not available) , 0.287	Depositor DCC
R_{free} test set	1377 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.016 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2094	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.74	26/1954 (1.3%)	1.28	7/2660 (0.3%)

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	162	ALA	CA-CB	-8.13	1.40	1.53
1	A	131	ALA	CA-CB	-7.23	1.42	1.53
1	A	211	THR	C-O	-6.20	1.17	1.25
1	A	165	LYS	C-O	-6.08	1.17	1.24
1	A	150	ALA	CA-CB	-6.01	1.44	1.53
1	A	85	GLU	N-CA	5.95	1.52	1.46
1	A	167	ILE	C-O	-5.67	1.17	1.24
1	A	70	TYR	C-O	-5.67	1.17	1.23
1	A	54	ASN	C-O	-5.46	1.16	1.23
1	A	48	LEU	C-O	-5.40	1.17	1.23
1	A	175	ALA	C-O	-5.39	1.17	1.24
1	A	132	LEU	C-O	-5.36	1.17	1.24
1	A	64	VAL	C-O	-5.34	1.18	1.24
1	A	151	LEU	C-O	-5.32	1.17	1.24
1	A	135	ALA	CA-CB	5.25	1.61	1.53
1	A	125	ILE	C-O	-5.25	1.18	1.24
1	A	82	PHE	CA-C	-5.22	1.46	1.52
1	A	12	SER	CA-C	5.20	1.59	1.52
1	A	68	ASN	CA-C	5.18	1.58	1.52
1	A	213	THR	C-O	-5.14	1.17	1.24
1	A	40	SER	C-O	-5.10	1.17	1.23
1	A	45	GLY	C-O	-5.08	1.18	1.24
1	A	46	ARG	C-O	-5.06	1.17	1.24
1	A	225	ILE	C-O	-5.05	1.18	1.24
1	A	126	PRO	C-O	-5.04	1.17	1.23
1	A	233	VAL	C-O	-5.00	1.18	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	233	VAL	N-CA-C	10.11	120.03	110.53
1	A	78	THR	N-CA-C	7.06	120.77	112.72
1	A	198	GLN	N-CA-C	6.47	118.33	111.28
1	A	143	ASP	N-CA-C	-5.45	98.13	107.61
1	A	21	LEU	N-CA-C	-5.35	105.35	111.07
1	A	217	ASP	N-CA-C	5.28	118.54	110.36
1	A	160	GLU	N-CA-C	5.13	116.95	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1918	0	1940	17	0
2	A	5	0	0	0	0
3	A	14	0	13	0	0
4	A	157	0	0	2	0
All	All	2094	0	1953	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:134:THR:HG22	4:A:506:HOH:O	1.87	0.73
1:A:90:LEU:O	1:A:93:GLN:HB2	1.94	0.67
1:A:231:ASN:HD22	1:A:231:ASN:H	1.42	0.65
1:A:54:ASN:HB2	1:A:133:ASP:OD1	2.01	0.59
1:A:231:ASN:HD22	1:A:231:ASN:N	2.04	0.54
1:A:185:THR:O	1:A:189:GLU:HG3	2.08	0.52
1:A:13:SER:HA	1:A:16[B]:MET:HE3	1.94	0.48
1:A:16[A]:MET:HE2	1:A:16[A]:MET:HA	1.94	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:77:LEU:HG	1:A:78:THR:HG23	1.97	0.46
1:A:121:PRO:HB2	1:A:123:GLU:OE2	2.16	0.45
1:A:123:GLU:H	1:A:123:GLU:CD	2.23	0.45
1:A:13:SER:HA	1:A:16[B]:MET:CE	2.47	0.44
1:A:93:GLN:HE21	1:A:93:GLN:HB3	1.59	0.43
1:A:222:ARG:HG2	1:A:223:VAL:N	2.34	0.41
1:A:71:ILE:HD12	1:A:155:ILE:HG12	2.02	0.41
1:A:231:ASN:ND2	4:A:446:HOH:O	2.53	0.41
1:A:72:MET:HE2	1:A:72:MET:HA	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	245/246 (100%)	238 (97%)	6 (2%)	1 (0%)	30 16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	69	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	210/209 (100%)	200 (95%)	10 (5%)	23 4

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	SER
1	A	69	VAL
1	A	93	GLN
1	A	97	ARG
1	A	100	ARG
1	A	123	GLU
1	A	124	LYS
1	A	165	LYS
1	A	224	GLN
1	A	231	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN
1	A	115	GLN
1	A	169	GLN
1	A	170	GLN
1	A	221	ASN
1	A	224	GLN
1	A	231	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	A	301	-	4,4,4	1.53	1 (25%)	6,6,6	1.39	1 (16%)
3	NAG	A	302	1	14,14,15	1.04	1 (7%)	17,19,21	2.73	8 (47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	302	1	-	1/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	NAG	C1-C2	3.29	1.56	1.52
2	A	301	PO4	P-O1	2.62	1.56	1.50

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	302	NAG	C1-O5-C5	6.19	120.48	112.19
3	A	302	NAG	C2-N2-C7	4.05	128.32	122.90
3	A	302	NAG	O5-C5-C4	3.95	120.44	110.83
3	A	302	NAG	O5-C1-C2	3.41	116.57	111.29
3	A	302	NAG	O3-C3-C2	3.15	115.94	109.40
3	A	302	NAG	C6-C5-C4	-2.90	105.90	113.02
3	A	302	NAG	O7-C7-N2	2.90	127.10	121.98
3	A	302	NAG	O7-C7-C8	-2.25	118.05	122.05
2	A	301	PO4	O4-P-O2	2.19	114.74	107.91

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	246/246 (100%)	0.68	20 (8%)	18 23	13, 27, 43, 53	2 (0%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	77	LEU	8.6
1	A	231	ASN	4.0
1	A	55	TYR	3.9
1	A	202	ALA	3.5
1	A	246	ILE	3.2
1	A	203	GLN	3.0
1	A	223	VAL	2.8
1	A	97	ARG	2.8
1	A	208	VAL	2.8
1	A	112	GLU	2.5
1	A	216	VAL	2.5
1	A	113	ARG	2.3
1	A	214	VAL	2.3
1	A	28	THR	2.2
1	A	222	ARG	2.2
1	A	220	GLY	2.2
1	A	32	TYR	2.2
1	A	134	THR	2.2
1	A	219	LYS	2.1
1	A	244	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	A	302	14/15	0.67	0.19	46,56,61,63	0
2	PO4	A	301	5/5	0.94	0.10	32,34,40,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.