



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:36 PM UTC

PDB ID : 5JTF / pdb_00005jtf
Title : Crystal structure of ArsN N-acetyltransferase from Pseudomonas putida KT2440
Authors : Venkadesh, S.; Dheeman, D.S.; Kandavelu, P.; Rosen, B.P.
Deposited on : 2016-05-09
Resolution : 2.16 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	7% 70% 15% 15%
1	B	207	4% 68% 15% 15%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2785 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative Phosphinothricin N-acetyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	176	Total	C	N	O	S	0	1	0
			1337	864	239	233	1			
1	B	175	Total	C	N	O	S	0	0	0
			1345	867	240	237	1			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	184	LEU	-	expression tag	UNP Q88LK7
A	185	GLY	-	expression tag	UNP Q88LK7
A	186	PRO	-	expression tag	UNP Q88LK7
A	187	GLU	-	expression tag	UNP Q88LK7
A	188	GLN	-	expression tag	UNP Q88LK7
A	189	LYS	-	expression tag	UNP Q88LK7
A	190	LEU	-	expression tag	UNP Q88LK7
A	191	ILE	-	expression tag	UNP Q88LK7
A	192	SER	-	expression tag	UNP Q88LK7
A	193	GLU	-	expression tag	UNP Q88LK7
A	194	GLU	-	expression tag	UNP Q88LK7
A	195	ASP	-	expression tag	UNP Q88LK7
A	196	LEU	-	expression tag	UNP Q88LK7
A	197	ASN	-	expression tag	UNP Q88LK7
A	198	SER	-	expression tag	UNP Q88LK7
A	199	ALA	-	expression tag	UNP Q88LK7
A	200	VAL	-	expression tag	UNP Q88LK7
A	201	ASP	-	expression tag	UNP Q88LK7
A	202	HIS	-	expression tag	UNP Q88LK7
A	203	HIS	-	expression tag	UNP Q88LK7
A	204	HIS	-	expression tag	UNP Q88LK7
A	205	HIS	-	expression tag	UNP Q88LK7
A	206	HIS	-	expression tag	UNP Q88LK7
A	207	HIS	-	expression tag	UNP Q88LK7
B	184	LEU	-	expression tag	UNP Q88LK7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	185	GLY	-	expression tag	UNP Q88LK7
B	186	PRO	-	expression tag	UNP Q88LK7
B	187	GLU	-	expression tag	UNP Q88LK7
B	188	GLN	-	expression tag	UNP Q88LK7
B	189	LYS	-	expression tag	UNP Q88LK7
B	190	LEU	-	expression tag	UNP Q88LK7
B	191	ILE	-	expression tag	UNP Q88LK7
B	192	SER	-	expression tag	UNP Q88LK7
B	193	GLU	-	expression tag	UNP Q88LK7
B	194	GLU	-	expression tag	UNP Q88LK7
B	195	ASP	-	expression tag	UNP Q88LK7
B	196	LEU	-	expression tag	UNP Q88LK7
B	197	ASN	-	expression tag	UNP Q88LK7
B	198	SER	-	expression tag	UNP Q88LK7
B	199	ALA	-	expression tag	UNP Q88LK7
B	200	VAL	-	expression tag	UNP Q88LK7
B	201	ASP	-	expression tag	UNP Q88LK7
B	202	HIS	-	expression tag	UNP Q88LK7
B	203	HIS	-	expression tag	UNP Q88LK7
B	204	HIS	-	expression tag	UNP Q88LK7
B	205	HIS	-	expression tag	UNP Q88LK7
B	206	HIS	-	expression tag	UNP Q88LK7
B	207	HIS	-	expression tag	UNP Q88LK7

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	33	Total	O	0	0
			33	33		
3	B	62	Total	O	0	0
			62	62		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative Phosphinothricin N-acetyltransferase



- Molecule 1: Putative Phosphinothricin N-acetyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	67.02Å 67.02Å 206.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.64 – 2.16 30.64 – 2.16	Depositor EDS
% Data completeness (in resolution range)	97.9 (30.64-2.16) 97.8 (30.64-2.16)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	11.04 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.238 , 0.266 0.239 , 0.265	Depositor DCC
R_{free} test set	1310 reflections (2.56%)	wwPDB-VP
Wilson B-factor (Å ²)	22.2	Xtriage
Anisotropy	0.367	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 31.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.25$, $\langle L^2 \rangle = 0.11$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	2785	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1372	0.83	4/1874 (0.2%)
1	B	0.34	0/1379	0.82	1/1879 (0.1%)
All	All	0.33	0/2751	0.82	5/3753 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	171	HIS	CA-C-N	6.79	128.33	119.84
1	A	171	HIS	C-N-CA	6.79	128.33	119.84
1	B	125	LEU	N-CA-C	5.69	122.39	109.81
1	A	76	ALA	N-CA-C	5.25	119.41	112.89
1	A	95	ARG	N-CA-C	5.17	116.60	111.07

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	125	LEU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1337	0	1269	12	0
1	B	1345	0	1288	18	0
2	B	8	0	12	1	0
3	A	33	0	0	2	0
3	B	62	0	0	3	0
All	All	2785	0	2569	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:176:LEU:HD22	2:B:301:TRS:H22	1.72	0.70
1:A:101:ARG:HG3	1:A:137:LEU:HD11	1.77	0.67
1:B:44:ARG:NH1	3:B:404:HOH:O	2.33	0.61
1:B:16:GLU:OE2	1:B:44:ARG:NH2	2.37	0.57
1:B:29:THR:O	1:B:151:LYS:NZ	2.31	0.54
1:A:75:ARG:NH1	3:A:306:HOH:O	2.41	0.53
1:B:25:ILE:HG23	1:B:31:ILE:HB	1.89	0.53
1:A:10:ALA:HB2	1:A:58:VAL:HG13	1.93	0.51
1:B:18:ILE:HG22	1:B:43:MET:HE3	1.94	0.50
1:A:139:PHE:HB3	1:A:161:TRP:HB3	1.94	0.50
1:A:55:PRO:HB3	1:A:176:LEU:HD13	1.93	0.49
1:A:17:GLU:O	1:A:21:ILE:HG12	2.11	0.49
1:A:25:ILE:O	1:A:29:THR:OG1	2.25	0.49
1:B:16:GLU:OE1	1:B:16:GLU:N	2.47	0.48
1:A:75:ARG:HD3	1:A:80:TYR:CZ	2.50	0.47
1:B:7:ILE:HG21	1:B:106:VAL:HG21	1.96	0.47
1:B:114:LEU:HD23	1:B:114:LEU:HA	1.76	0.45
1:B:42:GLN:NE2	3:B:414:HOH:O	2.50	0.45
1:B:82:TRP:HB3	1:B:117:ARG:HB2	1.98	0.45
1:A:97:SER:N	1:A:98:GLY:HA3	2.32	0.44
1:A:54:TYR:HB3	1:A:71:ALA:HB3	1.98	0.44
1:B:51:LEU:HD23	1:B:51:LEU:HA	1.84	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:55:PRO:HA	1:B:176:LEU:HG	2.00	0.43
1:B:21:ILE:O	1:B:24:PRO:HD2	2.19	0.43
1:A:42:GLN:HB3	3:A:308:HOH:O	2.17	0.43
1:B:142:ILE:HG13	1:B:160:TYR:O	2.18	0.43
1:B:144:THR:HA	1:B:159:GLY:HA2	2.00	0.43
1:B:8:ARG:HD2	3:B:418:HOH:O	2.19	0.42
1:B:10:ALA:HB2	1:B:58:VAL:HG13	2.01	0.42
1:A:68:TYR:CZ	1:A:89:TYR:HD2	2.40	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	175/207 (84%)	160 (91%)	13 (7%)	2 (1%)	11	7
1	B	171/207 (83%)	161 (94%)	6 (4%)	4 (2%)	5	1
All	All	346/414 (84%)	321 (93%)	19 (6%)	6 (2%)	7	2

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	126	PRO
1	A	93	GLY
1	B	53	THR
1	B	178	PHE
1	B	5	ILE
1	A	5	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/170 (72%)	117 (95%)	6 (5%)	22	19
1	B	127/170 (75%)	121 (95%)	6 (5%)	23	21
All	All	250/340 (74%)	238 (95%)	12 (5%)	23	20

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	ASP
1	A	53	THR
1	A	94	GLN
1	A	103	LEU
1	A	108	LEU
1	A	111	LEU
1	B	53	THR
1	B	103	LEU
1	B	108	LEU
1	B	111	LEU
1	B	125	LEU
1	B	176	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	140	GLN
1	A	147	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TRS	B	301	-	7,7,7	0.40	0	9,9,9	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	B	301	-	-	3/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	TRS	N-C-C2-O2
2	B	301	TRS	C3-C-C2-O2
2	B	301	TRS	C1-C-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	TRS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	176/207 (85%)	0.68	15 (8%) 16 18	15, 36, 69, 77	1 (0%)
1	B	175/207 (84%)	0.41	8 (4%) 37 42	9, 31, 52, 66	0
All	All	351/414 (84%)	0.54	23 (6%) 24 27	9, 34, 57, 77	1 (0%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	169	GLY	4.4
1	A	179	LEU	4.0
1	A	167	ASP	3.5
1	B	169	GLY	3.5
1	B	126	PRO	3.4
1	A	97	SER	2.9
1	A	176	LEU	2.7
1	A	66	VAL	2.7
1	A	96	ARG	2.7
1	A	170	LEU	2.7
1	B	100	ALA	2.5
1	B	104	TYR	2.4
1	B	79	ALA	2.3
1	B	178	PHE	2.3
1	A	79	ALA	2.3
1	A	63	GLY	2.3
1	A	177	GLY	2.3
1	A	178	PHE	2.2
1	B	176	LEU	2.2
1	B	166	GLY	2.2
1	A	77[A]	ARG	2.2
1	A	93	GLY	2.1
1	A	166	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	TRS	B	301	8/8	0.85	0.16	14,28,44,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.