



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 01:26 PM UTC

PDB ID : 8JT8 / pdb_00008jt8
Title : Crystal structure of 5-HT2AR in complex with (R)-IHCH-7179
Authors : Chen, Z.; Fan, L.; Wang, S.
Deposited on : 2023-06-21
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5-hydroxytryptamine receptor 2A, Soluble cytochrome b562.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	355	Total	C	N	O	S	0	0	0
			2720	1786	430	486	18			


There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	67	GLY	-	expression tag	UNP P28223
A	68	GLY	-	expression tag	UNP P28223
A	69	THR	-	expression tag	UNP P28223
A	162	LYS	SER	engineered mutation	UNP P28223
A	164	TRP	MET	engineered mutation	UNP P28223
A	1007	TRP	MET	engineered mutation	UNP P0ABE7
A	1061	GLY	-	linker	UNP P0ABE7
A	1062	SER	-	linker	UNP P0ABE7
A	1063	GLY	-	linker	UNP P0ABE7
A	1064	SER	-	linker	UNP P0ABE7
A	1065	GLY	-	linker	UNP P0ABE7
A	1098	ILE	ARG	engineered mutation	UNP P0ABE7
A	1102	ILE	HIS	engineered mutation	UNP P0ABE7
A	1106	GLY	ARG	engineered mutation	UNP P0ABE7
A	372	ASN	SER	engineered mutation	UNP P28223

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 1-(4-fluorophenyl)-4-[(7R)-2,5,11-triazatetracyclo[7.6.1.0²,7.0¹²,16]hexadeca-1(15),9,12(16),13-tetraen-5-yl]butan-1-one (CCD ID: EZX) (formula: C₂₃H₂₄FN₃O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			28	23	1	3	1		

- Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (CCD ID: OLC) (formula: $C_{21}H_{40}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			16	12	4		
4	A	1	Total	C	O	0	0
			16	12	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			16	12	4		
4	A	1	Total	C	O	0	0
			13	9	4		
4	A	1	Total	C	O	0	0
			14	10	4		
4	A	1	Total	C	O	0	0
			16	12	4		
4	A	1	Total	C	O	0	0
			16	12	4		

- Molecule 5 is CHOLESTEROL (CCD ID: CLR) (formula: $C_{27}H_{46}O$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			24	23	1		
5	A	1	Total	C	O	0	0
			20	19	1		

- Molecule 6 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			16	10	6		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		
7	A	1	Total	C	O	0	0
			7	4	3		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5-hydroxytryptamine receptor 2A, Soluble cytochrome b562



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 2 21	Depositor
Cell constants a, b, c, α , β , γ	48.89Å 54.03Å 177.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.19 – 2.70 47.19 – 2.70	Depositor EDS
% Data completeness (in resolution range)	96.0 (47.19-2.70) 96.0 (47.19-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.248 , 0.270 0.250 , 0.267	Depositor DCC
R_{free} test set	632 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	61.7	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2930	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, 1PE, OLC, EZX, MG, CLR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	0/2772	1.29	2/3774 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	344	ILE	CB-CA-C	5.68	116.06	111.06
1	A	344	ILE	N-CA-C	-5.54	107.42	112.96

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2720	0	2752	17	0
2	A	1	0	0	0	0
3	A	28	0	0	0	0
4	A	107	0	137	3	0
5	A	44	0	63	3	0
6	A	16	0	22	0	0
7	A	14	0	20	0	0
All	All	2930	0	2994	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:LYS:HE3	1:A:332:PHE:HE1	1.59	0.67
1:A:349:CYS:HB2	1:A:352:SER:HB2	1.85	0.58
1:A:93:ILE:HG21	5:A:1205:CLR:H191	1.88	0.55
1:A:1015:LYS:HA	1:A:1018:GLU:HG2	1.89	0.53
1:A:102:GLU:OE1	1:A:393:ARG:NH2	2.40	0.53
1:A:205:GLY:HA3	4:A:1203:OLC:H4	1.91	0.52
1:A:238:GLY:HA3	4:A:1213:OLC:H21A	1.91	0.52
1:A:90:ALA:HA	5:A:1205:CLR:H72	1.96	0.48
1:A:96:ILE:HA	1:A:114:MET:HG3	1.96	0.47
1:A:247:LEU:HD13	1:A:333:VAL:HG21	1.97	0.47
1:A:1091:ALA:HA	1:A:1094:LEU:HD12	1.97	0.47
1:A:107:ASN:OD1	1:A:110:ASN:ND2	2.48	0.46
1:A:342:THR:O	1:A:346:ALA:N	2.49	0.44
1:A:123:LEU:HD13	1:A:158:PHE:HB2	1.98	0.44
1:A:191:LYS:HA	1:A:191:LYS:HD3	1.83	0.44
1:A:209:PRO:HG2	4:A:1213:OLC:H4A	2.02	0.41
5:A:1204:CLR:H222	5:A:1204:CLR:H162	1.65	0.41
1:A:354:ASN:HB3	1:A:357:VAL:HB	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	349/376 (93%)	341 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/322 (89%)	286 (99%)	2 (1%)	76 90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	258	ILE
1	A	260	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 1 is monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	OLC	A	1207	-	15,15,24	0.68	0	16,16,25	0.76	0
4	OLC	A	1203	-	15,15,24	0.37	0	16,16,25	0.70	0
3	EZX	A	1202	-	32,32,32	0.90	2 (6%)	40,46,46	1.34	6 (15%)
7	PEG	A	1212	-	6,6,6	0.36	0	5,5,5	0.15	0
4	OLC	A	1213	-	15,15,24	0.42	0	16,16,25	0.64	0
4	OLC	A	1214	-	15,15,24	0.53	0	16,16,25	0.66	0
4	OLC	A	1208	-	12,12,24	0.61	0	13,13,25	0.54	0
4	OLC	A	1206	-	15,15,24	0.54	0	16,16,25	0.47	0
5	CLR	A	1204	-	27,27,31	0.35	0	43,43,48	0.63	0
6	1PE	A	1210	-	15,15,15	0.42	0	14,14,14	0.42	0
5	CLR	A	1205	-	23,23,31	0.45	0	37,37,48	0.69	0
4	OLC	A	1209	-	13,13,24	0.48	0	14,14,25	0.52	0
7	PEG	A	1211	-	6,6,6	0.17	0	5,5,5	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OLC	A	1207	-	-	5/15/15/24	-
4	OLC	A	1203	-	-	8/15/15/24	-
3	EZX	A	1202	-	-	5/10/31/31	0/4/5/5
7	PEG	A	1212	-	-	3/4/4/4	-
4	OLC	A	1213	-	-	8/15/15/24	-
4	OLC	A	1214	-	-	8/15/15/24	-
4	OLC	A	1208	-	-	4/12/12/24	-
4	OLC	A	1206	-	-	7/15/15/24	-
5	CLR	A	1204	-	-	4/6/64/68	0/4/4/4
6	1PE	A	1210	-	-	7/13/13/13	-
5	CLR	A	1205	-	-	-	0/4/4/4
4	OLC	A	1209	-	-	5/13/13/24	-
7	PEG	A	1211	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1202	EZX	C15-N2	2.61	1.50	1.47
3	A	1202	EZX	C7-N2	2.47	1.45	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1202	EZX	C14-C15-N2	3.77	115.55	109.77
3	A	1202	EZX	C6-N2-C15	-3.68	110.34	115.20
3	A	1202	EZX	C10-C11-N3	2.48	134.72	130.74
3	A	1202	EZX	C5-N1-C16	2.43	113.57	109.53
3	A	1202	EZX	C13-C12-N3	-2.26	107.82	110.31
3	A	1202	EZX	C14-C13-C17	-2.01	117.21	119.05

There are no chirality outliers.

All (66) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1203	OLC	C21-C22-C24-O25
4	A	1206	OLC	C21-C22-C24-O25
4	A	1206	OLC	O23-C22-C24-O25
4	A	1206	OLC	O20-C21-C22-O23
4	A	1209	OLC	C21-C22-C24-O25
4	A	1209	OLC	O23-C22-C24-O25
4	A	1214	OLC	C21-C22-C24-O25
4	A	1203	OLC	O19-C1-O20-C21
4	A	1213	OLC	O19-C1-O20-C21
4	A	1213	OLC	C2-C1-O20-C21
4	A	1203	OLC	C2-C1-O20-C21
6	A	1210	1PE	OH6-C15-C25-OH5
3	A	1202	EZX	C2-C3-C4-N1
4	A	1206	OLC	O20-C21-C22-C24
6	A	1210	1PE	OH5-C14-C24-OH4
7	A	1212	PEG	O2-C3-C4-O4
4	A	1203	OLC	C1-C2-C3-C4
4	A	1208	OLC	C1-C2-C3-C4
4	A	1209	OLC	O20-C21-C22-O23
4	A	1214	OLC	O20-C21-C22-O23
3	A	1202	EZX	C18-C1-C2-C3
4	A	1209	OLC	O20-C21-C22-C24
4	A	1214	OLC	O20-C21-C22-C24
4	A	1208	OLC	C2-C1-O20-C21
3	A	1202	EZX	O1-C1-C2-C3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	A	1211	PEG	O1-C1-C2-O2
7	A	1211	PEG	O2-C3-C4-O4
5	A	1204	CLR	C13-C17-C20-C22
4	A	1208	OLC	O19-C1-O20-C21
4	A	1214	OLC	O23-C22-C24-O25
6	A	1210	1PE	C16-C26-OH6-C15
6	A	1210	1PE	OH2-C12-C22-OH3
4	A	1207	OLC	O20-C21-C22-C24
4	A	1203	OLC	C2-C3-C4-C5
4	A	1203	OLC	C3-C4-C5-C6
5	A	1204	CLR	C16-C17-C20-C21
4	A	1213	OLC	C3-C4-C5-C6
4	A	1203	OLC	O23-C22-C24-O25
5	A	1204	CLR	C13-C17-C20-C21
4	A	1213	OLC	C4-C5-C6-C7
4	A	1206	OLC	C2-C3-C4-C5
4	A	1214	OLC	C4-C5-C6-C7
4	A	1203	OLC	C5-C6-C7-C8
5	A	1204	CLR	C16-C17-C20-C22
4	A	1213	OLC	C5-C6-C7-C8
4	A	1208	OLC	C3-C4-C5-C6
4	A	1207	OLC	O20-C21-C22-O23
4	A	1213	OLC	O20-C21-C22-O23
7	A	1212	PEG	C4-C3-O2-C2
4	A	1207	OLC	C1-C2-C3-C4
4	A	1214	OLC	C1-C2-C3-C4
3	A	1202	EZX	C1-C2-C3-C4
4	A	1206	OLC	C6-C7-C8-C9
4	A	1213	OLC	C2-C3-C4-C5
6	A	1210	1PE	C23-C13-OH4-C24
4	A	1209	OLC	C4-C5-C6-C7
4	A	1214	OLC	C3-C4-C5-C6
3	A	1202	EZX	C3-C4-N1-C16
4	A	1206	OLC	C3-C4-C5-C6
4	A	1214	OLC	C5-C6-C7-C8
6	A	1210	1PE	OH4-C13-C23-OH3
4	A	1213	OLC	C6-C7-C8-C9
4	A	1207	OLC	O23-C22-C24-O25
7	A	1212	PEG	O1-C1-C2-O2
6	A	1210	1PE	C24-C14-OH5-C25
4	A	1207	OLC	C5-C6-C7-C8

There are no ring outliers.


4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1203	OLC	1	0
4	A	1213	OLC	2	0
5	A	1204	CLR	1	0
5	A	1205	CLR	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand EZX A 1202







Ligand OLC A 1213



Ligand OLC A 1214







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	355/376 (94%)	0.50	25 (7%) 22 19	40, 66, 102, 126	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1068	LEU	4.3
1	A	180	PRO	3.4
1	A	1003	LEU	3.2
1	A	188	SER	3.1
1	A	74	LYS	3.1
1	A	1066	ASP	3.1
1	A	336	TRP	3.1
1	A	163	ILE	3.0
1	A	1098	ILE	2.7
1	A	243	PHE	2.7
1	A	1101	TYR	2.6
1	A	1037	ALA	2.6
1	A	1096	THR	2.5
1	A	1006	ASN	2.5
1	A	264	GLU	2.5
1	A	401	GLU	2.3
1	A	76	TRP	2.3
1	A	112	PHE	2.3
1	A	84	VAL	2.2
1	A	179	ASN	2.2
1	A	164	TRP	2.2
1	A	375	VAL	2.2
1	A	120	ASP	2.1
1	A	240	PHE	2.1
1	A	245	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	1PE	A	1210	16/16	0.76	0.18	72,113,121,123	0
4	OLC	A	1214	16/25	0.78	0.21	70,89,131,169	0
4	OLC	A	1208	13/25	0.79	0.15	65,93,118,127	0
4	OLC	A	1207	16/25	0.80	0.26	60,81,98,99	0
4	OLC	A	1209	14/25	0.81	0.15	68,85,116,117	0
4	OLC	A	1206	16/25	0.82	0.18	62,76,110,112	0
4	OLC	A	1213	16/25	0.82	0.21	65,79,122,127	0
7	PEG	A	1211	7/7	0.83	0.13	70,92,98,101	0
5	CLR	A	1205	20/28	0.84	0.17	86,94,113,119	0
3	EZX	A	1202	28/28	0.86	0.13	62,74,85,92	0
5	CLR	A	1204	24/28	0.87	0.16	72,106,129,141	0
4	OLC	A	1203	16/25	0.87	0.17	45,77,92,98	0
7	PEG	A	1212	7/7	0.90	0.13	68,77,84,96	0
2	MG	A	1201	1/1	0.99	0.05	20,20,20,20	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OLC A 1214:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 1208:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 1207:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around OLC A 1209:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around OLC A 1206:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around OLC A 1213:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around EZX A 1202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around CLR A 1204:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.