



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 24, 2026 – 11:12 PM EDT

PDB ID : 1JU4 / pdb_00001ju4
Title : BACTERIAL COCAINE ESTERASE COMPLEX WITH PRODUCT
Authors : Larsen, N.A.; Turner, J.M.; Stevens, J.; Rosser, S.J.; Basran, A.; Lerner, R.A.;
Bruce, N.C.; Wilson, I.A.
Deposited on : 2001-08-23
Resolution : 1.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	583	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BEZ	A	584	-	X	-	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4796 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called cocaine esterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	570	4351	2751	741	845	14	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	575	PRO	-	cloning artifact	UNP Q9L9D7
A	576	LEU	-	cloning artifact	UNP Q9L9D7
A	577	GLU	-	cloning artifact	UNP Q9L9D7
A	578	HIS	-	expression tag	UNP Q9L9D7
A	579	HIS	-	expression tag	UNP Q9L9D7
A	580	HIS	-	expression tag	UNP Q9L9D7
A	581	HIS	-	expression tag	UNP Q9L9D7
A	582	HIS	-	expression tag	UNP Q9L9D7
A	583	HIS	-	expression tag	UNP Q9L9D7

- Molecule 2 is BENZOIC ACID (CCD ID: BEZ) (formula: C₇H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			9	7	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	436	Total	O	0	0
			436	436		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: cocaine esterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	106.22Å 106.22Å 222.21Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.63 20.00 – 1.63	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-1.63) 94.5 (20.00-1.63)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.60Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.190 , 0.219 0.182 , 0.180	Depositor DCC
R_{free} test set	4390 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	16.2	Xtriage
Anisotropy	0.168	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4796	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BEZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/4459	1.52	48/6100 (0.8%)

There are no bond length outliers.

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	539	GLY	CA-C-N	15.06	139.60	121.83
1	A	539	GLY	C-N-CA	15.06	139.60	121.83
1	A	292	GLY	CA-C-N	14.48	142.03	122.77
1	A	292	GLY	C-N-CA	14.48	142.03	122.77
1	A	536	SER	CA-C-N	13.67	141.47	122.19
1	A	536	SER	C-N-CA	13.67	141.47	122.19
1	A	87	HIS	ND1-CE1-NE2	8.85	117.25	108.40
1	A	292	GLY	O-C-N	-8.03	111.92	122.28
1	A	236	GLN	CB-CG-CD	7.73	125.74	112.60
1	A	555	ASN	OD1-CG-ND2	-7.68	114.92	122.60
1	A	454	SER	CA-C-N	7.21	135.61	122.92
1	A	454	SER	C-N-CA	7.21	135.61	122.92
1	A	459	ASP	CA-CB-CG	7.09	119.69	112.60
1	A	539	GLY	O-C-N	-7.05	113.44	122.39
1	A	323	ARG	CD-NE-CZ	6.75	133.85	124.40
1	A	292	GLY	CA-C-O	6.61	126.04	119.04
1	A	490	VAL	CA-C-N	6.60	131.14	123.15
1	A	490	VAL	C-N-CA	6.60	131.14	123.15
1	A	535	ASN	OD1-CG-ND2	-6.38	116.22	122.60
1	A	409	HIS	CA-C-O	-6.28	113.51	120.60
1	A	176	THR	CA-C-N	-6.21	112.29	122.54
1	A	176	THR	C-N-CA	-6.21	112.29	122.54
1	A	422	HIS	N-CA-C	6.16	118.80	111.71
1	A	87	HIS	CE1-NE2-CD2	-6.16	102.84	109.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	275	ASN	CA-CB-CG	-6.05	106.55	112.60
1	A	454	SER	CA-C-O	5.97	126.75	119.11
1	A	535	ASN	CB-CG-OD1	5.83	132.46	120.80
1	A	536	SER	O-C-N	-5.71	114.10	122.43
1	A	491	ASN	CA-C-O	5.66	128.26	121.02
1	A	41	ARG	CD-NE-CZ	5.60	132.24	124.40
1	A	539	GLY	CA-C-O	5.51	125.14	119.02
1	A	533	ASP	CA-CB-CG	5.47	118.07	112.60
1	A	151	TRP	CD1-NE1-CE2	5.40	118.61	108.90
1	A	151	TRP	CG-CD1-NE1	-5.34	103.26	110.20
1	A	72	GLN	OE1-CD-NE2	5.21	127.81	122.60
1	A	230	ASP	CA-CB-CG	5.21	117.81	112.60
1	A	376	GLY	N-CA-C	5.20	117.44	111.36
1	A	189	PHE	CA-C-N	5.18	127.28	120.60
1	A	189	PHE	C-N-CA	5.18	127.28	120.60
1	A	304	THR	CA-C-O	-5.15	115.52	121.19
1	A	224	GLN	OE1-CD-NE2	-5.14	117.46	122.60
1	A	408	PHE	CA-CB-CG	-5.12	108.68	113.80
1	A	168	ALA	CA-C-O	5.10	125.96	120.55
1	A	175	ILE	CB-CA-C	-5.08	105.38	111.88
1	A	97	THR	CA-CB-OG1	-5.06	102.01	109.60
1	A	151	TRP	CB-CG-CD1	-5.06	119.31	126.90
1	A	219	PRO	CA-C-N	5.01	127.25	120.38
1	A	219	PRO	C-N-CA	5.01	127.25	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4351	0	4192	15	0
2	A	9	0	5	0	0
3	A	436	0	0	3	0
All	All	4796	0	4197	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:452:SER:OG	1:A:558:HIS:HE1	1.88	0.57
1:A:65:ASP:OD2	1:A:315:LYS:HD2	2.07	0.55
1:A:172:THR:HG23	1:A:189:PHE:CE2	2.46	0.51
1:A:546:GLN:OE1	1:A:548:GLU:OE2	2.29	0.51
1:A:161:GLU:HG3	3:A:2425:HOH:O	2.10	0.50
1:A:179:SER:O	1:A:180:ASP:HB2	2.13	0.48
1:A:422:HIS:NE2	1:A:521:MET:HE1	2.29	0.47
1:A:368:ALA:HB2	3:A:2311:HOH:O	2.15	0.47
1:A:351:TRP:CH2	1:A:443:GLY:HA2	2.51	0.46
1:A:538:THR:OG1	1:A:545:GLU:OE1	2.30	0.45
1:A:546:GLN:O	1:A:549:GLU:HG2	2.17	0.45
1:A:448:ARG:NH1	1:A:502:GLU:OE2	2.50	0.45
1:A:310:ALA:O	1:A:314:HIS:HD2	2.00	0.44
1:A:5:ASN:N	3:A:2130:HOH:O	2.50	0.44
1:A:354:PRO:O	1:A:355:ASP:HB2	2.18	0.43

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	568/583 (97%)	549 (97%)	19 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	462/475 (97%)	452 (98%)	10 (2%)	45 18

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	ASP
1	A	175	ILE
1	A	236	GLN
1	A	240	LEU
1	A	252	LEU
1	A	301	ILE
1	A	383	SER
1	A	387	SER
1	A	448	ARG
1	A	535	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	GLN
1	A	104	GLN
1	A	224	GLN
1	A	236	GLN
1	A	314	HIS
1	A	409	HIS
1	A	535	ASN
1	A	558	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BEZ	A	584	-	9,9,9	2.99	5 (55%)	11,11,11	3.11	5 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEZ	A	584	-	-	0/4/4/4	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	584	BEZ	C3-C2	6.42	1.49	1.38
2	A	584	BEZ	C6-C1	4.43	1.46	1.39
2	A	584	BEZ	O2-C	-2.46	1.23	1.30
2	A	584	BEZ	C5-C6	2.22	1.42	1.38
2	A	584	BEZ	C4-C3	2.19	1.43	1.38

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	584	BEZ	C3-C2-C1	-6.29	114.18	120.36
2	A	584	BEZ	C4-C5-C6	-4.51	114.67	120.24
2	A	584	BEZ	C2-C1-C	-3.56	113.33	120.37
2	A	584	BEZ	C5-C6-C1	3.54	123.85	120.36
2	A	584	BEZ	C6-C1-C2	2.83	122.16	118.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	570/583 (97%)	0.05	22 (3%)	43 48	12, 19, 38, 81	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	175	ILE	6.0
1	A	179	SER	5.4
1	A	176	THR	4.7
1	A	180	ASP	4.6
1	A	32	ASP	4.4
1	A	31	ALA	3.6
1	A	189	PHE	3.6
1	A	30	ASP	3.5
1	A	174	LEU	3.0
1	A	178	ARG	2.7
1	A	326	THR	2.5
1	A	366	SER	2.5
1	A	183	PRO	2.5
1	A	386	GLU	2.4
1	A	382	ILE	2.4
1	A	173	GLY	2.4
1	A	327	ASP	2.4
1	A	354	PRO	2.2
1	A	181	ALA	2.2
1	A	287	HIS	2.2
1	A	187	ALA	2.1
1	A	330	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BEZ	A	584	9/9	0.97	0.04	14,18,21,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.