



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:37 PM UTC

PDB ID : 4JUZ / pdb_00004juz
Title : Ternary complex of gamma-OHPDG adduct modified dna (zero primer) with dna polymerase iv and incoming dgtp
Authors : Banerjee, S.; Shanmugam, G.; Stone, M.P.
Deposited on : 2013-03-25
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	 73% 22% . .
2	B	15	 73% 20% 7%
3	C	14	 64% 36%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3411 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	0	0
			2744	1760	472	505	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q97W02
A	-4	HIS	-	expression tag	UNP Q97W02
A	-3	HIS	-	expression tag	UNP Q97W02
A	-2	HIS	-	expression tag	UNP Q97W02
A	-1	HIS	-	expression tag	UNP Q97W02
A	0	HIS	-	expression tag	UNP Q97W02


- Molecule 2 is a DNA chain called DNA (5'-D(P*CP*(KAG)P*GP*AP*AP*TP*CP*CP*TP*TP*CP*CP*CP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	15	Total	C	N	O	P	0	0	0
			302	145	50	92	15			

- Molecule 3 is a DNA chain called DNA (5'-D(*GP*GP*GP*GP*GP*AP*AP*GP*GP*AP*TP*TP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	14	Total	C	N	O	P	0	0	0
			292	138	60	81	13			

- Molecule 4 is 2'-DEOXYGUANOSINE-5'-TRIPHOSPHATE (CCD ID: DGT) (formula: C₁₀H₁₆N₅O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Ca	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	24	Total	O	0	0
			24	24		
6	B	10	Total	O	0	0
			10	10		
6	C	6	Total	O	0	0
			6	6		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase IV



- Molecule 2: DNA (5'-D(P*CP*(KAG)P*GP*AP*AP*TP*CP*CP*TP*TP*CP*CP*CP*CP*C)-3')



- Molecule 3: DNA (5'-D(*GP*GP*GP*GP*GP*AP*AP*GP*GP*AP*TP*TP*CP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	94.91Å 104.36Å 52.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.04 – 2.65 42.04 – 2.65	Depositor EDS
% Data completeness (in resolution range)	98.6 (42.04-2.65) 94.2 (42.04-2.65)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.65Å)	Xtriage
Refinement program	PHENIX 1.7.2 _869	Depositor
R, R_{free}	0.204 , 0.269 0.196 , 0.258	Depositor DCC
R_{free} test set	1549 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	56.2	Xtriage
Anisotropy	0.272	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 28.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3411	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DGT, CA, KAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/2783	1.00	4/3736 (0.1%)
2	B	0.52	0/305	1.32	1/463 (0.2%)
3	C	0.42	0/329	1.22	1/508 (0.2%)
All	All	0.64	0/3417	1.06	6/4707 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	11	PHE	N-CA-C	6.95	119.44	111.11
1	A	293	LEU	N-CA-C	6.52	120.51	111.90
3	C	27	DG	C4'-C3'-O3'	-5.43	101.85	110.00
1	A	74	LEU	CA-C-N	-5.39	114.42	120.14
1	A	74	LEU	C-N-CA	-5.39	114.42	120.14
2	B	6	DG	O5'-C5'-C4'	-5.26	102.91	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2744	0	2889	48	0
2	B	302	0	171	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	292	0	158	5	0
4	A	31	0	12	4	0
5	A	2	0	0	0	0
6	A	24	0	0	5	1
6	B	10	0	0	1	0
6	C	6	0	0	3	1
All	All	3411	0	3230	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:4:DC:H2''	2:B:5:KAG:H5''	1.60	0.83
1:A:36:ARG:NH2	1:A:254:ASN:OD1	2.13	0.80
1:A:106:GLU:OE1	6:A:1101:HOH:O	1.99	0.78
3:C:28:DA:N3	6:C:101:HOH:O	2.19	0.75
3:C:23:DG:OP2	6:C:106:HOH:O	2.04	0.73
1:A:51:ARG:NH2	4:A:1000:DGT:O1B	2.21	0.73
1:A:273:TYR:HA	1:A:276:LEU:HD12	1.70	0.73
1:A:79:GLU:H	1:A:79:GLU:CD	1.99	0.69
1:A:2:ILE:O	6:A:1121:HOH:O	2.11	0.68
1:A:112:SER:N	6:A:1121:HOH:O	2.31	0.62
1:A:78:LYS:NZ	2:B:5:KAG:H2B	2.15	0.62
1:A:242:ARG:HH12	2:B:7:DA:H5''	1.66	0.60
2:B:4:DC:H6	2:B:4:DC:H5''	1.67	0.59
1:A:95:TYR:HD2	1:A:124:LEU:HD11	1.68	0.58
1:A:168:ASP:N	1:A:168:ASP:OD1	2.34	0.58
1:A:199:ILE:HD11	1:A:208:ILE:HG13	1.86	0.57
3:C:23:DG:H2'	3:C:24:DA:C8	2.39	0.57
1:A:199:ILE:HG23	1:A:204:ASP:HB2	1.86	0.57
1:A:289:VAL:HG22	1:A:295:ILE:HG12	1.90	0.54
1:A:105:ASP:OD1	4:A:1000:DGT:H5'	2.08	0.53
1:A:111:ILE:N	6:A:1121:HOH:O	2.43	0.52
1:A:289:VAL:HB	1:A:332:ARG:HB2	1.92	0.52
4:A:1000:DGT:H8	3:C:32:DC:H2''	1.92	0.52
1:A:144:ILE:HB	1:A:165:VAL:HG22	1.92	0.52
1:A:67:ILE:HG22	1:A:68:LEU:HG	1.92	0.50
1:A:242:ARG:NH1	2:B:7:DA:H5''	2.26	0.50
4:A:1000:DGT:O1A	6:A:1101:HOH:O	2.20	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:87:ARG:NE	1:A:136:GLU:OE2	2.33	0.49
1:A:95:TYR:CD2	1:A:124:LEU:HD11	2.46	0.49
1:A:180:ILE:HD13	1:A:194:LEU:HD13	1.95	0.48
1:A:230:ARG:O	1:A:232:GLU:HG3	2.13	0.48
1:A:167:ASP:OD2	1:A:169:GLU:HG2	2.14	0.47
1:A:99:ILE:HD12	1:A:109:LEU:HD21	1.96	0.47
1:A:156:ASP:HA	1:A:159:LYS:HE3	1.96	0.47
1:A:204:ASP:N	1:A:204:ASP:OD1	2.49	0.46
1:A:270:GLU:OE2	1:A:312:TYR:OH	2.21	0.46
1:A:98:LYS:HB2	1:A:110:ASP:HB3	1.99	0.45
1:A:173:ARG:O	1:A:177:GLU:HB2	2.16	0.45
2:B:4:DC:O5'	6:B:103:HOH:O	2.20	0.45
2:B:4:DC:H5''	2:B:4:DC:C6	2.49	0.45
1:A:100:GLU:OE1	1:A:238:ARG:HD2	2.17	0.44
1:A:210:PHE:HE2	1:A:214:LYS:HE3	1.82	0.44
1:A:12:TYR:OH	1:A:78:LYS:HE3	2.17	0.44
1:A:115:VAL:HG22	1:A:120:GLU:HB3	2.00	0.44
1:A:100:GLU:HB2	1:A:237:ILE:HG23	2.00	0.43
3:C:23:DG:H5''	6:C:106:HOH:O	2.19	0.43
1:A:276:LEU:O	1:A:279:ARG:HB2	2.19	0.42
1:A:9:ASP:O	1:A:10:TYR:C	2.61	0.42
1:A:173:ARG:NH2	1:A:177:GLU:HG2	2.35	0.42
1:A:225:LEU:HD23	1:A:225:LEU:HA	1.73	0.41
1:A:27:PRO:HB2	1:A:50:ALA:HB2	2.02	0.41
1:A:78:LYS:HZ2	2:B:5:KAG:H2B	1.82	0.41
1:A:257:ASN:HB3	1:A:260:GLU:HB2	2.02	0.41
1:A:78:LYS:HZ1	2:B:5:KAG:H2B	1.83	0.41
1:A:166:ILE:HG23	1:A:170:GLU:HG2	2.03	0.41
1:A:221:LYS:HB2	1:A:221:LYS:HE2	1.81	0.41
1:A:251:MET:HA	1:A:264:TYR:CE1	2.56	0.40
1:A:260:GLU:O	1:A:263:PRO:HD2	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1108:HOH:O	6:C:104:HOH:O[3_547]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/347 (98%)	319 (94%)	19 (6%)	1 (0%)	36	52

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	277	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/306 (98%)	271 (90%)	29 (10%)	8	12

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	26	LYS
1	A	63	GLU
1	A	79	GLU
1	A	115	VAL
1	A	119	ARG
1	A	131	LYS
1	A	138	ILE
1	A	168	ASP
1	A	173	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	177	GLU
1	A	178	LEU
1	A	192	GLU
1	A	195	LYS
1	A	199	ILE
1	A	200	ASN
1	A	204	ASP
1	A	211	ASP
1	A	238	ARG
1	A	241	VAL
1	A	242	ARG
1	A	247	ARG
1	A	277	ASP
1	A	278	LYS
1	A	279	ARG
1	A	294	ASP
1	A	324	GLU
1	A	325	GLU
1	A	327	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	KAG	B	5	2,3	24,28,30	2.76	10 (41%)	32,39,44	2.39	14 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	KAG	B	5	2,3	-	2/11/26/29	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	5	KAG	O6-C6	6.80	1.36	1.23
2	B	5	KAG	C2-N2	5.64	1.46	1.34
2	B	5	KAG	C2'-C3'	-5.56	1.38	1.52
2	B	5	KAG	O4'-C4'	-3.18	1.37	1.45
2	B	5	KAG	C3'-C4'	-3.18	1.44	1.53
2	B	5	KAG	O2G-C3G	2.91	1.36	1.20
2	B	5	KAG	C2-N3	2.85	1.37	1.32
2	B	5	KAG	O4'-C1'	-2.36	1.37	1.42
2	B	5	KAG	O5'-C5'	-2.27	1.37	1.44
2	B	5	KAG	C2-N1	-2.15	1.33	1.36

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	5	KAG	C1A-C2B-C3G	6.08	121.87	112.62
2	B	5	KAG	C2-N3-C4	5.65	119.07	112.00
2	B	5	KAG	C5-C4-N3	-4.21	121.69	128.39
2	B	5	KAG	O4'-C1'-N9	3.95	114.88	107.86
2	B	5	KAG	C6-C5-N7	3.36	136.40	130.29
2	B	5	KAG	C4-C5-N7	-2.86	106.14	110.67
2	B	5	KAG	N9-C8-N7	-2.62	108.53	113.40
2	B	5	KAG	C2-N1-C6	-2.41	121.64	124.55
2	B	5	KAG	C8-N7-C5	2.34	108.43	104.26
2	B	5	KAG	N9-C4-N3	2.32	130.60	125.95
2	B	5	KAG	C5'-C4'-C3'	-2.25	102.06	114.64
2	B	5	KAG	O6-C6-C5	-2.07	121.06	126.53
2	B	5	KAG	C5-C6-N1	2.05	118.47	113.25
2	B	5	KAG	C1'-N9-C4	-2.00	120.10	125.50

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	5	KAG	C3'-C4'-C5'-O5'
2	B	5	KAG	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	5	KAG	4	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	DGT	A	1000	5	32,33,33	1.84	8 (25%)	48,52,52	2.01	19 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DGT	A	1000	5	-	3/22/34/34	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1000	DGT	PB-O3A	-4.76	1.54	1.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1000	DGT	O6-C6	4.24	1.31	1.23
4	A	1000	DGT	C2-N2	3.73	1.42	1.34
4	A	1000	DGT	PB-O3B	-3.12	1.56	1.59
4	A	1000	DGT	C6-N1	-2.65	1.33	1.38
4	A	1000	DGT	C2'-C3'	-2.44	1.46	1.52
4	A	1000	DGT	C3'-C4'	-2.36	1.46	1.53
4	A	1000	DGT	PG-O3G	2.20	1.57	1.50

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1000	DGT	C5-C4-N3	-5.55	119.56	128.39
4	A	1000	DGT	C2-N3-C4	4.52	120.08	112.30
4	A	1000	DGT	N9-C4-N3	3.86	133.67	125.95
4	A	1000	DGT	O1A-PA-O3A	3.70	117.28	107.27
4	A	1000	DGT	O5'-C5'-C4'	3.21	119.94	108.99
4	A	1000	DGT	C8-N7-C5	2.87	109.38	104.26
4	A	1000	DGT	O3B-PB-O2B	2.79	119.09	110.70
4	A	1000	DGT	N9-C8-N7	-2.77	108.26	113.40
4	A	1000	DGT	C4-C5-N7	-2.69	106.41	110.67
4	A	1000	DGT	C1'-N9-C4	-2.69	118.26	125.50
4	A	1000	DGT	O3A-PA-O2A	-2.55	103.02	110.70
4	A	1000	DGT	C6-C5-N7	2.48	134.79	130.29
4	A	1000	DGT	O1B-PB-O3A	2.44	113.87	107.27
4	A	1000	DGT	C2-N1-C6	-2.36	120.83	125.11
4	A	1000	DGT	O4'-C1'-N9	2.36	112.05	107.86
4	A	1000	DGT	C5-C6-N1	2.26	119.01	113.25
4	A	1000	DGT	C1'-N9-C8	2.07	132.55	127.91
4	A	1000	DGT	O1G-PG-O3B	2.06	111.53	104.64
4	A	1000	DGT	O6-C6-C5	-2.01	121.23	126.53

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1000	DGT	PG-O3B-PB-O1B
4	A	1000	DGT	PB-O3A-PA-O2A
4	A	1000	DGT	PG-O3B-PB-O2B

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1000	DGT	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/347 (98%)	-0.19	0 100 100	35, 54, 76, 91	0
2	B	14/15 (93%)	-0.16	0 100 100	49, 62, 70, 73	0
3	C	14/14 (100%)	-0.31	0 100 100	50, 55, 77, 77	0
All	All	369/376 (98%)	-0.19	0 100 100	35, 55, 76, 91	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	KAG	B	5	26/28	0.90	0.12	50,58,67,68	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	CA	A	1002	1/1	0.82	0.09	77,77,77,77	0
5	CA	A	1001	1/1	0.93	0.07	58,58,58,58	0
4	DGT	A	1000	31/31	0.93	0.08	42,48,70,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.