



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:17 AM UTC

PDB ID : 6JU5 / pdb_00006ju5
Title : Aspergillus oryzae pro-tyrosinase C92A/F513Y mutant
Authors : Fujieda, N.; Umakoshi, K.; Nishikawa, Y.; Kurisu, G.; Itoh, S.
Deposited on : 2019-04-13
Resolution : 1.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2194 (1.36-1.32)
Clashscore	190562	2222 (1.36-1.32)
Ramachandran outliers	187476	2197 (1.36-1.32)
Sidechain outliers	187428	2197 (1.36-1.32)
RSRZ outliers	180081	2193 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	621	
1	B	621	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9515 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	542	Total	C	N	O	S	0	5	0
			4276	2762	726	776	12			
1	B	554	Total	C	N	O	S	0	4	0
			4402	2841	750	799	12			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP A0A1S9DK56
A	-2	PRO	-	expression tag	UNP A0A1S9DK56
A	-1	GLY	-	expression tag	UNP A0A1S9DK56
A	0	GLY	-	expression tag	UNP A0A1S9DK56
A	1	SER	-	expression tag	UNP A0A1S9DK56
A	92	ALA	CYS	engineered mutation	UNP A0A1S9DK56
A	513	TYR	PHE	engineered mutation	UNP A0A1S9DK56
A	513	TYR	PHE	microheterogeneity	UNP A0A1S9DK56
A	513	DAH	PHE	microheterogeneity	UNP A0A1S9DK56
B	-3	GLY	-	expression tag	UNP A0A1S9DK56
B	-2	PRO	-	expression tag	UNP A0A1S9DK56
B	-1	GLY	-	expression tag	UNP A0A1S9DK56
B	0	GLY	-	expression tag	UNP A0A1S9DK56
B	1	SER	-	expression tag	UNP A0A1S9DK56
B	92	ALA	CYS	engineered mutation	UNP A0A1S9DK56
B	513	TYR	PHE	engineered mutation	UNP A0A1S9DK56
B	513	TYR	PHE	microheterogeneity	UNP A0A1S9DK56
B	513	DAH	PHE	microheterogeneity	UNP A0A1S9DK56

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cu	0	2
			4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total 4	Cu 4	0	2

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	402	Total 408	O 408	0	7
3	B	412	Total 421	O 421	0	9

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.34Å 118.21Å 79.36Å 90.00° 91.13° 90.00°	Depositor
Resolution (Å)	30.00 – 1.34 30.00 – 1.34	Depositor EDS
% Data completeness (in resolution range)	99.9 (30.00-1.34) 99.8 (30.00-1.34)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 1.34Å)	Xtriage
Refinement program	SHELXL 2017/1	Depositor
R, R_{free}	0.157 , 0.202 (Not available) , 0.196	Depositor DCC
R_{free} test set	11180 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	12.5	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9515	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: DAH, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/4410	1.05	8/6016 (0.1%)
1	B	0.43	0/4529	1.01	4/6177 (0.1%)
All	All	0.43	0/8939	1.03	12/12193 (0.1%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	224	ASP	CA-CB-CG	8.01	120.61	112.60
1	A	372	HIS	CA-CB-CG	-7.89	105.91	113.80
1	B	372	HIS	CA-CB-CG	-7.64	106.16	113.80
1	A	36	ARG	CD-NE-CZ	6.66	133.72	124.40
1	A	125	ASN	CA-C-N	-6.59	116.98	123.04
1	A	125	ASN	C-N-CA	-6.59	116.98	123.04
1	B	281	ASP	CA-CB-CG	6.29	118.89	112.60
1	A	317	ALA	CA-C-N	5.65	125.76	119.32
1	A	317	ALA	C-N-CA	5.65	125.76	119.32
1	A	359	VAL	CB-CA-C	-5.57	108.41	113.70
1	A	264	HIS	CA-CB-CG	5.32	119.12	113.80
1	B	355	HIS	CA-CB-CG	-5.06	108.74	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4276	0	3925	23	0
1	B	4402	0	4078	28	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	408	0	0	3	0
3	B	421	0	0	6	0
All	All	9515	0	8003	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:359:VAL:HG22	1:B:513[A]:TYR:CZ	2.14	0.83
1:B:359:VAL:HG22	1:B:513[A]:TYR:CE2	2.14	0.82
1:B:284:PHE:CE1	1:B:288:THR:HG21	2.25	0.71
1:B:131:LYS:O	1:B:135:GLU:HG3	1.98	0.63
1:A:302:TYR:CE2	1:A:473:ARG:NH1	2.68	0.62
1:B:162:ASP:N	3:B:803:HOH:O	2.34	0.59
1:B:586:THR:HG22	3:B:837:HOH:O	2.05	0.56
1:A:240:GLU:O	1:A:241:ASN:HB2	2.05	0.56
1:A:27:LEU:O	1:A:31:SER:HB2	2.06	0.56
1:B:166[A]:LEU:HD22	1:B:367:ILE:CD1	2.36	0.56
1:A:547:TYR:CZ	1:A:551:GLN:HG3	2.42	0.55
1:A:267:TYR:OH	1:A:543:LEU:HB3	2.08	0.53
1:B:-3:GLY:HA2	3:B:808:HOH:O	2.07	0.53
1:B:284:PHE:O	1:B:288:THR:HG23	2.09	0.53
1:B:349:LYS:HD3	3:B:1081:HOH:O	2.10	0.53
1:A:346:HIS:O	1:A:349:LYS:NZ	2.42	0.52
1:A:74:TRP:CE3	1:A:75:ASP:HB2	2.46	0.51
1:A:496:LYS:HE2	1:A:611:ASP:OD2	2.10	0.50
1:B:556:VAL:O	1:B:556:VAL:HG13	2.11	0.50
1:A:333:ASN:HB2	1:A:511:PHE:CD2	2.47	0.50
1:A:207:ARG:HA	3:A:1152:HOH:O	2.13	0.49
1:B:166[A]:LEU:HD22	1:B:367:ILE:HD12	1.95	0.48
1:B:54:LEU:HD23	3:B:823:HOH:O	2.13	0.47
1:B:333:ASN:HB2	1:B:511:PHE:CD2	2.49	0.47
1:A:359:VAL:HG22	1:A:513[A]:TYR:CE2	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:339:MET:HE2	3:B:886:HOH:O	2.14	0.47
1:A:302:TYR:CZ	1:A:473:ARG:NH1	2.82	0.47
1:A:441:ALA:O	1:A:444:ILE:HG22	2.14	0.47
1:B:14:HIS:HE1	1:B:136:ASN:OD1	1.98	0.47
1:B:14:HIS:CE1	1:B:136:ASN:OD1	2.68	0.46
1:A:36:ARG:HG3	1:A:178:LEU:HG	1.97	0.46
1:B:240:GLU:O	1:B:241:ASN:CB	2.64	0.45
1:B:233:THR:HA	1:B:355:HIS:CE1	2.52	0.45
1:A:233:THR:HA	1:A:355:HIS:CE1	2.52	0.44
1:A:245:ASP:OD2	1:B:30:THR:OG1	2.32	0.44
1:A:333:ASN:HB2	1:A:511:PHE:CG	2.53	0.44
1:A:339:MET:HE2	3:A:825:HOH:O	2.17	0.44
1:B:409:ARG:HB3	1:B:420:PHE:CE1	2.54	0.43
1:A:1:SER:HA	1:A:2:PRO:C	2.42	0.43
1:A:359:VAL:HB	1:A:360:PRO:HD3	2.01	0.42
1:B:363:ALA:HA	1:B:368:PHE:CG	2.54	0.42
1:B:169:LEU:HD13	1:B:193:LEU:HD12	2.01	0.42
1:A:400:LYS:NZ	1:A:423:ASP:OD2	2.51	0.41
1:A:363:ALA:HA	1:A:368:PHE:CG	2.56	0.41
1:B:542:ARG:HD3	1:B:542:ARG:C	2.44	0.41
1:A:118:MET:HE1	1:A:141:TRP:CD2	2.56	0.41
3:A:1065:HOH:O	1:B:347:ASP:CB	2.68	0.41
1:B:479:LEU:HD12	1:B:479:LEU:HA	1.91	0.41
1:B:556:VAL:O	1:B:556:VAL:CG1	2.69	0.41
1:B:365:ASP:OD1	1:B:367:ILE:HG22	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	529/621 (85%)	516 (98%)	13 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	540/621 (87%)	533 (99%)	7 (1%)	0	100	100
All	All	1069/1242 (86%)	1049 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	424/536 (79%)	422 (100%)	2 (0%)	81	61
1	B	441/536 (82%)	440 (100%)	1 (0%)	87	72
All	All	865/1072 (81%)	862 (100%)	3 (0%)	86	70

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	ASN
1	A	328	HIS
1	B	328	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	ASN
1	A	241	ASN
1	A	333	ASN
1	A	371	HIS
1	B	50	GLN
1	B	199	GLN
1	B	280	GLN
1	B	371	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	DAH	A	513[B]	2,1	12,13,14	0.30	0	12,17,19	0.38	0
1	DAH	B	513[B]	2,1	12,13,14	0.24	0	12,17,19	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DAH	A	513[B]	2,1	-	1/5/6/8	0/1/1/1
1	DAH	B	513[B]	2,1	-	0/5/6/8	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	513[B]	DAH	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	541/621 (87%)	-0.06	21 (3%)	43 55	9, 17, 38, 69	4 (0%)
1	B	553/621 (89%)	-0.09	17 (3%)	51 62	7, 16, 35, 65	3 (0%)
All	All	1094/1242 (88%)	-0.07	38 (3%)	47 59	7, 17, 37, 69	7 (0%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	590	ALA	5.7
1	A	212	THR	5.2
1	A	32	ALA	5.1
1	B	223	PHE	5.0
1	B	222	SER	4.2
1	B	597	ASP	4.1
1	A	526	ALA	4.1
1	B	30	THR	4.0
1	A	223	PHE	4.0
1	A	272	GLY	3.6
1	B	224	ASP	3.4
1	A	33	GLY	3.2
1	B	-2	PRO	3.2
1	B	212	THR	3.1
1	B	89	THR	3.0
1	A	222	SER	3.0
1	A	317	ALA	3.0
1	B	-1	GLY	3.0
1	A	181	GLY	2.9
1	A	273	TRP	2.8
1	A	398	VAL	2.8
1	B	514	SER	2.7
1	B	0	GLY	2.7
1	A	31	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	35	ASN	2.6
1	B	556	VAL	2.5
1	A	183	PRO	2.5
1	B	153	ARG	2.4
1	A	89	THR	2.4
1	B	88	PRO	2.4
1	A	441	ALA	2.4
1	B	317	ALA	2.3
1	A	453	GLY	2.3
1	A	598	ASP	2.3
1	A	182	GLN	2.2
1	A	34	SER	2.0
1	A	387	SER	2.0
1	B	441	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	DAH	A	513[B]	13/14	-	-	12,13,17,20	13
1	DAH	B	513[B]	13/14	-	-	10,13,18,20	13

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CU	B	701[A]	1/1	0.99	0.04	16,16,16,16	1
2	CU	B	701[B]	1/1	0.99	0.04	17,17,17,17	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CU	A	702[A]	1/1	1.00	0.02	11,11,11,11	1
2	CU	A	702[B]	1/1	1.00	0.02	13,13,13,13	1
2	CU	A	701[A]	1/1	1.00	0.01	15,15,15,15	1
2	CU	A	701[B]	1/1	1.00	0.01	17,17,17,17	1
2	CU	B	702[A]	1/1	1.00	0.01	12,12,12,12	1
2	CU	B	702[B]	1/1	1.00	0.01	13,13,13,13	1

6.5 Other polymers [i](#)

There are no such residues in this entry.