



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 10:43 AM UTC

PDB ID : 7JU5 / pdb_00007ju5
Title : Structure of RET protein tyrosine kinase in complex with pralsetinib
Authors : Terzyan, S.S.; Shen, T.; Wu, J.; Mooers, B.H.M.
Deposited on : 2020-08-19
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	314	5% 76% 14% • 9%
1	B	314	4% 79% 12% • 8%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5106 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase receptor Ret.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	287	Total	C	N	O	S	0	5	0
			2314	1487	405	406	16			
1	B	290	Total	C	N	O	S	0	14	0
			2438	1557	425	442	14			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	700	GLY	-	expression tag	UNP P07949
A	701	PRO	-	expression tag	UNP P07949
A	702	LEU	-	expression tag	UNP P07949
A	703	SER	-	expression tag	UNP P07949
A	704	LEU	-	expression tag	UNP P07949
B	700	GLY	-	expression tag	UNP P07949
B	701	PRO	-	expression tag	UNP P07949
B	702	LEU	-	expression tag	UNP P07949
B	703	SER	-	expression tag	UNP P07949
B	704	LEU	-	expression tag	UNP P07949

- Molecule 2 is Pralsetinib (CCD ID: Q4J) (formula: C₂₇H₃₂FN₉O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 39	C 27	F 1	N 9	O 2	0	0
2	B	1	Total 39	C 27	F 1	N 9	O 2	0	0

- Molecule 3 is FORMIC ACID (CCD ID: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 3	C 1	O 2	0	0
3	A	1	Total 3	C 1	O 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	127	Total	O	0	0
			127	127		
4	B	137	Total	O	0	0
			137	137		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase receptor Ret



- Molecule 1: Proto-oncogene tyrosine-protein kinase receptor Ret



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.45Å 80.33Å 79.84Å 90.00° 100.25° 90.00°	Depositor
Resolution (Å)	78.57 – 1.90 78.57 – 1.90	Depositor EDS
% Data completeness (in resolution range)	94.9 (78.57-1.90) 95.1 (78.57-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.188 , 0.247 (Not available) , 0.242	Depositor DCC
R_{free} test set	2392 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	25.1	Xtriage
Anisotropy	0.467	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5106	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: Q4J, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	1/2372 (0.0%)	1.08	4/3191 (0.1%)
1	B	1.08	2/2499 (0.1%)	1.07	3/3365 (0.1%)
All	All	1.07	3/4871 (0.1%)	1.08	7/6556 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	952	TYR	C-O	7.41	1.27	1.23
1	A	797	ASP	CA-C	-7.39	1.44	1.52
1	B	906	VAL	C-O	-6.29	1.17	1.24

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	778	VAL	N-CA-C	8.45	119.25	110.72
1	B	991	GLU	CA-C-N	-7.59	111.90	119.56
1	B	991	GLU	C-N-CA	-7.59	111.90	119.56
1	A	797	ASP	CB-CA-C	-5.87	102.62	110.79
1	B	782	VAL	N-CA-C	5.13	116.07	108.62
1	A	955	ILE	CA-C-N	-5.06	115.17	120.38
1	A	955	ILE	C-N-CA	-5.06	115.17	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2314	0	2366	35	0
1	B	2438	0	2459	27	0
2	A	39	0	0	1	0
2	B	39	0	0	0	0
3	A	6	0	2	1	0
3	B	6	0	2	1	0
4	A	127	0	0	11	0
4	B	137	0	0	6	0
All	All	5106	0	4829	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:709:PHE:CZ	1:A:800:LEU:HD13	2.19	0.78
1:A:852:ILE:HG22	1:A:1009:MET:HE2	1.77	0.66
1:A:771:ASP:OD2	1:A:912:ARG:NH2	2.32	0.63
1:A:910:GLN:HG3	1:B:770:ARG:HD2	1.80	0.61
1:A:713:GLU:O	1:A:715:PRO:HD3	2.00	0.61
1:A:789:LYS:HE2	1:A:791:TYR:CE1	2.37	0.59
1:A:959:ARG:NH1	1:B:798:GLY:O	2.36	0.58
1:A:847:THR:O	1:A:850:ASP:HB2	2.03	0.58
1:A:850:ASP:O	1:A:853:SER:HB3	2.04	0.57
1:B:878:ARG:NH1	4:B:1201:HOH:O	2.37	0.56
1:A:737:LYS:HE2	1:A:759[A]:MET:HE2	1.88	0.56
1:B:991:GLU:HG3	4:B:1265:HOH:O	2.07	0.54
1:A:728:LYS:HD3	4:A:1321:HOH:O	2.09	0.53
1:B:896[B]:SER:O	1:B:897:ARG:NH1	2.41	0.52
1:A:715:PRO:HB2	4:A:1290:HOH:O	2.10	0.52
1:A:763:ASN:HB2	4:A:1274:HOH:O	2.10	0.51
1:A:782:VAL:HG23	4:A:1260:HOH:O	2.10	0.50
1:A:746:LEU:HD11	1:A:791:TYR:HB3	1.92	0.50
1:A:715:PRO:CB	4:A:1290:HOH:O	2.59	0.50
1:A:709:PHE:CE2	1:A:800:LEU:HD13	2.46	0.50
1:B:817:ARG:NH1	1:B:948:GLY:O	2.46	0.49
1:B:907:LYS:HG2	4:B:1207:HOH:O	2.13	0.49
1:B:813[A]:ARG:NH2	1:B:950:ASN:OD1	2.44	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:986:GLN:O	1:A:989:LYS:HG2	2.12	0.48
1:A:959:ARG:HD3	4:A:1322:HOH:O	2.13	0.48
1:B:746:LEU:O	1:B:747:LYS:C	2.57	0.47
1:A:853:SER:HA	1:A:1009:MET:HE1	1.95	0.47
2:A:1101:Q4J:CAA	2:A:1101:Q4J:N1	2.74	0.47
1:B:713:GLU:HB2	1:B:796:GLN:OE1	2.14	0.46
1:A:873:ARG:HD3	1:A:895:LEU:O	2.16	0.46
1:A:959:ARG:CD	4:A:1322:HOH:O	2.63	0.46
1:A:770:ARG:CD	4:A:1261:HOH:O	2.63	0.45
1:B:902:GLU:HA	1:B:902:GLU:OE1	2.17	0.45
1:B:907:LYS:N	4:B:1207:HOH:O	2.49	0.45
1:B:823:GLY:HA2	1:B:824:PRO:C	2.42	0.45
1:B:929:THR:OG1	1:B:931[B]:GLN:HB2	2.16	0.45
1:B:843:GLU:HG3	4:B:1217:HOH:O	2.16	0.45
1:B:878:ARG:NH1	1:B:914:PRO:HG2	2.33	0.44
1:B:728:LYS:NZ	3:B:1103:FMT:H	2.33	0.44
1:A:896[A]:SER:O	1:A:897:ARG:NH1	2.49	0.43
1:B:826:TYR:OH	1:B:848:MET:HE2	2.18	0.43
1:B:747:LYS:HE3	1:B:791:TYR:O	2.18	0.43
3:A:1103:FMT:H	1:B:923:LEU:O	2.19	0.43
1:A:713:GLU:HG3	1:A:714:ASP:N	2.34	0.42
1:A:878:ARG:NH1	4:A:1202:HOH:O	2.51	0.42
1:A:776:PHE:HB2	1:A:802:LEU:HD21	2.02	0.42
1:B:871[A]:VAL:HG13	1:B:873:ARG:HG3	2.00	0.42
1:B:979:GLU:HB2	4:B:1311:HOH:O	2.20	0.42
1:B:952:TYR:N	1:B:953:PRO:CD	2.83	0.42
1:A:762:GLU:HA	1:A:762:GLU:OE1	2.20	0.42
1:A:770:ARG:NH2	1:A:771:ASP:OD1	2.52	0.42
1:B:905:TYR:OH	1:B:907:LYS:HD3	2.20	0.42
1:A:899:VAL:HG12	4:A:1303:HOH:O	2.19	0.42
1:A:868[B]:MET:HE2	1:A:868[B]:MET:HB3	1.63	0.41
1:A:911:GLY:HA3	4:A:1280:HOH:O	2.20	0.40
1:A:745:HIS:HA	1:A:749:ARG:O	2.21	0.40
1:B:813[A]:ARG:CZ	1:B:817:ARG:HD2	2.51	0.40
1:B:986:GLN:HB3	1:B:996:PRO:HD3	2.03	0.40
1:A:733:GLY:O	1:A:761:LYS:NZ	2.48	0.40
1:B:901:GLU:O	1:B:902:GLU:CB	2.70	0.40
1:A:709:PHE:CE2	1:A:795:SER:CB	3.05	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/314 (91%)	281 (98%)	5 (2%)	0	100	100
1	B	300/314 (96%)	290 (97%)	9 (3%)	1 (0%)	36	29
All	All	586/628 (93%)	571 (97%)	14 (2%)	1 (0%)	43	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	902	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/272 (92%)	241 (96%)	9 (4%)	31	23
1	B	266/272 (98%)	261 (98%)	5 (2%)	50	47
All	All	516/544 (95%)	502 (97%)	14 (3%)	38	34

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	768	GLU
1	A	781	GLN
1	A	782	VAL
1	A	789	LYS
1	A	800	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	926	HIS
1	A	945	VAL
1	A	959	ARG
1	A	1004	ASP
1	B	711	ILE
1	B	737	LYS
1	B	769	LEU
1	B	884	GLU
1	B	901	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	860	GLN
1	A	863	GLN
1	B	910	GLN
1	B	962	ASN
1	B	986	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FMT	B	1102	-	2,2,2	1.11	0	1,1,1	0.61	0
3	FMT	A	1102	-	2,2,2	0.97	0	1,1,1	0.43	0
3	FMT	A	1103	-	2,2,2	1.21	0	1,1,1	0.06	0
3	FMT	B	1103	-	2,2,2	0.88	0	1,1,1	0.93	0
2	Q4J	A	1101	-	39,43,43	2.55	14 (35%)	47,62,62	3.09	26 (55%)
2	Q4J	B	1101	-	39,43,43	2.25	12 (30%)	47,62,62	3.46	22 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	Q4J	A	1101	-	-	6/29/41/41	0/5/5/5
2	Q4J	B	1101	-	-	6/29/41/41	1/5/5/5

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	Q4J	NAC-NAD	-9.28	1.17	1.36
2	B	1101	Q4J	NBG-NBH	-7.67	1.25	1.36
2	B	1101	Q4J	CAZ-CAY	-5.05	1.38	1.52
2	B	1101	Q4J	CBI-CBJ	-4.20	1.32	1.39
2	A	1101	Q4J	C6-NAF	-4.17	1.30	1.40
2	A	1101	Q4J	CBF-NBE	3.87	1.42	1.34
2	A	1101	Q4J	CAA-CAB	-3.83	1.29	1.37
2	B	1101	Q4J	CBD-NBG	-3.81	1.34	1.42
2	A	1101	Q4J	CBI-CBJ	-3.74	1.33	1.39
2	A	1101	Q4J	CBM-CAB	3.67	1.57	1.49
2	B	1101	Q4J	C6-NAF	-3.66	1.32	1.40
2	A	1101	Q4J	CAE-NAD	-3.42	1.29	1.33
2	A	1101	Q4J	CBD-NBG	-3.29	1.35	1.42
2	A	1101	Q4J	NBG-NBH	-3.25	1.32	1.36
2	A	1101	Q4J	CAA-CAE	-3.23	1.34	1.40
2	B	1101	Q4J	CBF-NBE	3.12	1.40	1.34
2	A	1101	Q4J	CAZ-CAY	-2.92	1.44	1.52
2	B	1101	Q4J	CAM-C4	-2.88	1.43	1.50
2	B	1101	Q4J	CAO-CAN	-2.77	1.46	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	Q4J	CAS-CAN	-2.63	1.46	1.53
2	A	1101	Q4J	CAP-CAQ	2.50	1.56	1.53
2	A	1101	Q4J	CAB-NAC	-2.40	1.30	1.34
2	B	1101	Q4J	CAA-CAE	-2.21	1.36	1.40
2	B	1101	Q4J	CAP-CAQ	2.20	1.56	1.53
2	B	1101	Q4J	FBL-CBJ	-2.13	1.30	1.35
2	B	1101	Q4J	CAY-NAW	2.10	1.51	1.47

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1101	Q4J	CBI-NBH-NBG	9.93	112.09	104.19
2	B	1101	Q4J	CAE-NAD-NAC	9.73	110.78	103.71
2	B	1101	Q4J	CAA-CAE-NAD	-8.59	106.00	112.14
2	A	1101	Q4J	CAE-NAD-NAC	7.60	109.24	103.71
2	A	1101	Q4J	CBI-NBH-NBG	7.23	109.95	104.19
2	B	1101	Q4J	C2-N1-C6	6.55	123.89	115.67
2	B	1101	Q4J	CBJ-CBI-NBH	-6.16	104.91	111.94
2	A	1101	Q4J	CBK-NBG-NBH	-5.59	106.14	112.14
2	A	1101	Q4J	C2-N1-C6	5.19	122.19	115.67
2	A	1101	Q4J	CAA-CAE-NAD	-4.97	108.58	112.14
2	A	1101	Q4J	CBJ-CBI-NBH	-4.92	106.32	111.94
2	A	1101	Q4J	NBE-CBD-NBG	4.87	122.10	114.84
2	B	1101	Q4J	N3-C2-N1	-4.81	119.40	126.15
2	A	1101	Q4J	N3-C2-N1	-4.59	119.71	126.15
2	A	1101	Q4J	CAR-CAS-CAN	4.28	115.53	111.46
2	B	1101	Q4J	CAZ-CBF-NBE	-3.97	119.64	124.47
2	B	1101	Q4J	CBK-NBG-NBH	-3.90	107.96	112.14
2	B	1101	Q4J	NBE-CBD-NBG	3.88	120.63	114.84
2	A	1101	Q4J	CAS-CAR-CAQ	3.77	116.76	111.72
2	B	1101	Q4J	CAB-NAC-NAD	-3.69	106.06	112.47
2	A	1101	Q4J	CAM-C4-N3	3.62	122.03	116.56
2	A	1101	Q4J	CAB-CAA-CAE	-3.47	102.36	104.78
2	B	1101	Q4J	NAF-C6-N1	3.37	121.50	113.33
2	A	1101	Q4J	CBD-NBG-NBH	3.35	126.23	120.71
2	B	1101	Q4J	CAA-CAB-NAC	3.23	110.81	106.38
2	B	1101	Q4J	CBK-CBJ-CBI	3.14	109.11	105.00
2	A	1101	Q4J	CBB-CBC-CBD	3.13	121.99	117.65
2	B	1101	Q4J	CAS-CAN-C2	3.08	118.31	111.85
2	A	1101	Q4J	CAP-CAO-CAN	2.99	114.30	111.46
2	B	1101	Q4J	C5-C6-N1	-2.95	117.23	123.35
2	A	1101	Q4J	CBK-CBJ-CBI	2.86	108.74	105.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	Q4J	CBM-CAB-CAA	-2.84	127.47	131.96
2	A	1101	Q4J	NAF-C6-N1	2.81	120.12	113.33
2	B	1101	Q4J	CBF-NBE-CBD	2.80	122.00	117.41
2	B	1101	Q4J	CAM-C4-N3	2.78	120.76	116.56
2	B	1101	Q4J	CBD-NBG-NBH	2.78	125.29	120.71
2	A	1101	Q4J	CAZ-CBF-NBE	-2.60	121.31	124.47
2	A	1101	Q4J	CBC-CBD-NBE	-2.58	118.37	123.46
2	A	1101	Q4J	C5-C4-N3	-2.55	117.95	121.50
2	A	1101	Q4J	C2-N3-C4	2.54	121.56	116.33
2	B	1101	Q4J	CAO-CAN-C2	2.43	116.95	111.85
2	B	1101	Q4J	CAB-CAA-CAE	2.23	106.34	104.78
2	B	1101	Q4J	CBC-CBD-NBE	-2.20	119.11	123.46
2	A	1101	Q4J	CBF-NBE-CBD	2.11	120.86	117.41
2	B	1101	Q4J	CAP-CAO-CAN	2.05	113.40	111.46
2	A	1101	Q4J	CAS-CAN-C2	2.03	116.10	111.85
2	A	1101	Q4J	CAP-CAQ-CAR	2.03	111.81	109.66
2	A	1101	Q4J	CBB-CAZ-CAY	2.01	124.97	120.75

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1101	Q4J	N3-C2-CAN-CAS
2	B	1101	Q4J	N3-C2-CAN-CAO
2	A	1101	Q4J	CAR-CAQ-CAT-NAW
2	A	1101	Q4J	CBA-CAY-CAZ-CBB
2	B	1101	Q4J	CBA-CAY-CAZ-CBB
2	A	1101	Q4J	CBA-CAY-CAZ-CBF
2	B	1101	Q4J	CBA-CAY-CAZ-CBF
2	A	1101	Q4J	N3-C2-CAN-CAO
2	B	1101	Q4J	N3-C2-CAN-CAS
2	B	1101	Q4J	CAR-CAQ-CAT-NAW
2	B	1101	Q4J	CAP-CAQ-CAT-NAW
2	A	1101	Q4J	N1-C2-CAN-CAO


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1101	Q4J	CAN-CAO-CAP-CAQ-CAR-CAS

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1103	FMT	1	0
3	B	1103	FMT	1	0
2	A	1101	Q4J	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	287/314 (91%)	0.44	17 (5%) 28 30	13, 35, 68, 104	5 (1%)
1	B	290/314 (92%)	0.13	12 (4%) 41 44	10, 28, 59, 103	14 (4%)
All	All	577/628 (91%)	0.28	29 (5%) 34 36	10, 31, 67, 104	19 (3%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	712	LEU	5.2
1	A	712	LEU	4.4
1	A	823	GLY	4.3
1	B	711	ILE	3.7
1	A	844	ARG	3.7
1	A	714	ASP	3.6
1	A	900	TYR	3.6
1	A	924	PHE	3.2
1	A	909	SER	3.0
1	A	899	VAL	2.9
1	A	822	VAL	2.6
1	B	735	PHE	2.5
1	A	821	LYS	2.5
1	A	852	ILE	2.5
1	A	711	ILE	2.4
1	B	770	ARG	2.4
1	B	710	LYS	2.4
1	A	1013	ARG	2.3
1	B	900	TYR	2.3
1	A	851	LEU	2.2
1	B	822	VAL	2.2
1	B	842	ASP	2.2
1	B	748	GLY	2.2
1	B	796	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1010	VAL	2.1
1	B	734	GLU	2.1
1	A	845	ALA	2.1
1	A	1011	LYS	2.0
1	A	885	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FMT	A	1102	3/3	0.82	0.17	54,54,61,63	0
3	FMT	B	1102	3/3	0.88	0.13	37,37,38,43	0
3	FMT	B	1103	3/3	0.88	0.17	38,38,39,41	0
2	Q4J	B	1101	39/39	0.95	0.07	19,27,37,39	0
2	Q4J	A	1101	39/39	0.95	0.07	21,30,36,39	0
3	FMT	A	1103	3/3	0.96	0.14	24,24,25,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Q4J B 1101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Q4J A 1101:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.