



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 11:11 AM UTC

PDB ID : 7JWP / pdb_00007jwp
Title : Fab CJ11 in complex IL-18 peptide liberated by Caspase cleavage
Authors : Payandeh, J.; Ho, H.
Deposited on : 2020-08-26
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	85% 10% 5%
1	C	219	85% 10% 5%
1	E	219	87% 9% 5%
1	G	219	91% 5% 5%
1	I	219	86% 8% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	219	88% 8% 5%
1	M	219	90% 5% 5%
1	O	219	91% 5% 5%
2	B	217	90% 9%
2	D	217	90% 9%
2	F	217	94% 6%
2	H	217	93% 7%
2	J	217	89% 10% .
2	L	217	89% 9% .
2	N	217	90% 8% .
2	P	217	92% 7% .
3	Q	6	83% 17%
3	R	6	83% 17%
3	S	6	100%
3	T	6	100%
3	U	6	100%
3	V	6	100%
3	W	6	67% 17% 17%
3	X	6	83% 17%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 50918 atoms, of which 25165 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab CJ11 Heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	208	3071	985	1528	250	302	6	0	0	0
1	C	208	3071	985	1528	250	302	6	0	0	0
1	E	209	3078	987	1531	251	303	6	0	0	0
1	G	209	3078	987	1531	251	303	6	0	0	0
1	I	208	3071	985	1528	250	302	6	0	0	0
1	K	209	3078	987	1531	251	303	6	0	0	0
1	M	208	3071	985	1528	250	302	6	0	0	0
1	O	209	3078	987	1531	251	303	6	0	0	0

- Molecule 2 is a protein called Fab CJ11 Light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	216	3226	1020	1589	276	336	5	0	0	0
2	D	216	3225	1020	1588	276	336	5	0	0	0
2	F	216	3226	1020	1589	276	336	5	0	0	0
2	H	216	3226	1020	1589	276	336	5	0	0	0
2	J	215	3212	1016	1583	275	333	5	0	0	0
2	L	214	3197	1011	1577	274	330	5	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	N	214	Total	C	H	N	O	S	0	0	0
			3197	1011	1577	274	330	5			
2	P	215	Total	C	H	N	O	S	0	0	0
			3211	1015	1583	275	333	5			


- Molecule 3 is a protein called IL-18 peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Q	6	Total	C	H	N	O	0	0	0
			76	24	32	6	14			
3	R	6	Total	C	H	N	O	0	0	0
			76	24	32	6	14			
3	S	6	Total	C	H	N	O	0	0	0
			76	24	32	6	14			
3	T	6	Total	C	H	N	O	0	0	0
			76	24	32	6	14			
3	U	6	Total	C	H	N	O	0	0	0
			76	24	32	6	14			
3	V	6	Total	C	H	N	O	0	0	0
			76	24	32	6	14			
3	W	5	Total	C	H	N	O	0	0	0
			70	22	30	5	13			
3	X	6	Total	C	H	N	O	0	0	0
			76	24	32	6	14			

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab CJ11 Heavy chain

Chain A: 





- Molecule 1: Fab CJ11 Heavy chain

Chain C: 





- Molecule 1: Fab CJ11 Heavy chain

Chain E: 





- Molecule 1: Fab CJ11 Heavy chain

Chain G: 




- Molecule 1: Fab CJ11 Heavy chain

Chain I: 



- Molecule 1: Fab CJ11 Heavy chain

Chain K: 




- Molecule 1: Fab CJ11 Heavy chain

Chain M: 90% 5% 5%




- Molecule 1: Fab CJ11 Heavy chain

Chain O: 91% 5% 5%



- Molecule 2: Fab CJ11 Light chain

Chain B: 90% 9%



- Molecule 2: Fab CJ11 Light chain

Chain D: 90% 9%



- Molecule 2: Fab CJ11 Light chain

Chain F: 94% 6%




- Molecule 2: Fab CJ11 Light chain

Chain H: 93% 7%



- Molecule 2: Fab CJ11 Light chain

Chain J: 89% 10%



- Molecule 2: Fab CJ11 Light chain

Chain L: 89% 9%




- Molecule 2: Fab CJ11 Light chain

Chain N: 90% 8%



- Molecule 2: Fab CJ11 Light chain

Chain P: 92% 7%



- Molecule 3: IL-18 peptide

Chain Q: 83% 17%



- Molecule 3: IL-18 peptide

Chain R: 83% 17%



- Molecule 3: IL-18 peptide

Chain S: 100%

There are no outlier residues recorded for this chain.

- Molecule 3: IL-18 peptide

Chain T: 100%

There are no outlier residues recorded for this chain.

- Molecule 3: IL-18 peptide

Chain U:  100%

There are no outlier residues recorded for this chain.

- Molecule 3: IL-18 peptide

Chain V:  100%


There are no outlier residues recorded for this chain.

- Molecule 3: IL-18 peptide

Chain W:  67% 17% 17%



- Molecule 3: IL-18 peptide

Chain X:  83% 17%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	81.35Å 83.52Å 145.79Å 107.24° 106.53° 89.98°	Depositor
Resolution (Å)	23.97 – 3.00 23.97 – 3.00	Depositor EDS
% Data completeness (in resolution range)	86.4 (23.97-3.00) 84.9 (23.97-3.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 1.68Å)	Xtrriage
Refinement program	PHENIX dev_2747	Depositor
R, R_{free}	0.235 , 0.260 0.235 , 0.264	Depositor DCC
R_{free} test set	12545 reflections (3.12%)	wwPDB-VP
Wilson B-factor (Å ²)	14.7	Xtrriage
Anisotropy	0.777	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 0.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.000 for k,-h,h+1 0.000 for -k,h,k+1 0.054 for h,-k,-h-l 0.000 for -h,k,-k-l 0.014 for k,h,-h-k-l 0.000 for -k,-h,-l 0.000 for -h,-k,h+k+1	Xtrriage
Reported twinning fraction	0.120 for h,-k,-h-l	Depositor
Outliers	0 of 252856 reflections	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	50918	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 83.02 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4133e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/1584	0.59	0/2168
1	C	0.21	0/1584	0.57	0/2168
1	E	0.21	0/1588	0.60	0/2173
1	G	0.23	0/1588	0.60	0/2173
1	I	0.24	0/1584	0.62	0/2168
1	K	0.24	0/1588	0.63	0/2173
1	M	0.24	0/1584	0.63	0/2168
1	O	0.24	0/1588	0.61	0/2173
2	B	0.21	0/1669	0.58	0/2266
2	D	0.19	0/1669	0.52	0/2266
2	F	0.21	0/1669	0.55	0/2266
2	H	0.19	0/1669	0.53	0/2266
2	J	0.22	0/1661	0.58	0/2255
2	L	0.21	0/1652	0.54	0/2243
2	N	0.22	0/1652	0.56	0/2243
2	P	0.21	0/1660	0.56	0/2254
3	Q	0.15	0/43	0.32	0/55
3	R	0.18	0/43	0.47	0/55
3	S	0.14	0/43	0.25	0/55
3	T	0.12	0/43	0.28	0/55
3	U	0.13	0/43	0.26	0/55
3	V	0.19	0/43	0.47	0/55
3	W	0.12	0/39	0.32	0/50
3	X	0.12	0/43	0.25	0/55
All	All	0.22	0/26329	0.58	0/35858

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1543	1528	1527	14	0
1	C	1543	1528	1527	13	0
1	E	1547	1531	1530	11	0
1	G	1547	1531	1530	5	0
1	I	1543	1528	1527	10	0
1	K	1547	1531	1530	9	0
1	M	1543	1528	1527	6	0
1	O	1547	1531	1530	6	0
2	B	1637	1589	1589	10	0
2	D	1637	1588	1589	10	0
2	F	1637	1589	1589	7	0
2	H	1637	1589	1589	8	0
2	J	1629	1583	1582	12	0
2	L	1620	1577	1576	10	0
2	N	1620	1577	1576	10	0
2	P	1628	1583	1583	8	0
3	Q	44	32	32	2	0
3	R	44	32	32	1	0
3	S	44	32	32	0	0
3	T	44	32	32	0	0
3	U	44	32	32	0	0
3	V	44	32	32	0	0
3	W	40	30	29	1	0
3	X	44	32	32	1	0
All	All	25753	25165	25154	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:J:67:SER:OG	2:J:74:THR:OG1	2.06	0.73
1:C:30:ARG:NH1	1:C:31:TYR:OH	2.25	0.70
2:D:69:SER:O	2:D:71:THR:N	2.25	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:154:ASP:OD1	2:D:192:HIS:ND1	2.24	0.69
1:C:146:ASP:OD2	1:C:173:GLN:NE2	2.27	0.67

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	204/219 (93%)	198 (97%)	6 (3%)	0	100	100
1	C	204/219 (93%)	200 (98%)	4 (2%)	0	100	100
1	E	205/219 (94%)	201 (98%)	4 (2%)	0	100	100
1	G	205/219 (94%)	201 (98%)	4 (2%)	0	100	100
1	I	204/219 (93%)	200 (98%)	3 (2%)	1 (0%)	24	60
1	K	205/219 (94%)	201 (98%)	4 (2%)	0	100	100
1	M	204/219 (93%)	199 (98%)	5 (2%)	0	100	100
1	O	205/219 (94%)	201 (98%)	4 (2%)	0	100	100
2	B	214/217 (99%)	204 (95%)	8 (4%)	2 (1%)	14	48
2	D	214/217 (99%)	204 (95%)	9 (4%)	1 (0%)	24	60
2	F	214/217 (99%)	204 (95%)	9 (4%)	1 (0%)	24	60
2	H	214/217 (99%)	206 (96%)	8 (4%)	0	100	100
2	J	213/217 (98%)	205 (96%)	7 (3%)	1 (0%)	24	60
2	L	212/217 (98%)	203 (96%)	8 (4%)	1 (0%)	24	60
2	N	212/217 (98%)	202 (95%)	9 (4%)	1 (0%)	24	60
2	P	213/217 (98%)	205 (96%)	7 (3%)	1 (0%)	24	60
3	Q	4/6 (67%)	4 (100%)	0	0	100	100
3	R	4/6 (67%)	4 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	S	4/6 (67%)	4 (100%)	0	0	100	100
3	T	4/6 (67%)	4 (100%)	0	0	100	100
3	U	4/6 (67%)	4 (100%)	0	0	100	100
3	V	4/6 (67%)	4 (100%)	0	0	100	100
3	W	3/6 (50%)	3 (100%)	0	0	100	100
3	X	4/6 (67%)	4 (100%)	0	0	100	100
All	All	3373/3536 (95%)	3265 (97%)	99 (3%)	9 (0%)	36	70

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	70	GLY
2	B	70	GLY
2	F	70	GLY
2	J	70	GLY
2	N	70	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/185 (95%)	175 (100%)	0	100	100
1	C	175/185 (95%)	175 (100%)	0	100	100
1	E	175/185 (95%)	175 (100%)	0	100	100
1	G	175/185 (95%)	175 (100%)	0	100	100
1	I	175/185 (95%)	175 (100%)	0	100	100
1	K	175/185 (95%)	175 (100%)	0	100	100
1	M	175/185 (95%)	175 (100%)	0	100	100
1	O	175/185 (95%)	174 (99%)	1 (1%)	78	88
2	B	184/185 (100%)	184 (100%)	0	100	100
2	D	184/185 (100%)	182 (99%)	2 (1%)	65	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	184/185 (100%)	184 (100%)	0	100	100
2	H	184/185 (100%)	184 (100%)	0	100	100
2	J	183/185 (99%)	182 (100%)	1 (0%)	81	89
2	L	182/185 (98%)	181 (100%)	1 (0%)	81	89
2	N	182/185 (98%)	182 (100%)	0	100	100
2	P	183/185 (99%)	183 (100%)	0	100	100
3	Q	5/5 (100%)	5 (100%)	0	100	100
3	R	5/5 (100%)	5 (100%)	0	100	100
3	S	5/5 (100%)	5 (100%)	0	100	100
3	T	5/5 (100%)	5 (100%)	0	100	100
3	U	5/5 (100%)	5 (100%)	0	100	100
3	V	5/5 (100%)	5 (100%)	0	100	100
3	W	5/5 (100%)	5 (100%)	0	100	100
3	X	5/5 (100%)	5 (100%)	0	100	100
All	All	2906/3000 (97%)	2901 (100%)	5 (0%)	87	92

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	183	THR
2	D	186	LYS
2	J	216	GLU
2	L	183	THR
1	O	162	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
2	J	141	ASN
1	O	201	ASN
2	L	127	GLN
2	P	27	GLN
2	F	140	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	208/219 (94%)	-1.46	0 100 100	15, 44, 75, 98	0
1	C	208/219 (94%)	-1.60	0 100 100	11, 29, 63, 81	0
1	E	209/219 (95%)	-1.58	0 100 100	10, 27, 55, 77	0
1	G	209/219 (95%)	-1.61	0 100 100	8, 23, 58, 86	0
1	I	208/219 (94%)	-1.64	0 100 100	3, 11, 26, 39	0
1	K	209/219 (95%)	-1.69	0 100 100	3, 10, 21, 35	0
1	M	208/219 (94%)	-1.66	0 100 100	2, 10, 24, 53	0
1	O	209/219 (95%)	-1.69	0 100 100	2, 9, 21, 47	0
2	B	216/217 (99%)	-1.44	0 100 100	18, 43, 81, 123	0
2	D	216/217 (99%)	-1.54	0 100 100	10, 31, 73, 100	0
2	F	216/217 (99%)	-1.62	0 100 100	9, 24, 59, 88	0
2	H	216/217 (99%)	-1.53	0 100 100	11, 35, 73, 124	0
2	J	215/217 (99%)	-1.64	0 100 100	4, 14, 29, 50	0
2	L	214/217 (98%)	-1.65	0 100 100	4, 14, 29, 56	0
2	N	214/217 (98%)	-1.68	0 100 100	3, 14, 25, 41	0
2	P	215/217 (99%)	-1.66	0 100 100	3, 13, 28, 50	0
3	Q	6/6 (100%)	-1.52	0 100 100	39, 42, 43, 48	0
3	R	6/6 (100%)	-1.57	0 100 100	29, 33, 35, 40	0
3	S	6/6 (100%)	-1.49	0 100 100	35, 39, 39, 42	0
3	T	6/6 (100%)	-1.66	0 100 100	25, 33, 37, 37	0
3	U	6/6 (100%)	-1.52	0 100 100	26, 33, 36, 40	0
3	V	6/6 (100%)	-1.60	0 100 100	26, 31, 37, 44	0
3	W	5/6 (83%)	-1.47	0 100 100	26, 29, 32, 35	0
3	X	6/6 (100%)	-1.60	0 100 100	24, 30, 37, 42	0

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
All	All	3437/3536 (97%)	-1.60	0 100 100	2, 19, 63, 124	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.