



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:27 PM UTC

PDB ID : 9JWX / pdb_00009jwx
Title : NifS soaked with(2R,3R)-3-ethoxycarbonylaziridine-2-carboxylic acid
Authors : Otsuka, H.; Nakamura, R.; Fujishiro, T.
Deposited on : 2024-10-11
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cysteine desulfurase IscS.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	362	2817	1777	493	534	1	12	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	VAL	LEU	engineered mutation	UNP O25008
A	138	ARG	LYS	engineered mutation	UNP O25008
A	388	VAL	-	expression tag	UNP O25008
A	389	ASP	-	expression tag	UNP O25008
A	390	LEU	-	expression tag	UNP O25008
A	391	VAL	-	expression tag	UNP O25008
A	392	PRO	-	expression tag	UNP O25008
A	393	ARG	-	expression tag	UNP O25008
A	394	GLY	-	expression tag	UNP O25008
A	395	SER	-	expression tag	UNP O25008
A	396	HIS	-	expression tag	UNP O25008
A	397	HIS	-	expression tag	UNP O25008
A	398	HIS	-	expression tag	UNP O25008
A	399	HIS	-	expression tag	UNP O25008
A	400	HIS	-	expression tag	UNP O25008
A	401	HIS	-	expression tag	UNP O25008

- Molecule 2 is ISOPROPYL ALCOHOL (CCD ID: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	3	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cysteine desulfurase IscS



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	103.18Å 103.18Å 133.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.11 – 2.80 48.11 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.11-2.80) 99.9 (48.11-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.27 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.203 , 0.243 0.205 , 0.245	Depositor DCC
R_{free} test set	917 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	77.4	Xtriage
Anisotropy	0.173	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 66.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2825	wwPDB-VP
Average B, all atoms (Å ²)	99.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, LLP, IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/2843	1.54	30/3850 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	258	VAL	N-CA-CB	13.26	128.57	110.54
1	A	86	PHE	N-CA-CB	10.02	124.98	110.16
1	A	84	VAL	N-CA-CB	9.73	125.17	110.58
1	A	228	PHE	CA-CB-CG	8.67	122.47	113.80
1	A	378	VAL	N-CA-CB	7.73	121.05	110.54
1	A	180	ASP	CA-CB-CG	7.62	120.22	112.60
1	A	80	VAL	N-CA-CB	7.58	119.41	110.55
1	A	30	HIS	CA-CB-CG	-7.56	106.24	113.80
1	A	14	ARG	N-CA-CB	-6.97	99.30	109.83
1	A	120	GLU	CB-CA-C	-6.51	99.35	110.22
1	A	26	PHE	CA-CB-CG	-6.50	107.30	113.80
1	A	359	ARG	N-CA-CB	-6.47	98.54	110.32
1	A	42	THR	CA-CB-OG1	-6.03	100.55	109.60
1	A	4	ARG	CB-CA-C	6.00	119.70	109.80
1	A	244	VAL	O-C-N	5.99	124.25	120.42
1	A	82	LYS	CB-CA-C	5.97	120.70	110.79

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	56	TYR	N-CA-CB	5.90	118.56	110.01
1	A	372	VAL	N-CA-CB	5.83	118.46	110.54
1	A	368	LYS	CB-CA-C	5.70	119.82	110.88
1	A	18	LYS	N-CA-CB	5.62	118.39	110.12
1	A	40	PHE	CB-CA-C	5.60	119.75	110.90
1	A	24	ASP	CA-CB-CG	5.49	118.09	112.60
1	A	57	LYS	N-CA-CB	5.45	118.13	110.12
1	A	129	HIS	CA-CB-CG	5.33	119.13	113.80
1	A	167	GLY	CA-C-N	5.23	127.55	120.44
1	A	167	GLY	C-N-CA	5.23	127.55	120.44
1	A	50	GLU	N-CA-CB	5.20	117.77	110.12
1	A	171	LYS	N-CA-CB	-5.10	102.61	110.16
1	A	63	ASP	CA-CB-CG	5.03	117.63	112.60
1	A	110	THR	CA-CB-OG1	-5.00	102.10	109.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	14	ARG	Sidechain
1	A	359	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2817	0	2820	77	0
2	A	4	0	8	0	0
3	A	1	0	0	0	0
4	A	3	0	0	0	0
All	All	2825	0	2828	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:ILE:HG22	1:A:220:SER:HB2	1.40	1.00
1:A:204:ALA:HB2	1:A:250:MET:HE3	1.47	0.96
1:A:96:HIS:HD1	1:A:145:THR:HG22	1.38	0.88
1:A:141:ILE:HG21	1:A:173:LYS:HG2	1.62	0.80
1:A:96:HIS:ND1	1:A:145:THR:HG22	1.97	0.79
1:A:64:ILE:HG22	1:A:220:SER:CB	2.13	0.78
1:A:10:ASN:HB2	1:A:354:ARG:HH21	1.49	0.76
1:A:96:HIS:HD1	1:A:145:THR:CG2	1.98	0.76
1:A:214:GLY:HA2	1:A:250:MET:HE1	1.68	0.73
1:A:5:ILE:HD12	1:A:365:GLU:HG2	1.70	0.72
1:A:163:ILE:O	1:A:164:GLU:HB3	1.89	0.72
1:A:10:ASN:HB2	1:A:354:ARG:NH2	2.05	0.70
1:A:133:THR:O	1:A:134:ALA:HB3	1.91	0.70
1:A:98:VAL:CG1	1:A:148:VAL:HG22	2.23	0.69
1:A:244:VAL:HB	1:A:245:PRO:HD3	1.75	0.68
1:A:275:LEU:HD21	1:A:355:LEU:HD13	1.76	0.66
1:A:163:ILE:O	1:A:164:GLU:CB	2.45	0.64
1:A:141:ILE:CG2	1:A:173:LYS:HG2	2.28	0.63
1:A:142:THR:H	1:A:145:THR:HG23	1.65	0.61
1:A:141:ILE:HA	1:A:145:THR:HG21	1.83	0.59
1:A:284:ASP:HB2	1:A:304:ARG:HB2	1.84	0.59
1:A:312:LEU:CD1	1:A:324:THR:HG22	2.32	0.58
1:A:32:GLY:HA3	1:A:40:PHE:HB3	1.87	0.56
1:A:137:VAL:HG23	1:A:138:ARG:H	1.71	0.56
1:A:130:GLY:O	1:A:159:LEU:HD13	2.05	0.56
1:A:133:THR:OG1	1:A:136:GLN:HG3	2.06	0.56
1:A:262:ASP:O	1:A:266:GLU:HG2	2.06	0.55
1:A:312:LEU:HD12	1:A:324:THR:HG22	1.89	0.55
1:A:4:ARG:O	1:A:5:ILE:HD13	2.07	0.55
1:A:180:ASP:HA	1:A:201:SER:HB2	1.89	0.54
1:A:285:VAL:HG12	1:A:303:VAL:HA	1.90	0.54
1:A:311:MET:HE2	1:A:373:PHE:CE2	2.44	0.53
1:A:275:LEU:O	1:A:279:LEU:HG	2.07	0.53
1:A:142:THR:O	1:A:144:LYS:N	2.39	0.53
1:A:98:VAL:HG12	1:A:148:VAL:HG22	1.90	0.53
1:A:95:ASN:H	1:A:144:LYS:HG2	1.74	0.52
1:A:133:THR:O	1:A:134:ALA:CB	2.55	0.52
1:A:304:ARG:HB3	1:A:381:ARG:HH22	1.75	0.52
1:A:19:VAL:O	1:A:22:ILE:HG22	2.09	0.51
1:A:105:PRO:O	1:A:109:SER:HB2	2.09	0.51
1:A:204:ALA:CB	1:A:250:MET:HE3	2.31	0.51
1:A:5:ILE:HD12	1:A:365:GLU:CG	2.39	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:VAL:O	1:A:23:MET:HG3	2.12	0.49
1:A:284:ASP:CB	1:A:304:ARG:HB2	2.42	0.48
1:A:149:SER:HA	1:A:178:HIS:O	2.14	0.48
1:A:367:ASP:HA	1:A:370:ILE:HD12	1.96	0.47
1:A:85:TYR:HA	1:A:89:CYS:HB2	1.97	0.47
1:A:244:VAL:N	1:A:245:PRO:CD	2.76	0.47
1:A:137:VAL:O	1:A:141:ILE:HG13	2.15	0.46
1:A:178:HIS:HB2	1:A:199:PHE:HB2	1.98	0.46
1:A:45:HIS:N	1:A:46:PRO:HD2	2.31	0.46
1:A:243:ASN:O	1:A:247:ILE:HG13	2.16	0.45
1:A:5:ILE:CD1	1:A:365:GLU:HG2	2.42	0.45
1:A:124:LEU:HD22	1:A:136:GLN:HB3	1.98	0.45
1:A:306:ILE:HD13	1:A:380:LEU:HB2	1.98	0.45
1:A:134:ALA:O	1:A:169:ILE:HD12	2.17	0.44
1:A:147:LEU:HD12	1:A:176:LEU:O	2.17	0.44
1:A:115:GLU:HA	1:A:119:VAL:O	2.18	0.44
1:A:38:HIS:CE1	1:A:40:PHE:HB2	2.52	0.44
1:A:274:LYS:NZ	1:A:367:ASP:OD1	2.50	0.44
1:A:185:ILE:HD13	1:A:190:VAL:HG21	2.00	0.44
1:A:124:LEU:HD13	1:A:136:GLN:HB2	1.99	0.43
1:A:133:THR:HG23	1:A:136:GLN:OE1	2.18	0.43
1:A:142:THR:C	1:A:144:LYS:H	2.24	0.43
1:A:16:ASP:HB3	1:A:19:VAL:HG23	2.00	0.43
1:A:41:GLY:O	1:A:44:THR:OG1	2.35	0.43
1:A:5:ILE:HB	1:A:320:ILE:HG12	2.01	0.43
1:A:368:LYS:O	1:A:372:VAL:HG23	2.19	0.43
1:A:114:LEU:HD23	1:A:114:LEU:HA	1.86	0.42
1:A:265:LYS:HD2	1:A:265:LYS:HA	1.76	0.42
1:A:142:THR:C	1:A:144:LYS:N	2.77	0.42
1:A:261:LEU:HD23	1:A:261:LEU:HA	1.89	0.42
1:A:104:HIS:CG	1:A:105:PRO:HD2	2.55	0.42
1:A:96:HIS:NE2	1:A:122:THR:HG23	2.35	0.41
1:A:311:MET:HE2	1:A:373:PHE:CD2	2.56	0.41
1:A:279:LEU:HD22	1:A:373:PHE:CD1	2.56	0.40
1:A:304:ARG:HB3	1:A:381:ARG:NH2	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/401 (89%)	337 (94%)	19 (5%)	1 (0%)	36	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/336 (90%)	290 (95%)	14 (5%)	24	58

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ARG
1	A	53	ASP
1	A	64	ILE
1	A	100	THR
1	A	109	SER
1	A	116	SER
1	A	137	VAL
1	A	187	LYS
1	A	203	SER
1	A	226	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	277	GLU
1	A	324	THR
1	A	368	LYS
1	A	381	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	196	ASN
1	A	382	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	206	1	23,24,25	0.58	0	25,32,34	1.30	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	206	1	-	3/16/17/19	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	206	LLP	C5-C4-C4'	3.03	126.15	121.47
1	A	206	LLP	OP4-C5'-C5	2.31	113.68	109.36
1	A	206	LLP	CD-CG-CB	2.28	122.22	113.62
1	A	206	LLP	C3-C4-C4'	-2.23	116.38	120.40
1	A	206	LLP	O3-C3-C2	2.00	121.73	117.58

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	206	LLP	C-CA-CB-CG
1	A	206	LLP	C4-C4'-NZ-CE
1	A	206	LLP	C3-C4-C4'-NZ

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	IPA	A	501	-	3,3,3	0.55	0	3,3,3	0.51	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	361/401 (90%)	0.06	5 (1%) 73 64	51, 94, 159, 199	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	288	VAL	2.7
1	A	231	GLY	2.1
1	A	286	MET	2.0
1	A	328	CYS	2.0
1	A	86	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	206	24/25	0.98	0.06	55,64,78,91	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IPA	A	501	4/4	0.89	0.24	58,59,75,84	0
3	CL	A	502	1/1	0.94	0.17	83,83,83,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.