



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 05:05 PM UTC

PDB ID : 5JY7 / pdb_00005jy7
Title : Complex of Mycobacterium smegmatis trehalose synthase with maltokinase
Authors : Futterer, K.; Kermani, A.A.; Besra, G.S.
Deposited on : 2016-05-13
Resolution : 3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1747 (3.70-3.50)
Clashscore	190562	1827 (3.70-3.50)
Ramachandran outliers	187476	1773 (3.70-3.50)
Sidechain outliers	187428	1772 (3.70-3.50)
RSRZ outliers	180081	1745 (3.70-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	593	 86% 9% 5%
1	B	593	 85% 10% 5%
1	C	593	 87% 8% 5%
1	D	593	 86% 10% 5%
1	E	593	 83% 10% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	F	593		•
1	G	593		•
1	H	593		
2	I	441		••
2	J	441		••
2	K	441		
2	L	441		••
2	M	441		••
2	N	441		
2	O	441		
2	P	441		

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CA	F	601	-	-	-	X

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 56343 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Trehalose synthase/amylase TreS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	565	Total	C	N	O	S	0	0	0
			4610	2952	788	853	17			
1	B	571	Total	C	N	O	S	0	0	0
			4661	2981	800	863	17			
1	C	569	Total	C	N	O	S	0	0	0
			4630	2964	792	857	17			
1	D	571	Total	C	N	O	S	0	0	0
			4661	2981	800	863	17			
1	E	549	Total	C	N	O	S	0	0	0
			4496	2884	764	831	17			
1	F	571	Total	C	N	O	S	0	0	0
			4661	2981	800	863	17			
1	G	571	Total	C	N	O	S	0	0	0
			4661	2981	800	863	17			
1	H	544	Total	C	N	O	S	0	0	0
			4452	2858	756	821	17			

- Molecule 2 is a protein called Maltokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	435	Total	C	N	O	S	0	0	0
			2909	1811	523	569	6			
2	J	435	Total	C	N	O	S	0	0	0
			2899	1807	523	563	6			
2	K	353	Total	C	N	O	S	0	0	0
			2380	1481	432	462	5			
2	L	436	Total	C	N	O	S	0	0	0
			2913	1816	524	566	7			
2	M	431	Total	C	N	O	S	0	0	0
			2887	1799	519	563	6			
2	N	349	Total	C	N	O	S	0	0	0
			2347	1459	428	455	5			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	O	239	Total	C	N	O	S	0	0	0
			1584	974	291	315	4			
2	P	239	Total	C	N	O	S	0	0	0
			1584	974	291	315	4			


- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	1	Total	Ca	0	0
			1	1		
3	C	1	Total	Ca	0	0
			1	1		
3	D	1	Total	Ca	0	0
			1	1		
3	E	1	Total	Ca	0	0
			1	1		
3	F	1	Total	Ca	0	0
			1	1		
3	G	1	Total	Ca	0	0
			1	1		
3	H	1	Total	Ca	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Trehalose synthase/amylase TreS



- Molecule 1: Trehalose synthase/amylase TreS



- Molecule 1: Trehalose synthase/amylase TreS



GLY
ALA
GLN


• Molecule 1: Trehalose synthase/amilase TreS

Chain D:  86% 10% .




• Molecule 1: Trehalose synthase/amilase TreS

Chain E:  83% 10% 7%




• Molecule 1: Trehalose synthase/amilase TreS

Chain F:  86% 10% .





• Molecule 1: Trehalose synthase/amilase TreS



Chain G:  86% 10% .





• Molecule 1: Trehalose synthase/amylase TreS



• Molecule 2: Maltokinase



• Molecule 2: Maltokinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	315.68Å 315.68Å 124.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.90 – 3.60 49.90 – 3.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.90-3.60) 99.8 (49.90-3.60)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	0.17	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 3.57Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.261 , 0.281 0.254 , 0.271	Depositor DCC
R_{free} test set	7119 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	125.2	Xtriage
Anisotropy	0.002	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 79.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.036 for h,-k,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	56343	wwPDB-VP
Average B, all atoms (Å ²)	111.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section:
CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.50	0/4754	0.70	1/6485 (0.0%)
1	B	0.54	0/4807	0.71	1/6556 (0.0%)
1	C	0.50	0/4774	0.71	1/6513 (0.0%)
1	D	0.45	0/4807	0.68	1/6556 (0.0%)
1	E	0.41	0/4636	0.68	1/6324 (0.0%)
1	F	0.46	2/4807 (0.0%)	0.68	1/6556 (0.0%)
1	G	0.40	0/4807	0.67	1/6556 (0.0%)
1	H	0.38	0/4590	0.67	1/6259 (0.0%)
2	I	0.63	0/2973	0.94	5/4091 (0.1%)
2	J	0.58	0/2963	0.89	6/4078 (0.1%)
2	K	0.49	0/2432	0.86	4/3343 (0.1%)
2	L	0.64	2/2977 (0.1%)	0.88	6/4095 (0.1%)
2	M	0.52	0/2950	0.86	4/4058 (0.1%)
2	N	0.50	0/2398	0.88	4/3296 (0.1%)
2	O	0.47	0/1615	0.91	4/2219 (0.2%)
2	P	0.48	0/1615	0.92	4/2219 (0.2%)
All	All	0.49	4/57905 (0.0%)	0.76	45/79204 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	L	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	352	TYR	C-N	14.87	1.53	1.33
2	L	366	ASP	C-N	13.30	1.51	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	61	ARG	CZ-NH1	10.59	1.47	1.32
1	F	61	ARG	CD-NE	10.57	1.61	1.46

The worst 5 of 45 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	61	ARG	CG-CD-NE	8.27	130.19	112.00
1	A	61	ARG	CG-CD-NE	7.46	128.42	112.00
1	G	61	ARG	CG-CD-NE	7.40	128.28	112.00
1	E	61	ARG	CG-CD-NE	7.38	128.23	112.00
1	H	61	ARG	CG-CD-NE	7.38	128.23	112.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	L	365	ASP	Peptide,Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4610	0	4348	34	0
1	B	4661	0	4407	40	0
1	C	4630	0	4359	33	0
1	D	4661	0	4407	38	0
1	E	4496	0	4260	60	0
1	F	4661	0	4407	49	0
1	G	4661	0	4407	77	0
1	H	4452	0	4222	73	0
2	I	2909	0	2344	54	0
2	J	2899	0	2336	53	0
2	K	2380	0	1950	40	0
2	L	2913	0	2356	40	0
2	M	2887	0	2329	53	0
2	N	2347	0	1921	37	0
2	O	1584	0	1255	43	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	P	1584	0	1255	30	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
All	All	56343	0	50563	646	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 646 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:215:PHE:CB	2:I:361:GLN:HE22	1.62	1.11
2:O:215:PHE:CB	2:O:361:GLN:HE22	1.64	1.11
2:L:215:PHE:CB	2:L:361:GLN:HE22	1.64	1.10
2:N:215:PHE:CB	2:N:361:GLN:HE22	1.64	1.10
2:K:215:PHE:CB	2:K:361:GLN:HE22	1.64	1.10

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	561/593 (95%)	546 (97%)	13 (2%)	2 (0%)	30	61
1	B	569/593 (96%)	551 (97%)	16 (3%)	2 (0%)	30	61

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	565/593 (95%)	549 (97%)	14 (2%)	2 (0%)	30	61
1	D	569/593 (96%)	550 (97%)	16 (3%)	3 (0%)	24	57
1	E	545/593 (92%)	533 (98%)	12 (2%)	0	100	100
1	F	569/593 (96%)	553 (97%)	14 (2%)	2 (0%)	30	61
1	G	569/593 (96%)	548 (96%)	20 (4%)	1 (0%)	43	72
1	H	538/593 (91%)	528 (98%)	10 (2%)	0	100	100
2	I	431/441 (98%)	382 (89%)	43 (10%)	6 (1%)	9	38
2	J	431/441 (98%)	382 (89%)	41 (10%)	8 (2%)	6	33
2	K	349/441 (79%)	318 (91%)	25 (7%)	6 (2%)	7	35
2	L	432/441 (98%)	388 (90%)	35 (8%)	9 (2%)	5	31
2	M	425/441 (96%)	381 (90%)	35 (8%)	9 (2%)	5	31
2	N	345/441 (78%)	318 (92%)	24 (7%)	3 (1%)	14	46
2	O	235/441 (53%)	221 (94%)	13 (6%)	1 (0%)	30	61
2	P	235/441 (53%)	220 (94%)	14 (6%)	1 (0%)	30	61
All	All	7368/8272 (89%)	6968 (95%)	345 (5%)	55 (1%)	18	51

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	14	ILE
1	B	25	HIS
1	C	10	VAL
2	I	275	PRO
2	J	275	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	489/512 (96%)	485 (99%)	4 (1%)	73	77
1	B	495/512 (97%)	487 (98%)	8 (2%)	55	69

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	489/512 (96%)	484 (99%)	5 (1%)	68	75
1	D	495/512 (97%)	491 (99%)	4 (1%)	73	77
1	E	480/512 (94%)	477 (99%)	3 (1%)	78	79
1	F	495/512 (97%)	491 (99%)	4 (1%)	73	77
1	G	495/512 (97%)	491 (99%)	4 (1%)	73	77
1	H	475/512 (93%)	472 (99%)	3 (1%)	78	79
2	I	211/351 (60%)	195 (92%)	16 (8%)	12	39
2	J	208/351 (59%)	195 (94%)	13 (6%)	16	44
2	K	178/351 (51%)	174 (98%)	4 (2%)	45	65
2	L	210/351 (60%)	204 (97%)	6 (3%)	37	60
2	M	210/351 (60%)	205 (98%)	5 (2%)	43	63
2	N	175/351 (50%)	172 (98%)	3 (2%)	53	69
2	O	110/351 (31%)	108 (98%)	2 (2%)	51	68
2	P	110/351 (31%)	108 (98%)	2 (2%)	51	68
All	All	5325/6904 (77%)	5239 (98%)	86 (2%)	55	69

5 of 86 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	161	ASN
2	L	157	ASP
2	J	172	VAL
2	K	194	THR
2	M	55	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 67 such sidechains are listed below:

Mol	Chain	Res	Type
2	J	11	GLN
2	K	371	GLN
2	N	161	ASN
1	D	198	GLN
1	D	197	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	565/593 (95%)	-0.44	1 (0%) 91 80	34, 53, 96, 118	0
1	B	571/593 (96%)	-0.46	2 (0%) 88 70	28, 43, 64, 79	0
1	C	569/593 (95%)	-0.45	1 (0%) 91 80	35, 51, 79, 106	0
1	D	571/593 (96%)	-0.37	1 (0%) 91 80	44, 77, 111, 129	0
1	E	549/593 (92%)	-0.18	1 (0%) 91 80	67, 102, 176, 208	0
1	F	571/593 (96%)	-0.14	5 (0%) 81 56	68, 119, 249, 314	0
1	G	571/593 (96%)	-0.10	5 (0%) 81 56	80, 142, 218, 251	0
1	H	544/593 (91%)	0.17	14 (2%) 57 33	89, 159, 248, 316	0
2	I	435/441 (98%)	-0.30	3 (0%) 84 61	38, 62, 111, 129	0
2	J	435/441 (98%)	-0.04	5 (1%) 78 52	55, 92, 152, 171	0
2	K	353/441 (80%)	0.06	3 (0%) 82 58	106, 160, 228, 246	0
2	L	436/441 (98%)	0.06	7 (1%) 70 43	88, 119, 163, 181	0
2	M	431/441 (97%)	-0.05	3 (0%) 84 61	99, 131, 188, 228	0
2	N	349/441 (79%)	0.04	8 (2%) 61 35	116, 177, 233, 254	0
2	O	239/441 (54%)	0.12	7 (2%) 53 30	168, 210, 253, 266	0
2	P	239/441 (54%)	0.15	7 (2%) 53 30	150, 195, 247, 256	0
All	All	7428/8272 (89%)	-0.16	73 (0%) 79 54	28, 106, 220, 316	0

The worst 5 of 73 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	361	GLN	7.7
2	L	358	LEU	5.6
1	H	323	ALA	5.4
1	H	558	VAL	4.9
2	I	361	GLN	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	F	601	1/1	0.71	0.53	20,20,20,20	0
3	CA	D	601	1/1	0.84	0.06	20,20,20,20	0
3	CA	H	601	1/1	0.96	0.29	20,20,20,20	0
3	CA	E	601	1/1	0.97	0.18	20,20,20,20	0
3	CA	G	601	1/1	0.99	0.20	20,20,20,20	0
3	CA	C	601	1/1	1.00	0.02	20,20,20,20	0
3	CA	A	601	1/1	1.00	0.05	20,20,20,20	0
3	CA	B	601	1/1	1.00	0.01	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.