



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:15 AM UTC

PDB ID : 5JYM / pdb_00005jym
Title : Human P-cadherin EC12 with scFv TSP11 bound
Authors : Caaveiro, J.M.M.; Kudo, S.; Tsumoto, K.
Deposited on : 2016-05-14
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1190 (2.46-2.46)
Clashscore	190562	1229 (2.46-2.46)
Ramachandran outliers	187476	1218 (2.46-2.46)
Sidechain outliers	187428	1218 (2.46-2.46)
RSRZ outliers	180081	1190 (2.46-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	 92% 7%
1	C	213	 2% 94% 6%
2	B	257	 9% 85% 12%
2	D	257	 11% 79% 7% 13%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6874 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cadherin-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	0	0	0
			1638	1028	269	337	4			
1	C	213	Total	C	N	O	S	0	0	0
			1638	1028	269	337	4			

- Molecule 2 is a protein called Antibody scFv TSP11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	225	Total	C	N	O	S	0	0	0
			1742	1110	283	341	8			
2	D	223	Total	C	N	O	S	0	0	0
			1728	1102	281	337	8			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Ca	0	0
			4	4		
3	C	4	Total	Ca	0	0
			4	4		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	44	Total	O	0	0
			44	44		
4	B	22	Total	O	0	0
			22	22		
4	C	39	Total	O	0	1
			40	40		
4	D	14	Total	O	0	0
			14	14		

i


- Molecule 1: Cadherin-3




- Chain C:  2% 94% 6%



- Chain B: 9% 85% 12%



- Chain D:  11% 79% 7% 13%



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.70Å 83.43Å 199.82Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	99.91 – 2.45 99.91 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.6 (99.91-2.45) 99.6 (99.91-2.45)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.223 , 0.270 0.226 , 0.270	Depositor DCC
R_{free} test set	1832 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	43.5	Xtriage
Anisotropy	0.398	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6874	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/1673	0.96	2/2281 (0.1%)
1	C	0.84	1/1673 (0.1%)	0.93	2/2281 (0.1%)
2	B	0.70	0/1791	0.91	1/2439 (0.0%)
2	D	0.69	1/1776 (0.1%)	0.89	0/2417
All	All	0.77	2/6913 (0.0%)	0.92	5/9418 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	64	LYS	N-CA	5.43	1.51	1.45
2	D	101	ILE	N-CA	5.11	1.52	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	ASP	N-CA-C	8.65	123.62	112.89
2	B	203	ARG	NE-CZ-NH2	6.22	124.80	119.20
1	C	9	VAL	CB-CA-C	-5.36	104.57	110.52
1	C	127	VAL	CB-CA-C	-5.20	106.42	111.05
1	A	55	LYS	N-CA-C	5.18	117.60	111.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1638	0	1588	4	0
1	C	1638	0	1588	5	0
2	B	1742	0	1659	3	0
2	D	1728	0	1647	9	0
3	A	4	0	0	0	0
3	C	4	0	0	0	0
4	A	44	0	0	0	0
4	B	22	0	0	1	0
4	C	40	0	0	1	0
4	D	14	0	0	0	0
All	All	6874	0	6482	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:16:ALA:O	2:D:86:LEU:HD22	2.07	0.54
2:D:203:ARG:CB	2:D:203:ARG:HH11	2.23	0.51
1:A:102:ASN:HD22	1:A:196:MET:HE2	1.77	0.50
1:C:46:PRO:HA	1:C:47:PRO:C	2.36	0.50
2:D:81:MET:HE3	2:D:83:LEU:HD21	1.95	0.48
1:C:7:ILE:HG12	1:C:22:ASN:OD1	2.13	0.48
2:B:30:THR:HG23	4:B:319:HOH:O	2.13	0.48
1:C:102:ASN:HD22	1:C:196:MET:HE2	1.78	0.48
2:D:203:ARG:HH11	2:D:203:ARG:CG	2.29	0.46
1:A:46:PRO:HA	1:A:47:PRO:C	2.41	0.46
2:D:98:ARG:NE	2:D:99:GLU:O	2.40	0.44
2:B:62:GLU:HA	2:B:65:LYS:HE2	1.99	0.43
2:D:100:GLY:HA2	2:D:108:LEU:O	2.19	0.42
1:C:49:GLY:HA2	4:C:410:HOH:O	2.18	0.42
2:D:99:GLU:HA	2:D:108:LEU:O	2.20	0.42
1:A:15:GLY:CA	1:A:16:PRO:C	2.93	0.42
2:B:203:ARG:HH21	2:B:203:ARG:CG	2.33	0.41
2:D:11:LEU:HD23	2:D:11:LEU:C	2.46	0.41
1:C:48:GLU:OE2	2:D:32:TYR:OH	2.24	0.40
1:A:191:ILE:HD13	1:A:207:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/213 (99%)	207 (98%)	4 (2%)	0	100	100
1	C	211/213 (99%)	208 (99%)	3 (1%)	0	100	100
2	B	219/257 (85%)	208 (95%)	10 (5%)	1 (0%)	24	32
2	D	215/257 (84%)	204 (95%)	10 (5%)	1 (0%)	24	32
All	All	856/940 (91%)	827 (97%)	27 (3%)	2 (0%)	43	54

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	210	GLY
2	D	210	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/185 (100%)	178 (96%)	7 (4%)	29	43
1	C	185/185 (100%)	183 (99%)	2 (1%)	65	76
2	B	189/205 (92%)	186 (98%)	3 (2%)	55	69
2	D	188/205 (92%)	177 (94%)	11 (6%)	18	26
All	All	747/780 (96%)	724 (97%)	23 (3%)	35	50

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	GLU
1	A	107	LYS
1	A	110	GLN
1	A	176	SER
1	A	183	LYS
1	A	186	GLU
1	A	191	ILE
2	B	108	LEU
2	B	145	LEU
2	B	203	ARG
1	C	110	GLN
1	C	161	LEU
2	D	13	LYS
2	D	65	LYS
2	D	86	LEU
2	D	108	LEU
2	D	121	SER
2	D	145	LEU
2	D	167	SER
2	D	203	ARG
2	D	205	SER
2	D	232	GLN
2	D	247	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	174	ASN
2	B	232	GLN
1	C	84	ASN
2	D	231	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/213 (100%)	-0.02	1 (0%) 87 88	26, 39, 67, 89	0
1	C	213/213 (100%)	0.19	5 (2%) 61 63	26, 43, 78, 97	0
2	B	225/257 (87%)	0.58	23 (10%) 12 10	25, 56, 96, 120	0
2	D	223/257 (86%)	0.74	27 (12%) 8 7	30, 60, 101, 122	0
All	All	874/940 (92%)	0.38	56 (6%) 25 23	25, 48, 92, 122	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	178	TYR	5.2
2	D	11	LEU	4.5
2	B	41	PRO	4.4
2	B	120	VAL	4.2
2	D	148	SER	4.0
2	D	41	PRO	3.9
2	D	145	LEU	3.6
2	D	240	PHE	3.6
2	D	141	ILE	3.5
2	D	15	GLY	3.3
2	B	14	PRO	3.3
2	D	149	PRO	3.3
2	D	13	LYS	3.3
2	B	109	PHE	3.1
2	B	141	ILE	3.0
2	D	121	SER	3.0
2	B	199	GLY	2.8
2	D	161	THR	2.7
1	C	140	TYR	2.7
2	B	118	VAL	2.6
2	B	222	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	231	GLN	2.5
2	D	199	GLY	2.5
2	B	156	PRO	2.5
2	B	85	SER	2.5
2	B	219	SER	2.5
2	D	156	PRO	2.4
2	B	240	PHE	2.4
2	D	12	VAL	2.4
2	B	157	GLY	2.4
2	D	157	GLY	2.4
2	D	14	PRO	2.3
2	B	67	LYS	2.3
2	B	202	ALA	2.3
2	D	151	ILE	2.3
1	C	70	GLU	2.3
2	B	100	GLY	2.3
2	D	205	SER	2.3
2	D	239	THR	2.2
2	D	144	VAL	2.2
2	B	44	GLY	2.2
2	B	45	LEU	2.2
1	C	175	ILE	2.2
2	B	246	LEU	2.1
2	B	16	ALA	2.1
1	C	177	SER	2.1
2	B	88	SER	2.1
2	B	198	SER	2.1
2	D	119	THR	2.1
2	D	84	SER	2.1
1	A	4	VAL	2.0
2	D	154	ALA	2.0
1	C	184	VAL	2.0
2	D	159	LYS	2.0
2	D	182	PRO	2.0
2	D	198	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	C	304	1/1	0.84	0.11	84,84,84,84	0
3	CA	A	304	1/1	0.92	0.09	59,59,59,59	0
3	CA	A	303	1/1	0.99	0.02	37,37,37,37	0
3	CA	A	301	1/1	0.99	0.02	37,37,37,37	0
3	CA	A	302	1/1	0.99	0.03	34,34,34,34	0
3	CA	C	302	1/1	1.00	0.01	35,35,35,35	0
3	CA	C	303	1/1	1.00	0.01	35,35,35,35	0
3	CA	C	301	1/1	1.00	0.01	33,33,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.