



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 01:56 PM EDT

PDB ID : 6JYX / pdb_00006jyx
Title : Structure of CbpJ from Streptococcus Pneumoniae TIGR4
Authors : Xu, Q.; Zhang, J.W.; Li, Q.; Jiang, Y.L.
Deposited on : 2019-04-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	
1	B	303	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CHT	A	401	-	X	-	-
2	CHT	A	403	-	X	X	X
2	CHT	B	401	-	X	-	-
2	CHT	B	403	-	X	-	-
2	CHT	B	404	-	X	-	-
2	CHT	B	405	-	X	-	-
2	CHT	B	407	-	X	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5340 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Choline binding protein J.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	292	Total	C	N	O	S	0	0	0
			2449	1591	397	455	6			
1	B	293	Total	C	N	O	S	0	0	0
			2460	1597	401	456	6			

- Molecule 2 is CHOLINE ION (CCD ID: CHT) (formula: $C_5H_{14}NO$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			7	5	1	1		
2	A	1	Total	C	N	O	0	0
			7	5	1	1		
2	A	1	Total	C	N	O	0	0
			7	5	1	1		
2	A	1	Total	C	N	O	0	0
			7	5	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			7	5	1	1		
2	A	1	Total	C	N	O	0	0
			7	5	1	1		
2	A	1	Total	C	N	O	0	0
			7	5	1	1		
2	B	1	Total	C	N	O	0	0
			7	5	1	1		
2	B	1	Total	C	N	O	0	0
			7	5	1	1		
2	B	1	Total	C	N	O	0	0
			7	5	1	1		
2	B	1	Total	C	N	O	0	0
			7	5	1	1		
2	B	1	Total	C	N	O	0	0
			7	5	1	1		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	125	Total	O	0	0
			125	125		
4	B	194	Total	O	0	0
			194	194		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Choline binding protein J



• Molecule 1: Choline binding protein J



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.52Å 77.38Å 187.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.68 – 2.00 48.68 – 2.00	Depositor EDS
% Data completeness (in resolution range)	93.3 (48.68-2.00) 93.3 (48.68-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.16 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.214 , 0.256 0.216 , 0.256	Depositor DCC
R_{free} test set	2591 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.913	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5340	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, CHT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/2541	0.81	4/3456 (0.1%)
1	B	0.76	0/2552	0.78	0/3470
All	All	0.72	0/5093	0.80	4/6926 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	94	ILE	N-CA-C	9.35	128.79	109.34
1	A	117	VAL	CA-C-N	-6.58	114.32	122.75
1	A	117	VAL	C-N-CA	-6.58	114.32	122.75
1	A	97	GLU	N-CA-C	-6.01	101.57	110.46

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2449	0	2225	37	2
1	B	2460	0	2238	12	0
2	A	49	0	91	12	2
2	B	49	0	91	4	0
3	A	7	0	10	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	7	0	10	0	0
4	A	125	0	0	3	0
4	B	194	0	0	2	0
All	All	5340	0	4665	54	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:ARG:HD2	1:A:95:ARG:C	1.71	1.14
1:A:127:LYS:CE	1:A:183:GLU:HG2	1.85	1.07
1:A:127:LYS:HE3	1:A:183:GLU:HG2	1.38	1.01
1:A:127:LYS:NZ	1:A:183:GLU:HG2	1.80	0.95
1:A:95:ARG:HD2	1:A:95:ARG:O	1.69	0.92
1:A:95:ARG:C	1:A:95:ARG:CD	2.50	0.84
1:B:45:LYS:HD2	1:B:50:LYS:HG2	1.64	0.80
1:A:95:ARG:O	1:A:95:ARG:CD	2.32	0.77
1:A:127:LYS:HE3	1:A:183:GLU:CG	2.17	0.71
1:A:194:TRP:CE2	2:A:403:CHT:H83	2.28	0.69
1:A:127:LYS:HZ1	1:A:183:GLU:HG2	1.60	0.67
1:A:194:TRP:CE3	2:A:403:CHT:H81	2.31	0.66
1:A:95:ARG:O	1:A:95:ARG:NE	2.30	0.65
1:A:101:LYS:HG3	4:A:594:HOH:O	1.99	0.63
1:A:117:VAL:O	1:A:155:PHE:O	2.17	0.63
2:B:407:CHT:O6	2:B:407:CHT:H72	1.98	0.63
1:A:174:TYR:HB3	1:A:204:LEU:HD11	1.81	0.62
1:A:127:LYS:NZ	1:B:256:TRP:O	2.35	0.58
1:B:34:GLN:HG2	1:B:35:PHE:O	2.04	0.58
2:A:404:CHT:O6	2:A:404:CHT:H72	2.04	0.57
1:B:332:ARG:HB2	4:B:531:HOH:O	2.05	0.56
1:B:127:LYS:HG2	4:B:514:HOH:O	2.08	0.54
1:A:192:ALA:HB1	1:A:206:PRO:HA	1.90	0.54
1:A:325:ASN:ND2	1:A:329:GLU:HB2	2.23	0.53
1:A:118:GLY:HA3	4:A:523:HOH:O	2.10	0.52
1:A:194:TRP:CE2	2:A:403:CHT:C8	2.93	0.51
1:B:215:TRP:CE2	2:B:402:CHT:H61	2.45	0.51
1:B:246:ASP:HB2	1:B:253:LYS:HE3	1.93	0.51
1:B:306:TYR:CD2	1:B:319:VAL:HG11	2.46	0.50
1:A:178:THR:HA	1:A:186:LEU:O	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:325:ASN:HD21	1:A:329:GLU:HB2	1.78	0.49
1:A:194:TRP:CD2	2:A:403:CHT:C8	2.96	0.49
1:B:157:GLN:CD	1:B:157:GLN:H	2.21	0.48
2:B:407:CHT:O6	2:B:407:CHT:C7	2.61	0.48
1:A:194:TRP:CD2	2:A:403:CHT:H81	2.48	0.48
1:A:299:GLN:HA	1:A:303:ASN:O	2.14	0.48
1:A:159:HIS:HD2	3:A:408:PEG:H21	1.79	0.47
1:A:118:GLY:CA	4:A:523:HOH:O	2.63	0.47
1:A:322:TYR:CE2	2:A:406:CHT:H83	2.50	0.47
1:A:127:LYS:HZ1	1:A:183:GLU:CG	2.27	0.45
2:B:402:CHT:H62	2:B:402:CHT:HC41	1.60	0.45
1:A:127:LYS:HE3	1:A:183:GLU:CB	2.47	0.44
1:A:159:HIS:CD2	3:A:408:PEG:H31	2.53	0.43
2:A:403:CHT:H72	2:A:403:CHT:HC41	1.71	0.43
1:B:192:ALA:HB1	1:B:206:PRO:HA	1.99	0.43
1:A:41:ARG:HB2	1:A:43:TYR:CE1	2.54	0.43
1:A:194:TRP:CZ2	2:A:403:CHT:C8	3.02	0.43
1:A:231:MET:SD	2:A:403:CHT:H63	2.59	0.42
1:B:252:MET:HE2	1:B:252:MET:HB3	1.97	0.41
1:A:324:VAL:HA	1:A:329:GLU:O	2.20	0.41
1:B:147:LYS:HG3	1:B:147:LYS:O	2.20	0.41
2:A:407:CHT:HC41	2:A:407:CHT:H72	1.82	0.41
1:A:94:ILE:H	1:A:94:ILE:HG13	1.73	0.40
1:A:194:TRP:CZ3	2:A:403:CHT:H81	2.55	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:ARG:CB	2:A:403:CHT:C7[4_555]	1.77	0.43
1:A:95:ARG:CG	2:A:403:CHT:C7[4_555]	1.99	0.21

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/303 (95%)	279 (97%)	8 (3%)	1 (0%)	36	35
1	B	289/303 (95%)	282 (98%)	7 (2%)	0	100	100
All	All	577/606 (95%)	561 (97%)	15 (3%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	94	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/255 (96%)	239 (98%)	5 (2%)	48	54
1	B	245/255 (96%)	240 (98%)	5 (2%)	48	54
All	All	489/510 (96%)	479 (98%)	10 (2%)	48	54

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	95	ARG
1	A	96	LEU
1	A	286	LEU
1	A	287	ASN
1	A	331	VAL
1	B	38	GLU
1	B	180	ILE
1	B	238	GLU
1	B	241	THR
1	B	332	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	212	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CHT	A	407	-	6,6,6	2.28	1 (16%)	8,8,8	3.15	4 (50%)
2	CHT	A	403	-	6,6,6	2.85	4 (66%)	8,8,8	2.95	2 (25%)
2	CHT	B	407	-	6,6,6	2.18	1 (16%)	8,8,8	2.88	4 (50%)
2	CHT	B	403	-	6,6,6	2.23	2 (33%)	8,8,8	2.80	3 (37%)
2	CHT	B	405	-	6,6,6	2.12	1 (16%)	8,8,8	2.74	4 (50%)
2	CHT	A	401	-	6,6,6	2.14	1 (16%)	8,8,8	2.68	4 (50%)
2	CHT	A	402	-	6,6,6	2.21	1 (16%)	8,8,8	2.72	3 (37%)
2	CHT	A	404	-	6,6,6	2.20	1 (16%)	8,8,8	2.58	4 (50%)
2	CHT	A	406	-	6,6,6	2.20	1 (16%)	8,8,8	2.93	3 (37%)
2	CHT	A	405	-	6,6,6	2.20	1 (16%)	8,8,8	3.22	4 (50%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CHT	B	401	-	6,6,6	2.15	1 (16%)	8,8,8	3.16	4 (50%)
2	CHT	B	402	-	6,6,6	2.31	2 (33%)	8,8,8	2.68	2 (25%)
2	CHT	B	404	-	6,6,6	2.14	1 (16%)	8,8,8	2.95	4 (50%)
3	PEG	B	408	-	6,6,6	0.38	0	5,5,5	0.50	0
3	PEG	A	408	-	6,6,6	0.58	0	5,5,5	0.45	0
2	CHT	B	406	-	6,6,6	2.25	1 (16%)	8,8,8	2.63	3 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CHT	A	407	-	-	1/4/4/4	-
2	CHT	A	403	-	-	2/4/4/4	-
2	CHT	B	407	-	-	4/4/4/4	-
2	CHT	B	403	-	-	4/4/4/4	-
2	CHT	B	405	-	-	3/4/4/4	-
2	CHT	A	401	-	-	3/4/4/4	-
2	CHT	A	402	-	-	1/4/4/4	-
2	CHT	A	404	-	-	1/4/4/4	-
2	CHT	A	406	-	-	1/4/4/4	-
2	CHT	A	405	-	-	1/4/4/4	-
2	CHT	B	401	-	-	3/4/4/4	-
2	CHT	B	402	-	-	0/4/4/4	-
2	CHT	B	404	-	-	3/4/4/4	-
3	PEG	B	408	-	-	3/4/4/4	-
3	PEG	A	408	-	-	2/4/4/4	-
2	CHT	B	406	-	-	3/4/4/4	-

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	403	CHT	O6-C4	-4.84	1.17	1.42
2	B	402	CHT	O6-C4	-4.71	1.18	1.42
2	A	407	CHT	O6-C4	-4.55	1.18	1.42
2	A	402	CHT	O6-C4	-4.49	1.19	1.42
2	A	404	CHT	O6-C4	-4.49	1.19	1.42
2	A	401	CHT	O6-C4	-4.48	1.19	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	406	CHT	O6-C4	-4.47	1.19	1.42
2	B	407	CHT	O6-C4	-4.46	1.19	1.42
2	A	405	CHT	O6-C4	-4.45	1.19	1.42
2	B	406	CHT	O6-C4	-4.42	1.19	1.42
2	B	401	CHT	O6-C4	-4.35	1.19	1.42
2	B	405	CHT	O6-C4	-4.33	1.20	1.42
2	B	403	CHT	O6-C4	-4.33	1.20	1.42
2	B	404	CHT	O6-C4	-4.31	1.20	1.42
2	A	403	CHT	C5-N1	-3.17	1.41	1.51
2	A	403	CHT	C8-N1	-2.80	1.41	1.50
2	A	403	CHT	C6-N1	-2.32	1.43	1.50
2	B	402	CHT	C5-N1	-2.09	1.45	1.51
2	B	403	CHT	C6-N1	-2.04	1.44	1.50

All (48) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	405	CHT	C7-N1-C6	7.43	128.49	108.98
2	A	407	CHT	C7-N1-C6	7.29	128.11	108.98
2	B	401	CHT	C7-N1-C6	7.20	127.89	108.98
2	A	403	CHT	C7-N1-C6	7.18	127.83	108.98
2	A	406	CHT	C7-N1-C6	6.82	126.88	108.98
2	B	403	CHT	C7-N1-C6	6.61	126.33	108.98
2	B	404	CHT	C7-N1-C6	6.61	126.33	108.98
2	B	407	CHT	C7-N1-C6	6.60	126.31	108.98
2	B	402	CHT	C7-N1-C6	6.59	126.28	108.98
2	A	402	CHT	C7-N1-C6	6.37	125.70	108.98
2	A	401	CHT	C7-N1-C6	6.22	125.31	108.98
2	B	405	CHT	C7-N1-C6	6.21	125.28	108.98
2	B	406	CHT	C7-N1-C6	6.21	125.28	108.98
2	A	404	CHT	C7-N1-C6	5.93	124.55	108.98
2	A	407	CHT	C8-N1-C7	-3.75	99.12	108.98
2	A	405	CHT	C8-N1-C7	-3.60	99.51	108.98
2	A	403	CHT	C8-N1-C6	-3.38	100.09	108.98
2	B	404	CHT	C8-N1-C6	-3.24	100.47	108.98
2	B	401	CHT	C8-N1-C7	-3.07	100.91	108.98
2	A	406	CHT	C8-N1-C7	-3.01	101.06	108.98
2	B	401	CHT	C8-N1-C6	-3.00	101.09	108.98
2	B	407	CHT	C8-N1-C7	-2.99	101.13	108.98
2	B	405	CHT	C8-N1-C6	-2.79	101.66	108.98
2	A	405	CHT	C8-N1-C6	-2.71	101.85	108.98
2	B	405	CHT	C8-N1-C7	-2.65	102.00	108.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	CHT	C8-N1-C6	-2.61	102.12	108.98
2	A	401	CHT	C8-N1-C7	-2.52	102.35	108.98
2	A	401	CHT	C8-N1-C6	-2.52	102.36	108.98
2	B	404	CHT	C8-N1-C7	-2.47	102.50	108.98
2	A	407	CHT	C8-N1-C6	-2.46	102.52	108.98
2	A	406	CHT	C8-N1-C6	-2.46	102.53	108.98
2	B	405	CHT	O6-C4-C5	2.40	122.64	111.36
2	A	405	CHT	O6-C4-C5	2.40	122.64	111.36
2	B	407	CHT	C8-N1-C6	-2.38	102.71	108.98
2	B	406	CHT	C8-N1-C7	-2.36	102.78	108.98
2	B	404	CHT	O6-C4-C5	2.34	122.32	111.36
2	A	404	CHT	C8-N1-C6	-2.32	102.89	108.98
2	B	407	CHT	O6-C4-C5	2.31	122.17	111.36
2	A	401	CHT	O6-C4-C5	2.24	121.88	111.36
2	A	402	CHT	C8-N1-C7	-2.22	103.15	108.98
2	B	403	CHT	C8-N1-C6	-2.21	103.17	108.98
2	B	403	CHT	C8-N1-C7	-2.19	103.21	108.98
2	B	406	CHT	O6-C4-C5	2.14	121.38	111.36
2	A	407	CHT	O6-C4-C5	2.09	121.17	111.36
2	A	404	CHT	C8-N1-C7	-2.09	103.50	108.98
2	B	401	CHT	O6-C4-C5	2.06	121.04	111.36
2	B	402	CHT	C8-N1-C7	-2.06	103.57	108.98
2	A	404	CHT	O6-C4-C5	2.05	120.96	111.36

There are no chirality outliers.

All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	CHT	C4-C5-N1-C7
2	B	404	CHT	C4-C5-N1-C6
2	B	403	CHT	C4-C5-N1-C8
3	B	408	PEG	O1-C1-C2-O2
2	A	401	CHT	C4-C5-N1-C8
2	B	401	CHT	C4-C5-N1-C7
2	B	401	CHT	C4-C5-N1-C8
2	B	403	CHT	C4-C5-N1-C6
2	B	404	CHT	C4-C5-N1-C8
2	B	405	CHT	C4-C5-N1-C7
2	B	406	CHT	C4-C5-N1-C6
2	B	406	CHT	C4-C5-N1-C8
2	B	407	CHT	C4-C5-N1-C6
2	B	407	CHT	C4-C5-N1-C7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	407	CHT	C4-C5-N1-C8
2	A	401	CHT	C4-C5-N1-C6
2	B	401	CHT	C4-C5-N1-C6
2	B	405	CHT	C4-C5-N1-C6
2	B	405	CHT	C4-C5-N1-C8
3	A	408	PEG	O2-C3-C4-O4
2	B	404	CHT	C4-C5-N1-C7
2	A	404	CHT	O6-C4-C5-N1
2	A	405	CHT	O6-C4-C5-N1
2	B	403	CHT	O6-C4-C5-N1
2	B	406	CHT	C4-C5-N1-C7
3	A	408	PEG	O1-C1-C2-O2
2	A	403	CHT	O6-C4-C5-N1
2	B	407	CHT	O6-C4-C5-N1
3	B	408	PEG	O2-C3-C4-O4
3	B	408	PEG	C4-C3-O2-C2
2	A	406	CHT	C4-C5-N1-C7
2	B	403	CHT	C4-C5-N1-C7
2	A	407	CHT	C4-C5-N1-C6
2	A	403	CHT	C4-C5-N1-C6
2	A	402	CHT	C4-C5-N1-C7

There are no ring outliers.

7 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	407	CHT	1	0
2	A	403	CHT	9	2
2	B	407	CHT	2	0
2	A	404	CHT	1	0
2	A	406	CHT	1	0
2	B	402	CHT	2	0
3	A	408	PEG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	292/303 (96%)	0.76	21 (7%)	21 20	24, 43, 60, 71	8 (2%)
1	B	293/303 (96%)	0.32	16 (5%)	30 29	25, 36, 52, 65	3 (1%)
All	All	585/606 (96%)	0.54	37 (6%)	26 24	24, 40, 57, 71	11 (1%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	95	ARG	8.0
1	A	94	ILE	8.0
1	A	326	TYR	4.8
1	A	96	LEU	4.4
1	A	302	GLY	4.2
1	A	323	SER	4.0
1	A	331	VAL	3.6
1	A	288	ALA	3.3
1	B	149	TYR	3.2
1	B	326	TYR	3.2
1	B	182	GLY	3.0
1	B	199	SER	2.9
1	A	32	GLY	2.8
1	B	183	GLU	2.8
1	B	241	THR	2.8
1	A	137	LYS	2.6
1	A	321	GLY	2.6
1	A	322	TYR	2.5
1	A	324	VAL	2.4
1	B	157	GLN	2.4
1	B	320	ASP	2.4
1	A	93	GLY	2.3
1	B	180	ILE	2.3
1	A	180	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	97	GLU	2.2
1	A	127	LYS	2.2
1	A	330	TRP	2.2
1	B	302	GLY	2.2
1	B	288	ALA	2.1
1	A	263	TRP	2.1
1	A	57	ILE	2.1
1	A	147	LYS	2.1
1	B	287	ASN	2.0
1	B	247	HIS	2.0
1	B	184	ASN	2.0
1	B	261	ASN	2.0
1	A	118	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CHT	A	403	7/7	0.58	0.50	216,216,216,216	0
2	CHT	A	406	7/7	0.81	0.20	81,81,83,84	0
2	CHT	A	407	7/7	0.82	0.19	79,79,81,82	0
2	CHT	B	402	7/7	0.83	0.21	67,68,69,69	0
2	CHT	B	404	7/7	0.85	0.18	56,58,60,61	0
2	CHT	B	401	7/7	0.86	0.23	67,68,71,73	0
2	CHT	B	405	7/7	0.86	0.20	65,66,68,70	0
2	CHT	A	401	7/7	0.87	0.16	69,70,71,72	0
2	CHT	B	403	7/7	0.87	0.14	42,45,52,55	0
2	CHT	B	406	7/7	0.87	0.18	54,55,60,61	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PEG	A	408	7/7	0.87	0.14	56,59,62,63	0
3	PEG	B	408	7/7	0.88	0.15	53,53,55,58	0
2	CHT	B	407	7/7	0.89	0.16	54,56,61,64	0
2	CHT	A	402	7/7	0.89	0.17	62,64,67,69	0
2	CHT	A	404	7/7	0.89	0.19	67,68,69,70	0
2	CHT	A	405	7/7	0.93	0.15	62,63,65,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.