



wwPDB EM Validation Summary Report ⓘ

Mar 7, 2026 – 02:24 AM UTC

PDB ID : 9JYY / pdb_00009jyy
EMDB ID : EMD-61909
Title : core proteins of mature T7
Authors : Liu, H.R.; Chen, W.Y.
Deposited on : 2024-10-13
Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14081 (2.50 - 3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	88	 15% 5% 81%
1	B	88	 13% 7% 81%
1	C	88	 18% 81%
1	D	88	 13% 6% 82%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	88	
1	F	88	
1	a	88	
1	b	88	
2	S	196	
2	T	196	
2	c	196	
2	d	196	
2	e	196	
2	f	196	
2	g	196	
2	h	196	
3	G	747	
3	H	747	
3	M	747	
3	N	747	
3	O	747	
3	P	747	
3	Q	747	
3	R	747	
4	I	1318	
4	J	1318	
4	K	1318	
4	L	1318	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 87740 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein 6.7.

Mol	Chain	Residues	Atoms				AltConf	Trace
1	a	17	Total	C	N	O	0	0
			116	66	17	33		
1	b	16	Total	C	N	O	0	0
			112	64	16	32		
1	A	17	Total	C	N	O	0	0
			116	66	17	33		
1	D	16	Total	C	N	O	0	0
			112	64	16	32		
1	B	17	Total	C	N	O	0	0
			116	66	17	33		
1	E	16	Total	C	N	O	0	0
			112	64	16	32		
1	C	17	Total	C	N	O	0	0
			116	66	17	33		
1	F	16	Total	C	N	O	0	0
			112	64	16	32		

- Molecule 2 is a protein called Internal virion protein gp14.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	S	112	Total	C	N	O	S	0	0
			843	513	160	165	5		
2	T	111	Total	C	N	O	S	0	0
			835	507	159	164	5		
2	c	112	Total	C	N	O	S	0	0
			843	513	160	165	5		
2	d	111	Total	C	N	O	S	0	0
			835	507	159	164	5		
2	e	112	Total	C	N	O	S	0	0
			843	513	160	165	5		
2	f	111	Total	C	N	O	S	0	0
			835	507	159	164	5		
2	g	112	Total	C	N	O	S	0	0
			843	513	160	165	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
2	h	111	Total	C	N	O	S	0	0
			835	507	159	164	5		

- Molecule 3 is a protein called Internal virion protein gp15.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	M	671	Total	C	N	O	S	0	0
			5335	3300	940	1065	30		
3	N	671	Total	C	N	O	S	0	0
			5335	3300	940	1065	30		
3	O	671	Total	C	N	O	S	0	0
			5335	3300	940	1065	30		
3	P	671	Total	C	N	O	S	0	0
			5335	3300	940	1065	30		
3	Q	671	Total	C	N	O	S	0	0
			5335	3300	940	1065	30		
3	R	671	Total	C	N	O	S	0	0
			5335	3300	940	1065	30		
3	G	671	Total	C	N	O	S	0	0
			5335	3300	940	1065	30		
3	H	671	Total	C	N	O	S	0	0
			5335	3300	940	1065	30		


- Molecule 4 is a protein called Peptidoglycan transglycosylase gp16.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	I	1223	Total	C	N	O	S	0	0
			9359	5853	1654	1805	47		
4	J	1223	Total	C	N	O	S	0	0
			9359	5853	1654	1805	47		
4	K	1223	Total	C	N	O	S	0	0
			9359	5853	1654	1805	47		
4	L	1223	Total	C	N	O	S	0	0
			9359	5853	1654	1805	47		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Protein 6.7




• Molecule 1: Protein 6.7



• Molecule 1: Protein 6.7



• Molecule 1: Protein 6.7



LYS
ALA
SER
GLY
ASN
GLY
SER
ALA
MET
LYS
LYS
SER
SER
ILE
ARG
LYS
ALA
PHE
GLY
LYS
LYS

• Molecule 1: Protein 6.7

Chain B:  13% 7% 81%

MET CYS PHE SER GLY ASN PRO LYS ILE SER THR LYS MET LYS MET ASP THR ASN GLN ILE ARG ALA VAL GLU PRO PRO LEU THR GLN GLU VAL SER SER VAL GLU F34 E40 E41 D42 T43 E44 E47 V48 S49 G50 ARG LYS GLY LEU LYS VAL ARG ASP SER VAL LYS SER

LYS
ALA
SER
GLY
ASN
GLY
SER
ALA
MET
LYS
LYS
SER
SER
ILE
ARG
LYS
ALA
PHE
GLY
LYS
LYS

• Molecule 1: Protein 6.7

Chain E:  13% 6% 82%

MET CYS PHE SER GLY ASN PRO LYS ILE SER THR LYS MET LYS MET ASP THR ASN GLN ILE ARG ALA VAL GLU PRO PRO LEU THR GLN GLU VAL SER SER VAL GLU F34 D39 E40 T43 E44 G45 T46 E47 V48 S49 GLY ARG LYS GLY LEU LYS VAL ARG ASP SER VAL LYS SER

SER
LYS
ALA
SER
GLY
ASN
GLY
SER
ALA
MET
LYS
LYS
SER
SER
ILE
ARG
LYS
SER
PHE
GLY
LYS
LYS

• Molecule 1: Protein 6.7

Chain C:  18% 1% 81%

MET CYS PHE SER GLY ASN PRO LYS ILE SER THR LYS MET LYS MET ASP THR ASN GLN ILE ARG ALA VAL GLU PRO PRO LEU THR GLN GLU VAL SER SER VAL GLU F34 E44 E47 G50 ARG LYS GLY LEU LYS VAL ARG ASP SER VAL LYS SER ASN

GLY
SER
ALA
ARG
MET
LYS
LYS
SER
SER
ILE
ARG
LYS
LYS
SER
ALA
PHE
GLY
LYS
LYS


• Molecule 1: Protein 6.7

Chain F:  16% 1% 82%


MET CYS PHE SER GLY ASN PRO LYS ILE SER THR LYS MET LYS MET ASP THR ASN GLN ILE ARG ALA VAL GLU PRO PRO LEU THR GLN GLU VAL SER SER VAL GLU F34 E40 E44 G45 T46 E47 V48 S49 GLY ARG LYS GLY LEU LYS VAL ARG ASP SER VAL LYS SER

ALA
SER
GLY
ASN
GLY
SER
ALA
ARG
MET
LYS
LYS
SER
SER
ILE
ARG
LYS
SER
PHE
GLY
LYS
LYS


• Molecule 2: Internal virion protein gp14

Chain S:  12% 41% 16% 43%



MET CYS TRP A4 I7 P8 I9 A14 Q15 A16 I17 S18 G19 Q20 N21 A22 Q23 A24 K25 M26 I27 A28 R35 R36 Q37 A38 N39 E40 I41 I47 Q48 N49 A50 D51 L52 SS3 L54 Q55 A56 R57 L60 E61 E62 A63 S64 A65 E66 L67 T68 S69 M72 V75 Q76




• Molecule 2: Internal virion protein gp14



• Molecule 2: Internal virion protein gp14




• Molecule 2: Internal virion protein gp14




• Molecule 2: Internal virion protein gp14







• Molecule 3: Internal virion protein gp15




• Molecule 3: Internal virion protein gp15












• Molecule 3: Internal virion protein gp15







• Molecule 4: Peptidoglycan transglycosylase gp16




- Molecule 4: Peptidoglycan transglycosylase gp16





• Molecule 4: Peptidoglycan transglycosylase gp16





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	45731	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	35	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	4000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	19.307	Depositor
Minimum map value	-11.325	Depositor
Average map value	0.004	Depositor
Map value standard deviation	0.908	Depositor
Recommended contour level	2	Depositor
Map size (Å)	339.19998, 339.19998, 339.19998	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/116	0.33	0/155
1	B	0.18	0/116	0.38	0/155
1	C	0.14	0/116	0.34	0/155
1	D	0.21	0/112	0.42	0/150
1	E	0.21	0/112	0.43	0/150
1	F	0.20	0/112	0.46	0/150
1	a	0.18	0/116	0.31	0/155
1	b	0.20	0/112	0.41	0/150
2	S	0.21	0/846	0.42	0/1135
2	T	0.21	0/838	0.41	0/1124
2	c	0.19	0/846	0.42	0/1135
2	d	0.19	0/838	0.37	0/1124
2	e	0.18	0/846	0.39	0/1135
2	f	0.25	0/838	0.47	1/1124 (0.1%)
2	g	0.19	0/846	0.40	0/1135
2	h	0.18	0/838	0.38	0/1124
3	G	0.28	0/5415	0.52	5/7291 (0.1%)
3	H	0.27	0/5415	0.48	2/7291 (0.0%)
3	M	0.29	0/5415	0.53	6/7291 (0.1%)
3	N	0.28	0/5415	0.47	1/7291 (0.0%)
3	O	0.28	0/5415	0.50	5/7291 (0.1%)
3	P	0.27	0/5415	0.45	0/7291
3	Q	0.29	0/5415	0.52	5/7291 (0.1%)
3	R	0.27	0/5415	0.45	0/7291
4	I	0.29	0/9514	0.52	5/12824 (0.0%)
4	J	0.29	0/9514	0.54	11/12824 (0.1%)
4	K	0.29	0/9514	0.53	6/12824 (0.0%)
4	L	0.30	0/9514	0.55	11/12824 (0.1%)
All	All	0.28	0/89024	0.50	58/119880 (0.0%)

There are no bond length outliers.

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	Q	538	THR	N-CA-C	12.31	124.70	111.28
3	M	538	THR	N-CA-C	11.54	123.85	111.28
3	O	540	LYS	N-CA-C	10.10	122.29	111.28
3	G	540	LYS	N-CA-C	9.92	122.10	111.28
3	M	537	LEU	N-CA-C	9.13	121.00	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	116	0	91	4	0
1	B	116	0	91	4	0
1	C	116	0	91	1	0
1	D	112	0	88	5	0
1	E	112	0	88	6	0
1	F	112	0	88	3	0
1	a	116	0	91	1	0
1	b	112	0	88	5	0
2	S	843	0	852	29	0
2	T	835	0	841	22	0
2	c	843	0	852	22	0
2	d	835	0	841	18	0
2	e	843	0	852	21	0
2	f	835	0	841	16	0
2	g	843	0	852	22	0
2	h	835	0	841	23	0
3	G	5335	0	5232	164	0
3	H	5335	0	5232	164	0
3	M	5335	0	5232	168	0
3	N	5335	0	5232	188	0
3	O	5335	0	5232	170	0
3	P	5335	0	5232	186	0
3	Q	5335	0	5232	143	0
3	R	5335	0	5232	183	0
4	I	9359	0	9352	280	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	J	9359	0	9352	275	0
4	K	9359	0	9352	274	0
4	L	9359	0	9352	283	0
All	All	87740	0	86752	2346	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 2346 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:J:64:LEU:HD11	4:J:89:GLN:OE1	1.44	1.17
3:P:79:THR:H	3:P:82:GLN:HE21	1.14	0.95
4:K:669:VAL:HG11	4:K:799:ILE:HD12	1.49	0.95
4:I:669:VAL:HG11	4:I:799:ILE:HD12	1.51	0.91
2:f:42:MET:HG3	3:R:196:MET:HE3	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	15/88 (17%)	15 (100%)	0	0	100	100
1	B	15/88 (17%)	15 (100%)	0	0	100	100
1	C	15/88 (17%)	15 (100%)	0	0	100	100
1	D	14/88 (16%)	14 (100%)	0	0	100	100
1	E	14/88 (16%)	14 (100%)	0	0	100	100
1	F	14/88 (16%)	14 (100%)	0	0	100	100
1	a	15/88 (17%)	15 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	b	14/88 (16%)	14 (100%)	0	0	100	100
2	S	108/196 (55%)	106 (98%)	2 (2%)	0	100	100
2	T	107/196 (55%)	104 (97%)	3 (3%)	0	100	100
2	c	108/196 (55%)	104 (96%)	4 (4%)	0	100	100
2	d	107/196 (55%)	105 (98%)	2 (2%)	0	100	100
2	e	108/196 (55%)	105 (97%)	3 (3%)	0	100	100
2	f	107/196 (55%)	106 (99%)	1 (1%)	0	100	100
2	g	108/196 (55%)	106 (98%)	2 (2%)	0	100	100
2	h	107/196 (55%)	105 (98%)	2 (2%)	0	100	100
3	G	669/747 (90%)	642 (96%)	25 (4%)	2 (0%)	36	70
3	H	669/747 (90%)	636 (95%)	31 (5%)	2 (0%)	36	70
3	M	669/747 (90%)	642 (96%)	27 (4%)	0	100	100
3	N	669/747 (90%)	642 (96%)	24 (4%)	3 (0%)	30	65
3	O	669/747 (90%)	634 (95%)	34 (5%)	1 (0%)	48	80
3	P	669/747 (90%)	641 (96%)	25 (4%)	3 (0%)	30	65
3	Q	669/747 (90%)	640 (96%)	29 (4%)	0	100	100
3	R	669/747 (90%)	634 (95%)	31 (5%)	4 (1%)	21	56
4	I	1217/1318 (92%)	1118 (92%)	86 (7%)	13 (1%)	11	43
4	J	1217/1318 (92%)	1120 (92%)	85 (7%)	12 (1%)	12	45
4	K	1217/1318 (92%)	1112 (91%)	95 (8%)	10 (1%)	16	50
4	L	1217/1318 (92%)	1119 (92%)	84 (7%)	14 (1%)	10	40
All	All	11196/13520 (83%)	10537 (94%)	595 (5%)	64 (1%)	23	56

5 of 64 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	I	61	ALA
4	I	63	ALA
4	I	504	LYS
4	I	1060	LYS
4	J	61	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	13/73 (18%)	13 (100%)	0	100	100
1	B	13/73 (18%)	13 (100%)	0	100	100
1	C	13/73 (18%)	13 (100%)	0	100	100
1	D	13/73 (18%)	13 (100%)	0	100	100
1	E	13/73 (18%)	13 (100%)	0	100	100
1	F	13/73 (18%)	13 (100%)	0	100	100
1	a	13/73 (18%)	13 (100%)	0	100	100
1	b	13/73 (18%)	13 (100%)	0	100	100
2	S	83/149 (56%)	83 (100%)	0	100	100
2	T	82/149 (55%)	82 (100%)	0	100	100
2	c	83/149 (56%)	83 (100%)	0	100	100
2	d	82/149 (55%)	82 (100%)	0	100	100
2	e	83/149 (56%)	83 (100%)	0	100	100
2	f	82/149 (55%)	81 (99%)	1 (1%)	63	82
2	g	83/149 (56%)	83 (100%)	0	100	100
2	h	82/149 (55%)	82 (100%)	0	100	100
3	G	569/624 (91%)	565 (99%)	4 (1%)	76	86
3	H	569/624 (91%)	568 (100%)	1 (0%)	87	92
3	M	569/624 (91%)	567 (100%)	2 (0%)	84	90
3	N	569/624 (91%)	567 (100%)	2 (0%)	84	90
3	O	569/624 (91%)	569 (100%)	0	100	100
3	P	569/624 (91%)	568 (100%)	1 (0%)	87	92
3	Q	569/624 (91%)	567 (100%)	2 (0%)	84	90
3	R	569/624 (91%)	568 (100%)	1 (0%)	87	92
4	I	979/1059 (92%)	975 (100%)	4 (0%)	84	90
4	J	979/1059 (92%)	972 (99%)	7 (1%)	76	86

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	K	979/1059 (92%)	974 (100%)	5 (0%)	81	89
4	L	979/1059 (92%)	975 (100%)	4 (0%)	84	90
All	All	9232/11004 (84%)	9198 (100%)	34 (0%)	81	90

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	K	349	THR
4	K	1191	LEU
4	L	522	LEU
3	G	660	GLN
3	G	539	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 134 such sidechains are listed below:

Mol	Chain	Res	Type
4	J	846	HIS
4	K	567	GLN
4	L	875	ASN
3	P	82	GLN
3	O	695	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-61909. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 181




Y Index: 181




Z Index: 161


6.3.2 Raw map



X Index: 160



Y Index: 160




Z Index: 161


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 576 nm^3 ; this corresponds to an approximate mass of 521 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.00	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.02	3.76	3.09

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-61909 and PDB model 9JYY. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 2.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2).

9.4 Atom inclusion [i](#)



At the recommended contour level, 88% of all backbone atoms, 77% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7720	 0.5140
A	 0.6550	 0.4820
B	 0.7070	 0.5020
C	 0.6470	 0.4990
D	 0.6250	 0.4210
E	 0.6160	 0.4290
F	 0.6610	 0.4250
G	 0.7690	 0.5150
H	 0.7620	 0.5100
I	 0.8140	 0.5310
J	 0.8140	 0.5310
K	 0.8150	 0.5330
L	 0.8160	 0.5330
M	 0.7730	 0.5170
N	 0.7620	 0.5120
O	 0.7740	 0.5190
P	 0.7620	 0.5110
Q	 0.7710	 0.5190
R	 0.7630	 0.5110
S	 0.5990	 0.4300
T	 0.5860	 0.4220
a	 0.6720	 0.4780
b	 0.6340	 0.4370
c	 0.5810	 0.4180
d	 0.5670	 0.4190
e	 0.6010	 0.4290
f	 0.5730	 0.4100
g	 0.5960	 0.4210
h	 0.5900	 0.4160

