



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:31 PM UTC

PDB ID : 6JZ3 / pdb_00006jz3
Title : b-glucuronidase from Ruminococcus gnavus in complex with uronic deoxynojirimycin
Authors : Dashnyam, P.; Lin, H.Y.
Deposited on : 2019-04-30
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	627	
1	B	627	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10847 atoms, of which 20 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucuronidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	581	Total	C	N	O	S	0	1	0
			4770	3075	778	893	24			
1	B	578	Total	C	N	O	S	0	1	0
			4747	3059	776	888	24			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	initiating methionine	UNP Q6W7J7
A	-22	HIS	-	expression tag	UNP Q6W7J7
A	-21	HIS	-	expression tag	UNP Q6W7J7
A	-20	HIS	-	expression tag	UNP Q6W7J7
A	-19	HIS	-	expression tag	UNP Q6W7J7
A	-18	HIS	-	expression tag	UNP Q6W7J7
A	-17	HIS	-	expression tag	UNP Q6W7J7
A	-16	SER	-	expression tag	UNP Q6W7J7
A	-15	SER	-	expression tag	UNP Q6W7J7
A	-14	GLY	-	expression tag	UNP Q6W7J7
A	-13	VAL	-	expression tag	UNP Q6W7J7
A	-12	ASP	-	expression tag	UNP Q6W7J7
A	-11	LEU	-	expression tag	UNP Q6W7J7
A	-10	GLY	-	expression tag	UNP Q6W7J7
A	-9	THR	-	expression tag	UNP Q6W7J7
A	-8	GLU	-	expression tag	UNP Q6W7J7
A	-7	ASN	-	expression tag	UNP Q6W7J7
A	-6	LEU	-	expression tag	UNP Q6W7J7
A	-5	TYR	-	expression tag	UNP Q6W7J7
A	-4	PHE	-	expression tag	UNP Q6W7J7
A	-3	GLN	-	expression tag	UNP Q6W7J7
A	-2	SER	-	expression tag	UNP Q6W7J7
A	-1	ASN	-	expression tag	UNP Q6W7J7
A	0	GLY	-	expression tag	UNP Q6W7J7
B	-23	MET	-	initiating methionine	UNP Q6W7J7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-22	HIS	-	expression tag	UNP Q6W7J7
B	-21	HIS	-	expression tag	UNP Q6W7J7
B	-20	HIS	-	expression tag	UNP Q6W7J7
B	-19	HIS	-	expression tag	UNP Q6W7J7
B	-18	HIS	-	expression tag	UNP Q6W7J7
B	-17	HIS	-	expression tag	UNP Q6W7J7
B	-16	SER	-	expression tag	UNP Q6W7J7
B	-15	SER	-	expression tag	UNP Q6W7J7
B	-14	GLY	-	expression tag	UNP Q6W7J7
B	-13	VAL	-	expression tag	UNP Q6W7J7
B	-12	ASP	-	expression tag	UNP Q6W7J7
B	-11	LEU	-	expression tag	UNP Q6W7J7
B	-10	GLY	-	expression tag	UNP Q6W7J7
B	-9	THR	-	expression tag	UNP Q6W7J7
B	-8	GLU	-	expression tag	UNP Q6W7J7
B	-7	ASN	-	expression tag	UNP Q6W7J7
B	-6	LEU	-	expression tag	UNP Q6W7J7
B	-5	TYR	-	expression tag	UNP Q6W7J7
B	-4	PHE	-	expression tag	UNP Q6W7J7
B	-3	GLN	-	expression tag	UNP Q6W7J7
B	-2	SER	-	expression tag	UNP Q6W7J7
B	-1	ASN	-	expression tag	UNP Q6W7J7
B	0	GLY	-	expression tag	UNP Q6W7J7

- Molecule 2 is (2 {S},3 {R},4 {R},5 {S})-3,4,5-tris(oxidanyl)piperidine-2-carboxylic acid (CCD ID: CN0) (formula: C₆H₁₁NO₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			22	6	10	1	5		
2	B	1	Total	C	H	N	O	0	0
			22	6	10	1	5		

- Molecule 3 is (4R)-2-METHYLPENTANE-2,4-DIOL (CCD ID: MRD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			8	6	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	611	Total	O	0	0
			611	611		
4	B	667	Total	O	0	0
			667	667		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-glucuronidase



• Molecule 1: Beta-glucuronidase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	162.50Å 103.33Å 112.21Å 90.00° 130.92° 90.00°	Depositor
Resolution (Å)	28.38 – 1.50 28.38 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.1 (28.38-1.50) 98.0 (28.38-1.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.45 (at 1.50Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.149 , 0.169 (Not available) , 0.181	Depositor DCC
R_{free} test set	2000 reflections (0.90%)	wwPDB-VP
Wilson B-factor (Å ²)	13.7	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.010 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10847	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CN0, MRD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	17/4905 (0.3%)	0.97	1/6649 (0.0%)
1	B	1.02	18/4881 (0.4%)	0.96	0/6617
All	All	1.03	35/9786 (0.4%)	0.97	1/13266 (0.0%)

All (35) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	299	GLY	C-O	-8.30	1.14	1.23
1	B	297	GLY	C-O	-8.25	1.12	1.23
1	B	565	THR	C-O	-7.41	1.15	1.23
1	A	99	GLY	C-O	-7.28	1.16	1.23
1	B	528	MET	C-O	-6.85	1.15	1.23
1	A	518	MET	C-O	-6.69	1.15	1.24
1	B	285	LYS	C-O	-6.55	1.16	1.23
1	A	161	LEU	C-O	-6.51	1.16	1.23
1	A	364	ARG	C-O	-6.47	1.16	1.23
1	B	582	ASP	C-O	-6.35	1.15	1.24
1	B	580	THR	C-O	-6.27	1.15	1.23
1	A	11	GLN	C-O	-6.19	1.17	1.24
1	A	285	LYS	C-O	-6.11	1.16	1.23
1	B	307	LEU	C-O	-6.00	1.16	1.24
1	B	566	GLU	C-O	-5.90	1.16	1.23
1	A	280	ARG	C-O	-5.82	1.16	1.23
1	B	298	PHE	C-O	-5.63	1.17	1.23
1	B	279	VAL	C-O	-5.61	1.18	1.24
1	A	53	SER	C-O	-5.58	1.17	1.23
1	B	422	ILE	C-O	-5.52	1.16	1.24
1	B	479	ILE	C-O	-5.52	1.18	1.24
1	A	479	ILE	C-O	-5.47	1.18	1.24
1	A	312	HIS	C-O	-5.44	1.17	1.23
1	A	265	GLY	C-O	-5.39	1.18	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	213	GLU	C-O	-5.36	1.16	1.24
1	A	100	THR	C-O	-5.34	1.17	1.24
1	A	262	ASP	CA-C	-5.32	1.48	1.52
1	B	306	ILE	C-O	-5.30	1.17	1.24
1	B	248	VAL	C-O	-5.14	1.18	1.24
1	A	158	GLY	C-O	-5.11	1.16	1.23
1	A	453	LYS	C-O	-5.10	1.17	1.24
1	B	581	ARG	C-O	-5.10	1.17	1.24
1	B	567	GLY	C-O	-5.09	1.18	1.24
1	A	207	GLU	C-O	-5.04	1.17	1.23
1	A	48	SER	C-O	-5.00	1.17	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	157	ASN	N-CA-C	5.66	119.49	112.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4770	0	4556	52	0
1	B	4747	0	4540	47	0
2	A	12	10	0	0	0
2	B	12	10	0	0	0
3	B	8	0	13	3	0
4	A	611	0	0	6	0
4	B	667	0	0	8	0
All	All	10827	20	9109	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:263:GLY:CA	1:A:264:ASN:HB2	1.45	1.43
1:A:263:GLY:HA3	1:A:264:ASN:CB	1.49	1.31
1:B:590:VAL:HG23	4:B:1091:HOH:O	1.35	1.25
1:A:363:MET:CE	1:A:454:ASN:HD22	1.56	1.17
1:A:363:MET:HE1	1:A:454:ASN:ND2	1.60	1.16
1:A:590:VAL:HG23	4:A:896:HOH:O	1.43	1.15
1:B:248:VAL:HG23	1:B:288:LEU:CD2	1.87	1.04
1:B:248:VAL:HG23	1:B:288:LEU:HD23	1.38	1.04
1:B:248:VAL:CG2	1:B:288:LEU:CD2	2.37	1.02
1:A:363:MET:HE1	1:A:454:ASN:HD22	0.74	0.90
1:B:147:LEU:HD21	1:B:362:MET:HE3	1.54	0.90
1:A:82:GLU:OE1	1:B:82:GLU:HG2	1.74	0.86
1:B:363:MET:SD	1:B:452:GLU:OE1	2.37	0.82
1:A:312:HIS:HD2	1:B:312:HIS:HD2	1.29	0.79
1:B:248:VAL:CG2	1:B:288:LEU:HD23	2.06	0.78
1:A:263:GLY:CA	1:A:264:ASN:CB	2.30	0.77
1:B:147:LEU:CD2	1:B:362:MET:HE3	2.16	0.75
1:A:363:MET:CE	1:A:454:ASN:ND2	2.36	0.72
1:A:207:GLU:HG2	1:A:235:GLU:HG2	1.71	0.71
1:A:80:PRO:HB3	1:A:82:GLU:OE2	1.91	0.70
1:A:262:ASP:HB2	1:A:267:LEU:HD13	1.77	0.67
1:A:147:LEU:HD21	1:A:362:MET:HE3	1.76	0.66
1:B:147:LEU:HD21	1:B:362:MET:CE	2.23	0.66
1:A:312:HIS:HD2	1:B:312:HIS:CD2	2.13	0.66
1:A:312:HIS:CD2	1:B:312:HIS:HD2	2.14	0.64
1:A:147:LEU:CD2	1:A:362:MET:HE3	2.29	0.62
1:A:458:GLU:OE2	1:A:498:ARG:HD3	2.00	0.62
1:B:280:ARG:HB2	4:B:1302:HOH:O	1.99	0.61
1:B:118:VAL:HG11	4:B:1345:HOH:O	2.00	0.61
1:B:248:VAL:HG23	1:B:288:LEU:HD21	1.81	0.59
1:A:155:LEU:HD12	1:A:155:LEU:N	2.17	0.59
1:A:263:GLY:HA3	1:A:264:ASN:HB2	0.66	0.59
1:A:157:ASN:C	1:A:157:ASN:HD22	2.10	0.58
1:B:248:VAL:CG2	1:B:288:LEU:HD21	2.33	0.58
1:B:248:VAL:HG22	1:B:288:LEU:CD2	2.32	0.57
1:B:413:TRP:CE2	1:B:445:ARG:HD2	2.40	0.57
1:B:248:VAL:CG2	1:B:288:LEU:HD22	2.34	0.57
1:B:84:ARG:O	1:B:85:ASN:HB2	2.05	0.57
1:A:147:LEU:HD21	1:A:362:MET:CE	2.36	0.56
1:B:363:MET:HG2	1:B:420:GLU:N	2.21	0.56
1:A:420:GLU:OE2	1:A:423:THR:HG23	2.07	0.55
1:B:161:LEU:CD2	4:B:841:HOH:O	2.53	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:98:ARG:HG3	3:B:702:MRD:H4	1.89	0.54
1:A:384:LEU:N	1:A:384:LEU:HD12	2.22	0.54
1:A:565:THR:HG21	4:A:1354:HOH:O	2.07	0.54
1:A:458:GLU:OE2	1:A:498:ARG:CD	2.56	0.53
1:B:161:LEU:HD23	4:B:841:HOH:O	2.08	0.53
1:A:155:LEU:N	1:A:155:LEU:CD1	2.72	0.53
1:A:313:TRP:CD2	1:B:23:MET:HG2	2.46	0.51
1:A:364:ARG:NH2	4:A:805:HOH:O	2.31	0.51
1:A:154:ILE:CD1	1:A:160:LYS:HG2	2.41	0.51
1:B:472:ASN:O	1:B:473:ARG:HD3	2.11	0.51
1:A:137:LYS:HD3	4:A:926:HOH:O	2.12	0.50
1:A:36:LYS:NZ	4:A:821:HOH:O	2.44	0.49
1:A:157:ASN:O	1:A:157:ASN:ND2	2.46	0.49
1:B:99:GLY:HA2	3:B:702:MRD:H5C3	1.94	0.49
1:B:435:ALA:O	1:B:439:THR:HG23	2.13	0.48
1:A:263:GLY:HA2	1:A:264:ASN:HB2	1.75	0.48
1:A:231:THR:HG21	1:A:237:ILE:O	2.13	0.48
1:B:98:ARG:HD3	3:B:702:MRD:H1C1	1.95	0.47
1:A:262:ASP:HB2	1:A:267:LEU:CD1	2.44	0.47
1:A:289:ASN:O	1:A:290:ASP:HB2	2.13	0.47
1:A:388:GLU:HG3	4:A:911:HOH:O	2.15	0.46
1:A:435:ALA:O	1:A:439:THR:HG23	2.15	0.46
1:A:78:TYR:HB3	1:B:11:GLN:HG2	1.98	0.46
1:B:66:ASP:HB3	1:B:161:LEU:HD21	1.97	0.46
1:B:441:ASP:OD2	1:B:445:ARG:HG2	2.16	0.45
1:A:35:LYS:HA	1:A:35:LYS:HD2	1.72	0.45
1:B:461:LYS:HD3	4:B:1173:HOH:O	2.16	0.45
1:A:334:SER:HA	1:A:335:HIS:HA	1.84	0.45
1:B:362:MET:HE2	1:B:389:LEU:HD21	1.98	0.44
1:B:494:ASP:O	1:B:498:ARG:HG3	2.17	0.44
1:A:463:TYR:N	1:A:464:PRO:CD	2.81	0.44
1:B:420:GLU:OE2	1:B:422:ILE:HG12	2.17	0.43
1:B:84:ARG:NH2	4:B:817:HOH:O	2.47	0.43
1:A:23:MET:HG2	1:B:313:TRP:CD2	2.53	0.43
1:B:84:ARG:O	1:B:85:ASN:CB	2.65	0.43
1:B:458:GLU:OE2	1:B:498:ARG:NE	2.51	0.42
1:A:154:ILE:HD13	1:A:160:LYS:HG2	2.01	0.42
1:B:261:THR:HG23	4:B:1324:HOH:O	2.18	0.42
1:A:33:ILE:HD12	1:A:33:ILE:C	2.45	0.42
1:A:384:LEU:HD12	1:A:384:LEU:H	1.83	0.42
1:A:82:GLU:H	1:A:82:GLU:CD	2.26	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:463:TYR:N	1:B:464:PRO:CD	2.82	0.42
1:A:418:GLU:N	1:A:419:PRO:CD	2.82	0.41
1:A:297:GLY:HA3	1:A:330:CYS:O	2.20	0.41
1:A:144:GLU:OE2	1:A:364:ARG:NH2	2.54	0.41
1:B:334:SER:HA	1:B:335:HIS:HA	1.88	0.41
1:B:279:VAL:HG21	1:B:411:ILE:HG22	2.03	0.41
1:B:161:LEU:HD23	1:B:161:LEU:HA	1.79	0.40
1:B:61:ASP:OD1	1:B:64:GLU:OE2	2.39	0.40
1:A:470:CYS:HA	1:A:509:VAL:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	578/627 (92%)	556 (96%)	21 (4%)	1 (0%)	43	22
1	B	575/627 (92%)	554 (96%)	21 (4%)	0	100	100
All	All	1153/1254 (92%)	1110 (96%)	42 (4%)	1 (0%)	48	24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	264	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	507/546 (93%)	497 (98%)	10 (2%)	48	20
1	B	505/546 (92%)	502 (99%)	3 (1%)	78	62
All	All	1012/1092 (93%)	999 (99%)	13 (1%)	61	35

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	82	GLU
1	A	155	LEU
1	A	267	LEU
1	A	334	SER
1	A	364	ARG
1	A	452	GLU
1	A	511	THR
1	A	523	LYS
1	A	569	MET
1	B	363	MET
1	B	384	LEU
1	B	425	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	62	HIS
1	A	157	ASN
1	A	454	ASN
1	A	584	GLN
1	B	11	GLN
1	B	85	ASN
1	B	130	GLN
1	B	584	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MRD	B	702	-	7,7,7	1.61	1 (14%)	9,10,10	0.94	0
2	CN0	A	701	-	11,12,12	1.19	0	12,17,17	1.41	1 (8%)
2	CN0	B	701	-	11,12,12	1.44	2 (18%)	12,17,17	1.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MRD	B	702	-	-	3/5/5/5	-
2	CN0	A	701	-	-	0/3/21/21	0/1/1/1
2	CN0	B	701	-	-	1/3/21/21	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	702	MRD	O2-C2	-3.20	1.36	1.44
2	B	701	CN0	C08-C06	-2.45	1.46	1.52
2	B	701	CN0	C02-C08	-2.44	1.48	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	CN0	C03-C02-C08	3.78	114.86	110.25

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	702	MRD	C1-C2-C3-C4
3	B	702	MRD	O2-C2-C3-C4
2	B	701	CN0	N04-C05-C10-O11
3	B	702	MRD	CM-C2-C3-C4


There are no ring outliers.

1 monomer is involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	702	MRD	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand CN0 A 701



Ligand CN0 B 701



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	581/627 (92%)	-0.18	18 (3%)	51 55	8, 14, 31, 64	1 (0%)
1	B	578/627 (92%)	-0.29	16 (2%)	55 59	8, 13, 29, 53	1 (0%)
All	All	1159/1254 (92%)	-0.24	34 (2%)	53 58	8, 14, 30, 64	2 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	264	ASN	8.0
1	B	364	ARG	7.0
1	B	363	MET	6.8
1	B	383	ALA	6.2
1	B	161	LEU	5.4
1	A	384	LEU	5.3
1	B	425	TYR	5.2
1	A	263	GLY	5.1
1	B	384	LEU	5.0
1	A	364	ARG	4.8
1	A	363	MET	4.8
1	A	365	SER	4.7
1	A	425	TYR	4.7
1	A	381	PHE	4.2
1	A	157	ASN	4.0
1	A	383	ALA	3.8
1	A	82	GLU	3.4
1	A	265	GLY	3.3
1	A	155	LEU	3.2
1	B	84	ARG	3.2
1	A	156	ASN	2.9
1	A	262	ASP	2.8
1	B	156	ASN	2.7
1	B	290	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	85	ASN	2.5
1	A	382	GLU	2.4
1	A	290	ASP	2.4
1	B	498	ARG	2.1
1	B	82	GLU	2.1
1	B	422	ILE	2.1
1	B	224	GLU	2.1
1	B	486	GLU	2.1
1	B	244	ARG	2.1
1	B	362	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MRD	B	702	8/8	0.78	0.14	20,20,20,20	0
2	CN0	B	701	12/12	0.98	0.04	9,12,15,16	0
2	CN0	A	701	12/12	0.99	0.03	10,13,15,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around CN0 B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.