



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:45 PM UTC

PDB ID : 8JZI / pdb_00008jzi
Title : Mutant S-adenosylmethionine synthase from *C. glutamicum*
Authors : Lee, S.; Kim, K.J.
Deposited on : 2023-07-05
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	407	2% 85% 9% 6%
1	B	407	3% 85% 8% 7%
1	C	407	4% 86% 8% 6%
1	D	407	9% 86% 8% 6%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-adenosylmethionine synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	S	0	0	0
			2933	1840	516	570	7			
1	B	380	Total	C	N	O	S	0	0	0
			2913	1829	510	567	7			
1	C	382	Total	C	N	O	S	0	0	0
			2918	1832	513	566	7			
1	D	383	Total	C	N	O	S	0	0	0
			2927	1837	515	568	7			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	68	ALA	GLU	engineered mutation	UNP Q9K5E4
B	68	ALA	GLU	engineered mutation	UNP Q9K5E4
C	68	ALA	GLU	engineered mutation	UNP Q9K5E4
D	68	ALA	GLU	engineered mutation	UNP Q9K5E4

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Na	0	0
			2	2		
4	C	2	Total	Na	0	0
			2	2		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	353	Total	O	0	0
			353	353		
6	B	367	Total	O	0	0
			367	367		
6	C	316	Total	O	0	0
			316	316		
6	D	186	Total	O	0	0
			186	186		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: S-adenosylmethionine synthase




- Molecule 1: S-adenosylmethionine synthase




- Molecule 1: S-adenosylmethionine synthase



- Molecule 1: S-adenosylmethionine synthase





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	150.40Å 125.44Å 116.31Å 90.00° 128.91° 90.00°	Depositor
Resolution (Å)	31.55 – 1.76 31.55 – 1.76	Depositor EDS
% Data completeness (in resolution range)	99.2 (31.55-1.76) 73.5 (31.55-1.76)	Depositor EDS
R_{merge}	0.23	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.36 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.182 , 0.218 (Not available) , 0.221	Depositor DCC
R_{free} test set	8140 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.032	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.024 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12969	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MES, GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.14	6/2985 (0.2%)	1.27	3/4055 (0.1%)
1	B	1.10	0/2965	1.28	3/4028 (0.1%)
1	C	1.15	0/2970	1.29	3/4035 (0.1%)
1	D	1.15	3/2979 (0.1%)	1.36	5/4047 (0.1%)
All	All	1.13	9/11899 (0.1%)	1.30	14/16165 (0.1%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	265	LYS	C-O	6.42	1.30	1.23
1	A	78	ILE	C-O	6.07	1.31	1.24
1	A	277	HIS	CE1-NE2	5.71	1.38	1.32
1	A	264	ARG	C-O	5.44	1.31	1.24
1	A	258	ASP	C-O	5.32	1.29	1.23
1	A	220	GLU	C-O	5.25	1.30	1.24
1	A	44	VAL	C-O	-5.24	1.18	1.24
1	D	304	ASN	C-O	5.19	1.30	1.24
1	D	271	TYR	N-CA	5.18	1.51	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	270	THR	CA-C-O	-7.17	115.68	119.77
1	B	259	ALA	CA-C-N	6.82	126.89	122.18
1	B	259	ALA	C-N-CA	6.82	126.89	122.18
1	D	164	THR	CA-CB-OG1	-5.64	101.13	109.60
1	C	101	GLY	CA-C-O	-5.61	116.24	121.18
1	A	138	PHE	CA-C-N	5.60	125.92	121.61
1	A	138	PHE	C-N-CA	5.60	125.92	121.61
1	D	138	PHE	CA-C-N	5.33	125.72	121.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	138	PHE	C-N-CA	5.33	125.72	121.61
1	D	177	PRO	N-CA-C	5.26	120.84	113.53
1	A	133	ASP	CA-CB-CG	5.18	117.78	112.60
1	B	375	ASP	CA-CB-CG	5.12	117.72	112.60
1	C	138	PHE	CA-C-N	5.01	125.89	121.58
1	C	138	PHE	C-N-CA	5.01	125.89	121.58

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2933	0	2908	24	0
1	B	2913	0	2890	26	0
1	C	2918	0	2895	35	0
1	D	2927	0	2903	19	0
2	A	12	0	16	4	0
2	B	12	0	16	0	0
2	C	6	0	8	0	0
2	D	6	0	8	0	0
3	A	12	0	13	1	0
4	A	2	0	0	0	0
4	C	2	0	0	0	0
5	B	4	0	6	1	0
6	A	353	0	0	3	0
6	B	367	0	0	6	0
6	C	316	0	0	12	0
6	D	186	0	0	0	0
All	All	12969	0	11663	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:503:GOL:H32	6:B:710:HOH:O	1.60	0.99
1:A:49:VAL:HG21	1:B:49:VAL:HG21	1.50	0.92
1:A:104:SER:HB3	6:A:787:HOH:O	1.74	0.87
1:C:49:VAL:HG21	1:D:49:VAL:HG21	1.62	0.81
1:C:266:ILE:HG13	1:C:267:ILE:HD12	1.60	0.81
1:C:54:ILE:CG1	6:C:639:HOH:O	2.30	0.80
1:C:54:ILE:HG13	6:C:639:HOH:O	1.83	0.79
1:B:105:GLN:HG2	1:B:106:GLU:HG3	1.63	0.78
1:C:385:THR:HB	6:C:779:HOH:O	1.82	0.78
1:A:274:MET:SD	2:A:501:GOL:H32	2.26	0.76
1:A:260:GLY:C	1:A:261:LEU:HD12	2.14	0.73
1:C:267:ILE:HD13	1:D:279:GLY:HA2	1.72	0.71
1:A:127:ASP:OD1	1:A:364:ARG:NH1	2.23	0.70
1:A:104:SER:CB	6:A:787:HOH:O	2.35	0.64
1:A:151:LEU:HD21	1:A:235:LEU:HD13	1.80	0.63
1:C:252:LEU:HG	6:C:783:HOH:O	1.98	0.63
1:C:260:GLY:C	1:C:261:LEU:HD12	2.23	0.62
1:C:266:ILE:CG1	1:C:267:ILE:HD12	2.27	0.62
1:A:127:ASP:HA	1:A:360:ALA:HB3	1.84	0.60
1:B:75:ASN:O	1:B:79:GLU:HG3	2.01	0.60
1:C:142:THR:O	1:C:148:TYR:HA	2.02	0.59
1:D:49:VAL:HG13	1:D:261:LEU:CD1	2.34	0.57
1:C:76:LYS:NZ	1:C:79:GLU:OE1	2.30	0.57
1:C:266:ILE:HG13	1:C:267:ILE:CD1	2.33	0.56
1:A:142:THR:O	1:A:148:TYR:HA	2.05	0.56
1:B:78:ILE:HD11	1:B:92:ARG:HG3	1.87	0.55
1:B:142:THR:O	1:B:148:TYR:HA	2.05	0.55
1:C:54:ILE:HD11	6:C:639:HOH:O	2.06	0.55
1:B:204:GLN:HA	1:B:250:PHE:O	2.06	0.55
1:D:49:VAL:HG13	1:D:261:LEU:HD12	1.89	0.55
1:A:252:LEU:HD22	1:A:252:LEU:N	2.21	0.55
1:C:228:LYS:HE2	6:C:874:HOH:O	2.07	0.55
1:D:142:THR:O	1:D:148:TYR:HA	2.07	0.54
5:B:502:EDO:C2	6:B:708:HOH:O	2.55	0.54
1:C:266:ILE:CG1	1:C:267:ILE:CD1	2.86	0.54
1:B:92:ARG:HD2	6:B:905:HOH:O	2.08	0.53
1:C:54:ILE:CD1	6:C:639:HOH:O	2.55	0.53
1:C:228:LYS:HD2	6:C:733:HOH:O	2.08	0.53
1:D:201:ILE:HB	1:D:244:ILE:HD13	1.90	0.53
1:C:267:ILE:HD12	1:C:267:ILE:N	2.22	0.53
1:D:308:ALA:HA	1:D:397:VAL:HG13	1.90	0.53
1:A:285:LYS:NZ	2:A:503:GOL:O2	2.37	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:245:ASN:N	1:C:246:PRO:CD	2.73	0.51
1:C:127:ASP:OD2	1:C:364:ARG:NH1	2.44	0.51
2:A:503:GOL:H31	1:B:265:LYS:NZ	2.26	0.50
1:A:3:GLN:N	1:A:4:PRO:CD	2.75	0.49
1:B:155:LEU:HD23	1:B:183:VAL:HG11	1.94	0.49
1:B:259:ALA:N	6:B:611:HOH:O	2.45	0.49
1:B:49:VAL:HG13	1:B:261:LEU:CD2	2.43	0.49
1:D:127:ASP:HA	1:D:360:ALA:HB3	1.94	0.49
1:C:211:ARG:HG2	6:C:616:HOH:O	2.13	0.48
1:B:49:VAL:HG13	1:B:261:LEU:HD23	1.95	0.47
1:A:261:LEU:HD12	1:A:261:LEU:N	2.29	0.47
1:C:397:VAL:O	1:C:401:ARG:HG2	2.14	0.47
3:A:502:MES:H22	6:A:676:HOH:O	2.15	0.47
1:A:26:ALA:HB1	1:A:76:LYS:HE2	1.96	0.47
1:D:215:GLU:O	1:D:219:ARG:HG3	2.15	0.47
1:B:188:ASP:OD2	1:B:192:ARG:NH1	2.48	0.46
1:D:20:PRO:HA	1:D:23:ILE:HD12	1.98	0.46
1:C:267:ILE:HD13	1:D:279:GLY:CA	2.43	0.45
1:D:127:ASP:OD2	1:D:364:ARG:NH1	2.50	0.45
1:C:127:ASP:CG	1:C:364:ARG:NH1	2.75	0.45
1:D:204:GLN:HA	1:D:250:PHE:O	2.17	0.44
1:C:267:ILE:CD1	1:C:267:ILE:N	2.80	0.44
1:C:396:ARG:HD3	1:C:399:GLU:OE1	2.18	0.44
1:A:266:ILE:HD11	1:B:266:ILE:HD11	1.99	0.44
1:D:183:VAL:HG13	1:D:196:LEU:HD11	2.00	0.44
1:A:71:GLN:HG2	1:D:103:GLN:OE1	2.18	0.43
1:A:246:PRO:HG2	1:B:322:ILE:HB	2.01	0.43
1:C:76:LYS:HA	1:C:76:LYS:HD3	1.85	0.43
1:C:208:GLU:HG3	6:C:626:HOH:O	2.17	0.43
1:A:49:VAL:HG21	1:B:49:VAL:CG2	2.33	0.43
1:C:228:LYS:CE	6:C:874:HOH:O	2.66	0.43
1:C:247:SER:HB3	1:D:322:ILE:HG12	2.00	0.43
1:B:389:LEU:HD12	6:B:849:HOH:O	2.19	0.43
1:C:228:LYS:HE2	6:C:762:HOH:O	2.19	0.43
1:A:245:ASN:N	1:A:246:PRO:CD	2.82	0.43
1:B:102:GLU:C	1:B:103:GLN:HG3	2.44	0.43
1:A:243:LEU:HD21	6:B:902:HOH:O	2.19	0.43
1:B:236:ALA:O	1:B:240:ILE:HD11	2.19	0.43
1:B:105:GLN:HE22	1:C:74:ARG:HD2	1.84	0.42
1:B:102:GLU:O	1:B:103:GLN:HG3	2.19	0.42
1:A:261:LEU:HD11	1:B:47:GLU:HG3	2.01	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:276:ARG:HD2	1:B:273:GLY:O	2.19	0.42
1:B:26:ALA:HB1	1:B:76:LYS:HE3	2.00	0.42
1:B:192:ARG:NH1	1:B:192:ARG:HB2	2.34	0.42
1:D:51:THR:O	1:D:52:THR:C	2.63	0.42
1:B:51:THR:O	1:B:52:THR:C	2.62	0.41
1:C:215:GLU:O	1:C:219:ARG:HG3	2.21	0.41
1:D:183:VAL:HG13	1:D:196:LEU:CD1	2.51	0.41
1:C:127:ASP:CG	1:C:364:ARG:HH12	2.28	0.41
1:A:251:ILE:HG22	1:A:252:LEU:HD22	2.02	0.41
1:C:266:ILE:HG12	1:C:267:ILE:CD1	2.51	0.41
1:A:51:THR:O	1:A:52:THR:C	2.64	0.40
1:D:127:ASP:CG	1:D:364:ARG:NH1	2.79	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/407 (93%)	367 (97%)	13 (3%)	0	100	100
1	B	376/407 (92%)	361 (96%)	15 (4%)	0	100	100
1	C	378/407 (93%)	368 (97%)	10 (3%)	0	100	100
1	D	379/407 (93%)	366 (97%)	13 (3%)	0	100	100
All	All	1513/1628 (93%)	1462 (97%)	51 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	308/327 (94%)	302 (98%)	6 (2%)	50 31
1	B	308/327 (94%)	302 (98%)	6 (2%)	50 31
1	C	306/327 (94%)	302 (99%)	4 (1%)	61 46
1	D	307/327 (94%)	302 (98%)	5 (2%)	55 38
All	All	1229/1308 (94%)	1208 (98%)	21 (2%)	53 36

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	22	LYS
1	A	104	SER
1	A	136	LEU
1	A	261	LEU
1	A	326	LYS
1	B	3	GLN
1	B	22	LYS
1	B	105	GLN
1	B	106	GLU
1	B	136	LEU
1	B	256	MET
1	C	22	LYS
1	C	136	LEU
1	C	256	MET
1	C	267	ILE
1	D	22	LYS
1	D	136	LEU
1	D	249	SER
1	D	261	LEU
1	D	338	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	B	71	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	71	GLN
1	D	190	GLN
1	D	217	GLN
1	D	221	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	501	-	5,5,5	0.12	0	5,5,5	0.54	0
5	EDO	B	502	-	3,3,3	0.24	0	2,2,2	0.88	0
2	GOL	B	503	-	5,5,5	0.32	0	5,5,5	0.35	0
3	MES	A	502	-	12,12,12	0.77	0	15,16,16	0.57	0
2	GOL	D	501	-	5,5,5	0.10	0	5,5,5	0.25	0
2	GOL	A	503	-	5,5,5	0.19	0	5,5,5	0.67	0
2	GOL	A	501	-	5,5,5	0.18	0	5,5,5	1.32	1 (20%)
2	GOL	C	501	-	5,5,5	0.22	0	5,5,5	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	501	-	-	2/4/4/4	-
5	EDO	B	502	-	-	1/1/1/1	-
2	GOL	B	503	-	-	0/4/4/4	-
3	MES	A	502	-	-	3/6/14/14	0/1/1/1
2	GOL	D	501	-	-	2/4/4/4	-
2	GOL	A	503	-	-	4/4/4/4	-
2	GOL	A	501	-	-	3/4/4/4	-
2	GOL	C	501	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	GOL	O2-C2-C3	2.51	119.59	109.18

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	503	GOL	O1-C1-C2-O2
2	A	503	GOL	O1-C1-C2-C3
2	A	503	GOL	C1-C2-C3-O3
2	B	501	GOL	O1-C1-C2-C3
3	A	502	MES	C7-C8-S-O2S
3	A	502	MES	C7-C8-S-O3S
2	A	501	GOL	O1-C1-C2-O2
2	B	501	GOL	O1-C1-C2-O2
2	A	501	GOL	O1-C1-C2-C3
2	D	501	GOL	O1-C1-C2-C3
2	A	503	GOL	O2-C2-C3-O3
2	A	501	GOL	O2-C2-C3-O3
2	D	501	GOL	O1-C1-C2-O2
5	B	502	EDO	O1-C1-C2-O2
3	A	502	MES	C7-C8-S-O1S

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	502	EDO	1	0
3	A	502	MES	1	0
2	A	503	GOL	3	0
2	A	501	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	384/407 (94%)	0.06	10 (2%) 57 63	9, 15, 32, 54	0
1	B	380/407 (93%)	0.18	12 (3%) 50 57	10, 16, 32, 81	0
1	C	382/407 (93%)	0.34	18 (4%) 36 42	10, 18, 37, 73	0
1	D	383/407 (94%)	0.74	38 (9%) 13 15	12, 22, 43, 61	0
All	All	1529/1628 (93%)	0.33	78 (5%) 33 39	9, 18, 37, 81	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	257	GLY	6.0
1	C	257	GLY	4.9
1	B	105	GLN	4.7
1	D	126	ASP	4.5
1	B	66	TYR	4.3
1	C	126	ASP	4.1
1	D	213	TRP	4.1
1	D	257	GLY	4.0
1	B	258	ASP	4.0
1	A	66	TYR	3.9
1	D	212	ALA	3.8
1	D	252	LEU	3.7
1	D	195	HIS	3.7
1	D	4	PRO	3.5
1	D	103	GLN	3.4
1	D	209	VAL	3.4
1	D	66	TYR	3.3
1	B	132	GLY	3.3
1	A	103	GLN	3.3
1	C	323	GLY	3.2
1	D	235	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	211	ARG	3.2
1	A	129	ALA	3.2
1	D	250	PHE	3.2
1	C	252	LEU	3.1
1	A	407	ALA	3.1
1	D	407	ALA	3.1
1	B	3	GLN	3.1
1	C	4	PRO	3.0
1	D	214	LEU	3.0
1	C	128	ARG	3.0
1	D	173	PRO	3.0
1	C	385	THR	2.9
1	C	129	ALA	2.9
1	D	232	ILE	2.9
1	B	104	SER	2.9
1	D	253	GLY	2.9
1	D	3	GLN	2.9
1	D	234	ASP	2.8
1	C	66	TYR	2.8
1	D	171	ILE	2.7
1	C	130	GLY	2.7
1	B	252	LEU	2.6
1	B	106	GLU	2.6
1	C	258	ASP	2.6
1	D	172	VAL	2.6
1	A	3	GLN	2.6
1	A	4	PRO	2.6
1	D	239	GLU	2.6
1	D	385	THR	2.6
1	D	183	VAL	2.6
1	B	195	HIS	2.5
1	D	223	ILE	2.5
1	C	133	ASP	2.5
1	B	211	ARG	2.5
1	D	216	THR	2.5
1	D	127	ASP	2.4
1	A	104	SER	2.4
1	D	166	VAL	2.4
1	A	126	ASP	2.4
1	D	258	ASP	2.4
1	D	242	VAL	2.4
1	D	128	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	406	LEU	2.3
1	C	127	ASP	2.3
1	D	61	VAL	2.3
1	D	240	ILE	2.2
1	C	407	ALA	2.2
1	D	341	GLY	2.2
1	D	222	VAL	2.2
1	D	230	ALA	2.1
1	C	352	LEU	2.1
1	A	194	SER	2.1
1	C	35	LEU	2.1
1	D	227	ILE	2.1
1	B	236	ALA	2.0
1	C	405	LYS	2.0
1	A	234	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	D	501	6/6	0.82	0.17	49,51,54,54	0
3	MES	A	502	12/12	0.83	0.15	28,48,51,52	0
5	EDO	B	502	4/4	0.87	0.12	25,28,30,33	0
2	GOL	A	503	6/6	0.88	0.13	25,27,33,35	0
2	GOL	A	501	6/6	0.90	0.10	21,24,25,25	0
2	GOL	B	503	6/6	0.94	0.10	16,20,21,25	0
4	NA	A	504	1/1	0.94	0.09	8,8,8,8	0
2	GOL	B	501	6/6	0.94	0.09	22,26,30,31	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	C	501	6/6	0.95	0.06	19,25,27,28	0
4	NA	C	503	1/1	0.96	0.08	16,16,16,16	0
4	NA	C	502	1/1	0.98	0.14	10,10,10,10	0
4	NA	A	505	1/1	1.00	0.06	6,6,6,6	0

6.5 Other polymers [i](#)

There are no such residues in this entry.