



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:06 PM UTC

PDB ID : 6K00 / pdb_00006k00
Title : Crystal structure A of ceNAP1-H2A-H2B complex
Authors : Liu, Y.R.
Deposited on : 2019-05-05
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	308	11% 80% 5% 15%
1	B	308	8% 83% 5% 13%
2	D	207	11% 80% 7% 13%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleosome Assembly Protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	0	0
			2083	1341	336	402	4			
1	B	269	Total	C	N	O	S	0	0	0
			2136	1370	346	415	5			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	initiating methionine	UNP Q19007
A	-10	GLY	-	expression tag	UNP Q19007
A	-9	SER	-	expression tag	UNP Q19007
A	-8	SER	-	expression tag	UNP Q19007
A	-7	HIS	-	expression tag	UNP Q19007
A	-6	HIS	-	expression tag	UNP Q19007
A	-5	HIS	-	expression tag	UNP Q19007
A	-4	HIS	-	expression tag	UNP Q19007
A	-3	HIS	-	expression tag	UNP Q19007
A	-2	HIS	-	expression tag	UNP Q19007
A	-1	SER	-	expression tag	UNP Q19007
A	0	SER	-	expression tag	UNP Q19007
A	1	GLY	-	expression tag	UNP Q19007
A	2	GLU	-	expression tag	UNP Q19007
A	3	ASN	-	expression tag	UNP Q19007
A	4	LEU	-	expression tag	UNP Q19007
A	5	TYR	-	expression tag	UNP Q19007
A	6	PHE	-	expression tag	UNP Q19007
A	7	GLN	-	expression tag	UNP Q19007
A	8	HIS	-	expression tag	UNP Q19007
A	9	MET	-	expression tag	UNP Q19007
B	-11	MET	-	initiating methionine	UNP Q19007
B	-10	GLY	-	expression tag	UNP Q19007
B	-9	SER	-	expression tag	UNP Q19007
B	-8	SER	-	expression tag	UNP Q19007

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-7	HIS	-	expression tag	UNP Q19007
B	-6	HIS	-	expression tag	UNP Q19007
B	-5	HIS	-	expression tag	UNP Q19007
B	-4	HIS	-	expression tag	UNP Q19007
B	-3	HIS	-	expression tag	UNP Q19007
B	-2	HIS	-	expression tag	UNP Q19007
B	-1	SER	-	expression tag	UNP Q19007
B	0	SER	-	expression tag	UNP Q19007
B	1	GLY	-	expression tag	UNP Q19007
B	2	GLU	-	expression tag	UNP Q19007
B	3	ASN	-	expression tag	UNP Q19007
B	4	LEU	-	expression tag	UNP Q19007
B	5	TYR	-	expression tag	UNP Q19007
B	6	PHE	-	expression tag	UNP Q19007
B	7	GLN	-	expression tag	UNP Q19007
B	8	HIS	-	expression tag	UNP Q19007
B	9	MET	-	expression tag	UNP Q19007

- Molecule 2 is a protein called Histone H2B 1,Histone H2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	180	Total	C	N	O	S	0	0	0
			1369	858	255	254	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	28	HIS	-	expression tag	UNP P04255
D	29	MET	-	expression tag	UNP P04255


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	25	Total	O	0	0
			25	25		
3	B	19	Total	O	0	0
			19	19		
3	D	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Nucleosome Assembly Protein



• Molecule 1: Nucleosome Assembly Protein



• Molecule 2: Histone H2B 1,Histone H2A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.30Å 78.72Å 169.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.35 – 2.20 49.35 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.7 (49.35-2.20) 98.7 (49.35-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 2.20Å)	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, R_{free}	0.211 , 0.243 0.213 , 0.218	Depositor DCC
R_{free} test set	2365 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.8	Xtriage
Anisotropy	0.421	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5636	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.24	0/2127	0.43	1/2881 (0.0%)
1	B	0.16	0/2182	0.39	0/2954
2	D	0.14	0/1388	0.38	0/1875
All	All	0.19	0/5697	0.41	1/7710 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	GLN	CB-CA-C	-5.86	99.80	110.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2083	0	2044	12	1
1	B	2136	0	2066	10	1
2	D	1369	0	1388	7	0
3	A	25	0	0	0	0
3	B	19	0	0	0	0
3	D	4	0	0	0	0
All	All	5636	0	5498	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:187:ALA:O	1:A:190:GLN:HG3	1.76	0.86
1:B:253:ARG:HH12	1:B:263:GLU:HG2	1.47	0.79
1:B:250:LYS:HE2	1:B:253:ARG:HH21	1.52	0.74
1:A:264:GLU:O	1:A:268:LEU:HD12	1.90	0.70
1:A:247:PRO:HG2	1:A:265:PHE:HB2	1.79	0.64
1:A:261:GLN:H	1:A:261:GLN:CD	2.09	0.61
1:B:248:LYS:HD2	1:B:248:LYS:H	1.68	0.56
1:A:63:LYS:HE3	1:A:67:ASP:OD2	2.06	0.55
1:B:215:LYS:HG3	1:B:235:VAL:HG23	1.89	0.55
2:D:86:ILE:HG13	2:D:155:ALA:HB3	1.90	0.54
1:A:262:ALA:HA	1:A:265:PHE:CD1	2.44	0.52
1:A:196:VAL:HG21	1:A:270:TYR:CG	2.49	0.48
1:B:145:VAL:HG12	1:B:281:ILE:HD11	1.97	0.47
2:D:75:SER:HB2	2:D:154:TYR:HB3	1.98	0.46
1:B:250:LYS:O	1:B:250:LYS:HD3	2.16	0.45
2:D:87:SER:OG	2:D:88:SER:N	2.49	0.45
2:D:67:PHE:CD2	2:D:166:LEU:HD21	2.53	0.44
2:D:189:LYS:HB2	2:D:189:LYS:HE2	1.69	0.44
1:A:283:ARG:NH2	1:B:191:PHE:O	2.51	0.43
1:A:95:ASP:O	1:A:99:GLU:HG2	2.18	0.43
2:D:109:SER:O	2:D:113:LYS:HG3	2.18	0.43
1:B:292:LEU:HD23	1:B:292:LEU:HA	1.87	0.43
1:B:175:LYS:HE3	1:B:246:PRO:HB3	2.02	0.42
1:A:190:GLN:O	1:A:191:PHE:C	2.63	0.42
1:A:96:GLN:NE2	1:B:11:LEU:HD11	2.34	0.41
2:D:195:PRO:O	2:D:199:GLN:HG2	2.21	0.41
1:A:264:GLU:O	1:A:268:LEU:CD1	2.63	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:63:LYS:NZ	1:B:67:ASP:OD2[3_544]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/308 (83%)	254 (99%)	3 (1%)	0	100	100
1	B	263/308 (85%)	260 (99%)	3 (1%)	0	100	100
2	D	176/207 (85%)	169 (96%)	4 (2%)	3 (2%)	7	5
All	All	696/823 (85%)	683 (98%)	10 (1%)	3 (0%)	30	34

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	101	GLY
2	D	157	ARG
2	D	154	TYR

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	222/265 (84%)	222 (100%)	0	100	100
1	B	226/265 (85%)	226 (100%)	0	100	100
2	D	140/166 (84%)	140 (100%)	0	100	100
All	All	588/696 (84%)	588 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	134	HIS
1	A	161	HIS
1	A	274	GLN
1	B	161	HIS
2	D	156	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/308 (85%)	0.40	34 (12%) 7 5	23, 36, 70, 132	0
1	B	269/308 (87%)	0.39	25 (9%) 14 12	26, 37, 78, 111	0
2	D	180/207 (86%)	0.66	23 (12%) 7 5	29, 44, 96, 131	0
All	All	712/823 (86%)	0.46	82 (11%) 9 7	23, 39, 82, 132	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	262	ALA	8.4
1	B	217	VAL	7.9
1	A	296	ASP	7.2
2	D	155	ALA	6.4
1	B	260	GLU	6.0
1	A	10	GLY	5.7
2	D	85	THR	5.1
1	A	260	GLU	5.0
1	A	265	PHE	5.0
2	D	78	ALA	4.5
2	D	80	TYR	4.5
1	A	12	LEU	4.5
2	D	82	LYS	4.2
1	A	261	GLN	4.0
1	A	231	LEU	4.0
1	A	94	GLY	4.0
2	D	77	LEU	3.9
1	B	216	ALA	3.8
1	B	261	GLN	3.8
2	D	83	ARG	3.8
2	D	153	ASN	3.8
1	A	247	PRO	3.8
1	A	190	GLN	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	81	ASN	3.6
2	D	79	HIS	3.6
1	B	294	SER	3.6
1	B	91	GLY	3.4
1	B	5	TYR	3.3
1	B	250	LYS	3.3
2	D	156	GLN	3.3
1	B	92	LEU	3.2
1	A	16	PHE	3.2
1	B	253	ARG	3.1
1	A	93	GLU	3.1
2	D	84	SER	3.0
2	D	88	SER	3.0
1	A	99	GLU	3.0
1	A	151	LYS	3.0
2	D	154	TYR	2.9
1	A	92	LEU	2.9
1	A	11	LEU	2.8
1	B	90	GLU	2.8
2	D	203	ARG	2.8
2	D	128	LYS	2.8
1	B	95	ASP	2.7
1	B	252	GLU	2.7
1	B	293	GLN	2.7
1	A	263	GLU	2.7
2	D	89	ARG	2.7
1	A	152	ASP	2.6
1	A	267	GLU	2.6
1	A	91	GLY	2.6
1	A	295	ASP	2.5
1	B	98	ALA	2.5
2	D	31	LYS	2.5
1	A	95	ASP	2.5
2	D	122	LYS	2.5
1	A	14	THR	2.5
1	B	102	LYS	2.5
1	A	192	ASP	2.4
1	A	266	LEU	2.4
1	A	233	LYS	2.4
1	A	198	ARG	2.3
1	B	245	GLU	2.3
1	A	268	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	153	PRO	2.3
1	A	133	GLU	2.2
1	B	265	PHE	2.2
1	B	4	LEU	2.2
1	B	234	THR	2.2
1	B	103	ALA	2.2
2	D	76	ARG	2.2
1	A	264	GLU	2.1
2	D	157	ARG	2.1
2	D	87	SER	2.1
1	B	94	GLY	2.1
1	A	218	LYS	2.1
1	B	248	LYS	2.1
2	D	215	VAL	2.1
1	B	262	ALA	2.1
1	A	114	LYS	2.0
1	B	236	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.