



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 09:31 PM UTC

PDB ID : 8K0K / pdb_00008k0k
Title : Crystal structure of Csy complex
Authors : Feng, Y.; Wang, H.
Deposited on : 2023-07-09
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)
RNA backbone	3983	1109 (3.20-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	179	3% 77% 20% ..
2	B	248	% 76% 21% .
3	C	306	2% 83% 15% ..
3	D	306	4% 81% 16% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	306	2% 81% 17% ..
3	F	306	3% 80% 17% ..
3	G	306	2% 78% 20% ..
3	H	306	7% 74% 19% 7%
4	I	168	43% 49% 34% 14% ..
5	J	60	10% 27% 42% 30% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 19734 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Csy1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	177	Total	C	N	O	S	0	0	0
			1418	898	235	274	11			

- Molecule 2 is a protein called Csy2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	248	Total	C	N	O	S	0	0	0
			1919	1223	322	360	14			

- Molecule 3 is a protein called Csy3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	304	Total	C	N	O	S	0	0	0
			2333	1468	396	462	7			
3	D	302	Total	C	N	O	S	0	0	0
			2320	1460	393	460	7			
3	E	304	Total	C	N	O	S	0	0	0
			2332	1467	395	462	8			
3	F	303	Total	C	N	O	S	0	0	0
			2328	1465	394	461	8			
3	G	303	Total	C	N	O	S	0	0	0
			2328	1465	394	461	8			
3	H	285	Total	C	N	O	S	0	0	0
			2198	1391	367	432	8			

- Molecule 4 is a protein called Csy4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	I	166	Total	C	N	O	S	0	0	0
			1298	823	222	246	7			


- Molecule 5 is a RNA chain called RNA (60-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	J	60	Total	C	N	O	P	0	0	0
			1260	565	210	425	60			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Csy1




• Molecule 2: Csy2




• Molecule 3: Csy3





• Molecule 3: Csy3









• Molecule 3: Csy3





• Molecule 3: Csy3




• Molecule 3: Csy3





• Molecule 3: Csy3






● Molecule 4: Csy4



● Molecule 5: RNA (60-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	238.85Å 95.08Å 204.03Å 90.00° 93.16° 90.00°	Depositor
Resolution (Å)	46.30 – 3.00 46.30 – 3.00	Depositor EDS
% Data completeness (in resolution range)	92.4 (46.30-3.00) 92.4 (46.30-3.00)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.76 (at 3.01Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.198 , 0.238 0.200 , 0.238	Depositor DCC
R_{free} test set	4261 reflections (4.62%)	wwPDB-VP
Wilson B-factor (Å ²)	57.2	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	19734	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/1451	0.44	0/1959
2	B	0.19	0/1956	0.42	0/2645
3	C	0.20	0/2373	0.43	1/3221 (0.0%)
3	D	0.21	0/2360	0.42	0/3205
3	E	0.18	0/2372	0.39	0/3220
3	F	0.26	0/2368	0.47	0/3215
3	G	0.29	0/2368	0.48	1/3215 (0.0%)
3	H	0.19	0/2236	0.38	0/3035
4	I	0.42	0/1316	0.72	1/1767 (0.1%)
5	J	0.20	0/1404	0.56	1/2181 (0.0%)
All	All	0.24	0/20204	0.46	4/27663 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	130	TRP	CB-CA-C	-6.01	109.62	116.54
4	I	132	PRO	N-CA-CB	-5.56	97.42	103.25
3	G	209	SER	CB-CA-C	-5.47	108.67	115.89
5	J	34	U	P-O3'-C3'	5.36	128.23	120.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1418	0	1360	18	1
2	B	1919	0	1935	40	0
3	C	2333	0	2281	27	0
3	D	2320	0	2265	33	0
3	E	2332	0	2280	33	1
3	F	2328	0	2277	36	0
3	G	2328	0	2277	37	0
3	H	2198	0	2155	36	0
4	I	1298	0	1345	84	0
5	J	1260	0	641	53	0
All	All	19734	0	18816	335	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 335 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:182:MET:HE2	2:B:215:ILE:HG21	1.27	1.08
2:B:182:MET:CE	2:B:215:ILE:HG21	1.83	1.07
2:B:182:MET:HE2	2:B:215:ILE:CG2	1.84	1.07
4:I:1:MET:HG2	4:I:86:ASP:HB2	1.58	0.86
1:A:153:THR:HG22	1:A:156:CYS:H	1.44	0.81

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:GLU:OE1	3:E:146:ASN:OD1[1_565]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	175/179 (98%)	163 (93%)	11 (6%)	1 (1%)	21	56
2	B	246/248 (99%)	227 (92%)	19 (8%)	0	100	100
3	C	302/306 (99%)	282 (93%)	20 (7%)	0	100	100
3	D	300/306 (98%)	280 (93%)	19 (6%)	1 (0%)	36	70
3	E	302/306 (99%)	283 (94%)	19 (6%)	0	100	100
3	F	301/306 (98%)	275 (91%)	25 (8%)	1 (0%)	36	70
3	G	301/306 (98%)	281 (93%)	20 (7%)	0	100	100
3	H	281/306 (92%)	264 (94%)	16 (6%)	1 (0%)	30	65
4	I	164/168 (98%)	135 (82%)	21 (13%)	8 (5%)	1	10
All	All	2372/2431 (98%)	2190 (92%)	170 (7%)	12 (0%)	24	60

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	207	GLU
4	I	87	LYS
4	I	60	GLY
4	I	111	GLY
4	I	123	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	156/157 (99%)	146 (94%)	10 (6%)	16	48
2	B	211/211 (100%)	196 (93%)	15 (7%)	13	43
3	C	249/251 (99%)	235 (94%)	14 (6%)	19	52
3	D	248/251 (99%)	236 (95%)	12 (5%)	23	57
3	E	249/251 (99%)	231 (93%)	18 (7%)	13	43
3	F	249/251 (99%)	237 (95%)	12 (5%)	23	57
3	G	249/251 (99%)	236 (95%)	13 (5%)	21	55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	H	235/251 (94%)	227 (97%)	8 (3%)	32 66
4	I	149/151 (99%)	113 (76%)	36 (24%)	1 4
All	All	1995/2025 (98%)	1857 (93%)	138 (7%)	14 45

5 of 138 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	I	69	THR
4	I	93	LYS
4	I	128	LYS
3	D	245	THR
3	D	202	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
3	E	258	ASN
3	G	61	ASN
4	I	48	ASN
3	F	208	ASN
3	G	63	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
5	J	59/60 (98%)	27 (45%)	5 (8%)

5 of 27 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
5	J	-5	U
5	J	0	A
5	J	1	G
5	J	2	U
5	J	6	C

All (5) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
5	J	33	G
5	J	34	U
5	J	36	A
5	J	37	G
5	J	44	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	177/179 (98%)	0.08	6 (3%) 48 27	25, 44, 74, 100	0
2	B	248/248 (100%)	0.28	3 (1%) 76 55	24, 56, 91, 107	0
3	C	304/306 (99%)	0.37	6 (1%) 65 41	22, 55, 80, 110	0
3	D	302/306 (98%)	0.33	13 (4%) 40 21	23, 48, 82, 107	0
3	E	304/306 (99%)	-0.14	6 (1%) 65 41	21, 38, 67, 119	0
3	F	303/306 (99%)	-0.14	8 (2%) 57 34	21, 35, 70, 95	0
3	G	303/306 (99%)	0.38	7 (2%) 61 38	36, 58, 89, 114	0
3	H	285/306 (93%)	1.01	21 (7%) 20 10	51, 79, 102, 123	0
4	I	166/168 (98%)	2.10	72 (43%) 0 1	70, 94, 139, 141	0
5	J	60/60 (100%)	0.57	6 (10%) 12 7	26, 54, 145, 153	0
All	All	2452/2491 (98%)	0.41	148 (6%) 27 14	21, 54, 100, 153	0

The worst 5 of 148 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	I	145	PHE	6.2
4	I	133	TYR	5.7
4	I	132	PRO	5.5
4	I	137	ASN	5.4
3	H	211	LEU	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.