



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:28 PM UTC

PDB ID : 3K15 / pdb_00003k15
Title : Crystal Structure of BRCA1 BRCT D1840T in complex with a minimal recognition tetrapeptide with an amidated C-terminus
Authors : Campbell, S.J.; Edwards, R.A.; Glover, J.N.
Deposited on : 2009-09-25
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	20% 53% 40% • •
2	B	5	20% 60% 20% 20%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SEP	B	1	-	-	X	-
4	CL	B	6	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 1751 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Breast cancer type 1 susceptibility protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1694	1082	289	309	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1645	MET	-	expression tag	UNP P38398
A	1840	THR	ASP	engineered mutation	UNP P38398

- Molecule 2 is a protein called phospho peptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	5	Total	C	N	O	P	0	0	1
			36	21	5	9	1			

- Molecule 3 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ni	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	16	Total	O	0	0
			16	16		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Breast cancer type 1 susceptibility protein



- Molecule 2: phospho peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	114.57Å 114.57Å 123.19Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.50 – 2.80 37.50 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (37.50-2.80) 99.5 (37.50-2.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.81Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.250 , 0.280 (Not available) , 0.294	Depositor DCC
R_{free} test set	582 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	71.7	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 56.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	1751	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NI, SEP, CL, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/1735	0.82	2/2354 (0.1%)
2	B	0.37	0/26	0.56	0/34
All	All	0.46	0/1761	0.81	2/2388 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1748	GLY	CA-C-N	5.55	125.90	119.47
1	A	1748	GLY	C-N-CA	5.55	125.90	119.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1694	0	1674	104	1
2	B	36	0	28	6	0
3	A	1	0	0	0	0
4	B	1	0	0	5	0
5	A	16	0	0	6	1
5	B	3	0	0	0	0
All	All	1751	0	1702	109	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

All (109) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1649:ARG:CD	1:A:1650:MET:H	1.22	1.52
1:A:1649:ARG:HD3	1:A:1650:MET:N	1.39	1.37
1:A:1855:ILE:CG2	1:A:1856:PRO:HD2	1.74	1.18
1:A:1855:ILE:HG23	1:A:1856:PRO:HD2	1.16	1.12
1:A:1649:ARG:CD	1:A:1650:MET:N	2.04	1.05
1:A:1713:VAL:HG12	1:A:1736:VAL:CG1	1.98	0.94
1:A:1649:ARG:HD3	1:A:1650:MET:H	0.78	0.92
1:A:1658:THR:OG1	1:A:1661:GLU:HG3	1.69	0.92
1:A:1726:ARG:O	1:A:1727:LYS:HG2	1.71	0.91
1:A:1649:ARG:HD2	1:A:1650:MET:H	1.38	0.85
1:A:1855:ILE:CG2	1:A:1856:PRO:CD	2.56	0.83
1:A:1804:VAL:HG12	1:A:1805:HIS:H	1.45	0.82
1:A:1805:HIS:CE1	5:A:19:HOH:O	2.32	0.81
2:B:1:SEP:N	4:B:6:CL:CL	2.50	0.81
1:A:1763:GLY:O	1:A:1764:LEU:HD23	1.81	0.81
1:A:1805:HIS:HE1	5:A:19:HOH:O	1.64	0.80
1:A:1731:GLU:OE1	1:A:1731:GLU:N	2.15	0.78
1:A:1713:VAL:HG12	1:A:1736:VAL:HG11	1.66	0.78
1:A:1805:HIS:CE1	5:A:18:HOH:O	2.38	0.75
1:A:1855:ILE:HG23	1:A:1856:PRO:CD	2.09	0.75
1:A:1649:ARG:CG	1:A:1650:MET:N	2.52	0.73
2:B:1:SEP:O	4:B:6:CL:CL	2.45	0.72
1:A:1759:LYS:O	1:A:1762:ARG:HG3	1.88	0.72
1:A:1855:ILE:HG22	1:A:1856:PRO:HD2	1.69	0.71
1:A:1857:GLN:O	1:A:1859:PRO:HD3	1.91	0.71
1:A:1649:ARG:CG	1:A:1650:MET:H	2.03	0.71
1:A:1730:ASN:C	1:A:1730:ASN:HD22	1.99	0.70
1:A:1726:ARG:O	1:A:1727:LYS:CG	2.40	0.69
1:A:1650:MET:HE3	1:A:1686:HIS:CE1	2.27	0.69
1:A:1816:THR:HG22	1:A:1816:THR:O	1.93	0.68
1:A:1649:ARG:HD3	1:A:1649:ARG:C	2.09	0.67
1:A:1731:GLU:H	1:A:1731:GLU:CD	2.03	0.67
1:A:1802:THR:O	1:A:1803:GLY:C	2.37	0.66
1:A:1776:PRO:HG2	1:A:1779:GLN:CG	2.26	0.66
1:A:1713:VAL:HG12	1:A:1736:VAL:HG13	1.74	0.65
1:A:1773:THR:HG23	1:A:1814:ALA:HB2	1.79	0.63
1:A:1855:ILE:HG22	1:A:1856:PRO:CD	2.29	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1683:GLU:OE2	1:A:1683:GLU:N	2.32	0.62
1:A:1699:ARG:HH11	1:A:1742:ASN:HD21	1.47	0.61
1:A:1804:VAL:HG12	1:A:1805:HIS:N	2.15	0.61
1:A:1785:GLN:NE2	5:A:14:HOH:O	2.33	0.61
1:A:1776:PRO:HG2	1:A:1779:GLN:HG2	1.81	0.60
1:A:1649:ARG:O	1:A:1685:THR:HG21	2.01	0.60
1:A:1761:PHE:HB3	1:A:1789:ALA:HB2	1.84	0.60
1:A:1806:PRO:C	1:A:1807:ILE:HG13	2.27	0.59
1:A:1700:THR:O	1:A:1703:TYR:HB3	2.03	0.58
1:A:1650:MET:HE1	1:A:1718:TRP:HH2	1.69	0.57
1:A:1730:ASN:ND2	1:A:1732:HIS:HB2	2.19	0.57
1:A:1660:GLU:H	1:A:1660:GLU:CD	2.11	0.57
1:A:1653:VAL:HB	5:A:7:HOH:O	2.04	0.56
1:A:1672:HIS:HB2	1:A:1674:ILE:HD12	1.88	0.55
1:A:1832:VAL:O	1:A:1833:VAL:HG13	2.07	0.54
1:A:1785:GLN:HE21	1:A:1791:VAL:HG23	1.73	0.54
1:A:1692:ASP:OD1	1:A:1692:ASP:C	2.52	0.53
1:A:1858:ILE:HG22	1:A:1859:PRO:O	2.08	0.53
1:A:1838:VAL:O	1:A:1842:VAL:HG23	2.09	0.53
1:A:1725:GLU:O	1:A:1726:ARG:C	2.53	0.52
1:A:1694:GLU:HB2	1:A:1744:ARG:NH2	2.25	0.51
1:A:1730:ASN:C	1:A:1730:ASN:ND2	2.65	0.51
1:A:1730:ASN:HD21	1:A:1732:HIS:HB2	1.74	0.51
1:A:1722:SER:HB3	1:A:1727:LYS:O	2.12	0.50
1:A:1752:ALA:C	1:A:1754:GLU:N	2.68	0.50
1:A:1759:LYS:CB	1:A:1787:CYS:O	2.59	0.50
1:A:1704:PHE:CZ	1:A:1839:LEU:HB3	2.48	0.49
1:A:1773:THR:C	1:A:1775:MET:H	2.20	0.49
1:A:1750:LYS:O	1:A:1754:GLU:HG3	2.13	0.49
1:A:1761:PHE:HB2	1:A:1787:CYS:O	2.12	0.49
1:A:1810:VAL:O	1:A:1834:THR:HA	2.11	0.49
1:A:1850:LEU:O	1:A:1851:ASP:C	2.55	0.49
1:A:1759:LYS:HB2	1:A:1787:CYS:O	2.13	0.48
1:A:1752:ALA:O	1:A:1754:GLU:N	2.46	0.48
2:B:1:SEP:CA	4:B:6:CL:CL	2.99	0.48
1:A:1761:PHE:CD1	1:A:1787:CYS:HB2	2.49	0.47
1:A:1763:GLY:C	1:A:1764:LEU:HD23	2.40	0.47
1:A:1665:VAL:O	1:A:1668:PHE:HB3	2.13	0.47
1:A:1699:ARG:HB2	2:B:4:PHE:O	2.13	0.47
1:A:1722:SER:CB	1:A:1727:LYS:O	2.63	0.47
1:A:1766:ILE:HG13	1:A:1789:ALA:HB1	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1714:VAL:HG11	1:A:1731:GLU:HB3	1.96	0.47
1:A:1658:THR:HB	1:A:1660:GLU:OE2	2.15	0.46
1:A:1664:LEU:HD23	1:A:1664:LEU:HA	1.76	0.46
1:A:1729:LEU:O	1:A:1730:ASN:C	2.58	0.46
1:A:1816:THR:O	1:A:1816:THR:CG2	2.62	0.46
1:A:1752:ALA:O	1:A:1753:ARG:C	2.59	0.45
1:A:1759:LYS:O	1:A:1761:PHE:N	2.49	0.45
2:B:1:SEP:C	4:B:6:CL:CL	3.02	0.45
2:B:1:SEP:HB3	4:B:6:CL:CL	2.53	0.45
1:A:1811:GLN:HE21	1:A:1813:ASP:HB2	1.81	0.45
1:A:1855:ILE:HG22	1:A:1856:PRO:N	2.32	0.44
1:A:1832:VAL:O	1:A:1833:VAL:CG1	2.65	0.44
1:A:1752:ALA:C	1:A:1754:GLU:H	2.25	0.44
1:A:1759:LYS:C	1:A:1761:PHE:N	2.75	0.43
1:A:1658:THR:HG1	1:A:1661:GLU:HG3	1.76	0.43
1:A:1675:THR:HG22	1:A:1676:LEU:N	2.34	0.42
1:A:1696:VAL:HG21	1:A:1744:ARG:HG2	2.01	0.42
1:A:1759:LYS:C	1:A:1761:PHE:H	2.26	0.42
1:A:1765:GLU:OE2	1:A:1799:THR:HB	2.18	0.42
1:A:1679:LEU:HA	1:A:1679:LEU:HD23	1.71	0.42
1:A:1747:GLN:OE1	1:A:1747:GLN:CA	2.67	0.42
1:A:1750:LYS:O	1:A:1751:ARG:C	2.63	0.42
1:A:1841:SER:HA	1:A:1848:GLN:HG2	2.02	0.42
1:A:1759:LYS:HB3	1:A:1787:CYS:O	2.20	0.41
1:A:1837:TRP:O	1:A:1841:SER:OG	2.29	0.41
1:A:1727:LYS:HA	1:A:1727:LYS:HD3	1.69	0.41
1:A:1759:LYS:H	1:A:1759:LYS:HG2	1.69	0.41
1:A:1806:PRO:O	1:A:1831:PRO:HD2	2.20	0.41
1:A:1846:GLN:HE21	1:A:1846:GLN:HB2	1.52	0.41
1:A:1653:VAL:HA	5:A:7:HOH:O	2.20	0.40
1:A:1804:VAL:CG1	1:A:1805:HIS:N	2.84	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1673:HIS:NE2	5:A:18:HOH:O[6_555]	1.82	0.38

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/215 (97%)	178 (85%)	26 (12%)	5 (2%)	4	17
2	B	3/5 (60%)	3 (100%)	0	0	100	100
All	All	212/220 (96%)	181 (85%)	26 (12%)	5 (2%)	4	17

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1803	GLY
1	A	1847	CYS
1	A	1758	ARG
1	A	1748	GLY
1	A	1760	ILE

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/192 (98%)	179 (95%)	9 (5%)	23	56
2	B	3/3 (100%)	3 (100%)	0	100	100
All	All	191/195 (98%)	182 (95%)	9 (5%)	23	57

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1660	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1714	VAL
1	A	1730	ASN
1	A	1736	VAL
1	A	1794	GLU
1	A	1799	THR
1	A	1804	VAL
1	A	1819	ASN
1	A	1846	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1721	GLN
1	A	1730	ASN
1	A	1742	ASN
1	A	1774	ASN
1	A	1779	GLN
1	A	1785	GLN
1	A	1846	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SEP	B	1	2	8,9,10	1.51	1 (12%)	7,12,14	1.61	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SEP	B	1	2	-	5/6/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	SEP	P-O1P	3.41	1.61	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	SEP	OG-CB-CA	2.30	110.39	108.14

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	SEP	N-CA-CB-OG
2	B	1	SEP	C-CA-CB-OG
2	B	1	SEP	CB-OG-P-O1P
2	B	1	SEP	CB-OG-P-O3P
2	B	1	SEP	CB-OG-P-O2P

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	SEP	5	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	211/215 (98%)	1.19	44 (20%) 2 2	31, 48, 62, 70	1 (0%)
2	B	3/5 (60%)	1.58	1 (33%) 1 1	47, 47, 48, 48	3 (100%)
All	All	214/220 (97%)	1.20	45 (21%) 2 2	31, 48, 62, 70	4 (1%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1820	GLY	5.2
1	A	1816	THR	5.1
1	A	1858	ILE	4.8
1	A	1859	PRO	3.6
1	A	1856	PRO	3.6
1	A	1851	ASP	3.2
1	A	1855	ILE	3.1
1	A	1686	HIS	3.0
1	A	1857	GLN	2.8
1	A	1734	PHE	2.8
1	A	1822	HIS	2.8
1	A	1849	GLU	2.7
1	A	1663	MET	2.7
1	A	1823	ALA	2.6
1	A	1815	TRP	2.5
1	A	1650	MET	2.5
1	A	1802	THR	2.5
1	A	1733	ASP	2.5
1	A	1812	PRO	2.5
1	A	1848	GLN	2.5
1	A	1824	ILE	2.4
1	A	1717	PHE	2.4
1	A	1821	PHE	2.4
1	A	1712	TRP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	4	PHE	2.4
1	A	1742	ASN	2.4
1	A	1825	GLY	2.4
1	A	1846	GLN	2.4
1	A	1850	LEU	2.4
1	A	1730	ASN	2.3
1	A	1819	ASN	2.3
1	A	1682	GLU	2.3
1	A	1813	ASP	2.3
1	A	1710	GLY	2.3
1	A	1729	LEU	2.2
1	A	1854	LEU	2.1
1	A	1760	ILE	2.1
1	A	1811	GLN	2.1
1	A	1732	HIS	2.1
1	A	1741	VAL	2.1
1	A	1833	VAL	2.1
1	A	1769	TYR	2.0
1	A	1651	SER	2.0
1	A	1656	GLY	2.0
1	A	1832	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SEP	B	1	10/11	0.88	0.18	50,53,57,57	10

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NI	A	1	1/1	0.80	0.21	83,83,83,83	1
4	CL	B	6	1/1	0.84	0.31	74,74,74,74	1

6.5 Other polymers [i](#)

There are no such residues in this entry.