



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 04:55 PM UTC

PDB ID : 5K1V / pdb_00005k1v
Title : Crystal structure of Endoplasmic Reticulum aminopeptidase 2 (ERAP2) in complex with a diaminobenzoic acid derivative ligand.
Authors : Saridakis, E.; Papakyriakou, A.; Giastas, P.; Mpakali, A.; Mavridis, I.M.; Stratikos, E.
Deposited on : 2016-05-18
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	967	0% 59% 29% • 7%
1	B	967	2% 52% 30% 5% 12%
2	C	2	100%
2	D	2	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	E	4	 A horizontal bar chart showing the quality of chain 3. The bar is divided into two segments: a yellow segment on the left representing 25% and an orange segment on the right representing 75%. The labels '25%' and '75%' are positioned below their respective segments.

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14637 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoplasmic reticulum aminopeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	900	Total	C	N	O	S	0	4	1
			7345	4732	1225	1356	32			
1	B	849	Total	C	N	O	S	0	2	0
			6937	4482	1147	1281	27			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	961	ARG	-	expression tag	UNP Q6P179
A	962	HIS	-	expression tag	UNP Q6P179
A	963	HIS	-	expression tag	UNP Q6P179
A	964	HIS	-	expression tag	UNP Q6P179
A	965	HIS	-	expression tag	UNP Q6P179
A	966	HIS	-	expression tag	UNP Q6P179
A	967	HIS	-	expression tag	UNP Q6P179
B	961	ARG	-	expression tag	UNP Q6P179
B	962	HIS	-	expression tag	UNP Q6P179
B	963	HIS	-	expression tag	UNP Q6P179
B	964	HIS	-	expression tag	UNP Q6P179
B	965	HIS	-	expression tag	UNP Q6P179
B	966	HIS	-	expression tag	UNP Q6P179
B	967	HIS	-	expression tag	UNP Q6P179

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		

- Molecule 5 is methyl N-[4-amino-3-(L-arginylamino)benzene-1-carbonyl]-L-tyrosinate (CCD ID: 6PX) (formula: C₂₃H₃₁N₇O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			35	23	7	5		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	30	Total	O	0	0
			30	30		
7	B	14	Total	O	0	0
			14	14		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Endoplasmic reticulum aminopeptidase 2



- Molecule 1: Endoplasmic reticulum aminopeptidase 2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 

MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1
MAG2

- Molecule 3: alpha-D-mannopyranose-(1-4)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  25% 75%

MAG1
MAG2
MAN3
MAN4

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.40Å 135.17Å 127.49Å 90.00° 90.15° 90.00°	Depositor
Resolution (Å)	48.34 – 2.90 48.34 – 2.90	Depositor EDS
% Data completeness (in resolution range)	98.3 (48.34-2.90) 99.6 (48.34-2.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.91Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.205 , 0.273 0.215 , 0.280	Depositor DCC
R_{free} test set	2793 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	76.4	Xtriage
Anisotropy	0.239	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14637	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ZN, MAN, 6PX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/7537	1.00	18/10210 (0.2%)
1	B	0.55	0/7114	0.92	11/9636 (0.1%)
All	All	0.63	0/14651	0.96	29/19846 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	55	VAL	N-CA-C	8.81	119.34	108.06
1	A	889	LEU	N-CA-C	7.70	119.31	111.07
1	B	62	PHE	CA-C-N	6.57	128.06	119.84
1	B	62	PHE	C-N-CA	6.57	128.06	119.84
1	A	177	GLU	N-CA-C	6.52	119.09	109.24

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	54	PRO	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7345	0	7282	198	0
1	B	6937	0	6887	256	0
2	C	28	0	25	1	0
2	D	28	0	25	0	0
3	E	50	0	43	2	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	35	0	0	3	0
6	A	84	0	78	0	0
6	B	84	0	78	2	0
7	A	30	0	0	3	0
7	B	14	0	0	0	0
All	All	14637	0	14418	453	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 453 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:592:TYR:HE1	1:B:601:HIS:HB2	1.13	1.07
1:B:599:VAL:HA	1:B:600:ILE:HG13	1.38	1.01
1:B:599:VAL:HG13	1:B:600:ILE:HB	1.42	0.98
1:B:599:VAL:HA	1:B:600:ILE:CG1	1.92	0.98
1:A:383:MET:H	1:A:383:MET:HE3	1.32	0.95

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	898/967 (93%)	788 (88%)	96 (11%)	14 (2%)	7	27
1	B	841/967 (87%)	725 (86%)	105 (12%)	11 (1%)	9	32
All	All	1739/1934 (90%)	1513 (87%)	201 (12%)	25 (1%)	9	30

5 of 25 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	176	PHE
1	A	582	GLU
1	A	715	ILE
1	B	140	LEU
1	B	216	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	812/870 (93%)	738 (91%)	74 (9%)	9	28
1	B	764/870 (88%)	699 (92%)	65 (8%)	10	31
All	All	1576/1740 (91%)	1437 (91%)	139 (9%)	9	29

5 of 139 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	592	TYR
1	B	622	LYS
1	B	848	LYS
1	A	674	ARG
1	A	660	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	447	GLN
1	B	568	GLN
1	B	959	ASN
1	B	879	ASN
1	A	708	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	2,1	14,14,15	0.47	0	17,19,21	0.39	0
2	NAG	C	2	2	14,14,15	0.30	0	17,19,21	0.69	1 (5%)
2	NAG	D	1	2,1	14,14,15	0.40	0	17,19,21	1.61	3 (17%)
2	NAG	D	2	2	14,14,15	1.12	1 (7%)	17,19,21	1.20	1 (5%)
3	NAG	E	1	1,3	14,14,15	1.15	1 (7%)	17,19,21	0.88	1 (5%)
3	NAG	E	2	3	14,14,15	0.90	1 (7%)	17,19,21	1.04	1 (5%)
3	MAN	E	3	3	11,11,12	1.31	1 (9%)	15,15,17	1.31	2 (13%)
3	MAN	E	4	3	11,11,12	1.31	1 (9%)	15,15,17	0.92	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	3/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
3	NAG	E	1	1,3	-	4/6/23/26	0/1/1/1
3	NAG	E	2	3	-	1/6/23/26	0/1/1/1
3	MAN	E	3	3	-	2/2/19/22	0/1/1/1
3	MAN	E	4	3	-	2/2/19/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2	NAG	C1-C2	3.99	1.57	1.52
3	E	1	NAG	O5-C1	-3.73	1.37	1.43
3	E	4	MAN	O5-C1	-3.49	1.37	1.43
3	E	2	NAG	O5-C1	3.20	1.49	1.43
3	E	3	MAN	C4-C5	2.76	1.58	1.53

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	NAG	C1-O5-C5	4.14	117.74	112.19
2	D	2	NAG	C1-O5-C5	3.99	117.53	112.19
2	D	1	NAG	O4-C4-C5	3.71	118.47	109.32
3	E	3	MAN	C1-O5-C5	3.54	116.94	112.19
3	E	2	NAG	C1-O5-C5	3.31	116.62	112.19

There are no chirality outliers.

5 of 16 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	O5-C5-C6-O6
3	E	3	MAN	O5-C5-C6-O6
3	E	4	MAN	O5-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6


There are no ring outliers.


4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	2	NAG	1	0
2	C	1	NAG	1	0
3	E	3	MAN	1	0
3	E	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	6PX	A	1002	4	36,36,36	2.25	6 (16%)	46,48,48	1.77	17 (36%)
6	NAG	B	1011	1	14,14,15	1.56	2 (14%)	17,19,21	1.42	1 (5%)
6	NAG	B	1010	1	14,14,15	0.33	0	17,19,21	0.41	0
6	NAG	B	1009	1	14,14,15	0.41	0	17,19,21	0.87	1 (5%)
6	NAG	A	1009	1	14,14,15	0.45	0	17,19,21	0.59	0
6	NAG	A	1007	1	14,14,15	0.53	0	17,19,21	0.52	0
6	NAG	A	1011	1	14,14,15	0.24	0	17,19,21	0.42	0
6	NAG	A	1008	1	14,14,15	0.65	1 (7%)	17,19,21	0.51	0
6	NAG	A	1012	1	14,14,15	0.84	1 (7%)	17,19,21	0.60	0
6	NAG	B	1002	1	14,14,15	0.60	0	17,19,21	0.66	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	B	1007	1	14,14,15	0.61	1 (7%)	17,19,21	0.40	0
6	NAG	B	1008	1	14,14,15	0.81	1 (7%)	17,19,21	0.51	0
6	NAG	A	1010	1	14,14,15	0.98	1 (7%)	17,19,21	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	6PX	A	1002	4	-	16/33/33/33	0/2/2/2
6	NAG	B	1011	1	-	2/6/23/26	0/1/1/1
6	NAG	B	1010	1	-	2/6/23/26	0/1/1/1
6	NAG	B	1009	1	-	2/6/23/26	0/1/1/1
6	NAG	A	1009	1	-	0/6/23/26	0/1/1/1
6	NAG	A	1007	1	-	2/6/23/26	0/1/1/1
6	NAG	A	1011	1	-	2/6/23/26	0/1/1/1
6	NAG	A	1008	1	-	3/6/23/26	0/1/1/1
6	NAG	A	1012	1	-	1/6/23/26	0/1/1/1
6	NAG	B	1002	1	-	3/6/23/26	0/1/1/1
6	NAG	B	1007	1	-	2/6/23/26	0/1/1/1
6	NAG	B	1008	1	-	2/6/23/26	0/1/1/1
6	NAG	A	1010	1	-	0/6/23/26	0/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1002	6PX	C6-N2	8.05	1.48	1.33
5	A	1002	6PX	C7-N7	5.55	1.47	1.34
5	A	1002	6PX	C2-N5	5.35	1.47	1.35
6	B	1011	NAG	O5-C1	5.27	1.52	1.43
5	A	1002	6PX	O5-C15	4.02	1.43	1.33

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	1011	NAG	C1-O5-C5	5.69	119.81	112.19
5	A	1002	6PX	C12-C10-C11	-4.63	115.85	120.80
5	A	1002	6PX	C11-C7-N7	-3.81	109.97	117.04

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1002	6PX	C18-C20-C22	-3.34	116.35	119.88
6	B	1009	NAG	C1-O5-C5	2.98	116.18	112.19

There are no chirality outliers.

5 of 37 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	1002	6PX	C3-C1-C2-O1
5	A	1002	6PX	C3-C1-C2-N5
5	A	1002	6PX	O2-C7-N7-C14
5	A	1002	6PX	N7-C14-C16-C17
5	A	1002	6PX	C11-C7-N7-C14

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1002	6PX	3	0
6	B	1002	NAG	1	0
6	B	1007	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	900/967 (93%)	-0.36	5 (0%) 85 81	24, 55, 99, 147	4 (0%)
1	B	849/967 (87%)	0.31	20 (2%) 59 50	26, 96, 133, 156	2 (0%)
All	All	1749/1934 (90%)	-0.04	25 (1%) 73 65	24, 70, 127, 156	6 (0%)

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	54	PRO	4.2
1	B	640	TRP	3.8
1	B	925	GLY	3.2
1	B	592	TYR	3.2
1	B	623	PHE	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MAN	E	3	11/12	0.51	0.13	112,137,143,143	0
3	MAN	E	4	11/12	0.54	0.12	140,150,157,158	0
2	NAG	D	2	14/15	0.68	0.12	77,86,98,99	0
3	NAG	E	2	14/15	0.84	0.10	57,88,107,124	0
2	NAG	C	2	14/15	0.86	0.09	86,93,99,101	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	E	1	14/15	0.90	0.09	50,63,74,80	0
2	NAG	D	1	14/15	0.93	0.07	42,55,66,71	0
2	NAG	C	1	14/15	0.96	0.05	48,66,86,86	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	NAG	A	1012	14/15	0.36	0.13	90,117,127,130	0
6	NAG	B	1011	14/15	0.42	0.14	103,119,134,137	0
6	NAG	B	1008	14/15	0.48	0.13	124,136,145,146	0
6	NAG	A	1010	14/15	0.50	0.12	107,119,131,136	0
6	NAG	B	1010	14/15	0.51	0.12	108,118,127,127	0
6	NAG	B	1002	14/15	0.51	0.10	114,127,135,135	0
6	NAG	A	1008	14/15	0.55	0.13	105,114,120,130	0
6	NAG	A	1009	14/15	0.59	0.13	96,107,123,125	0
6	NAG	B	1009	14/15	0.64	0.12	100,119,123,124	0
6	NAG	B	1007	14/15	0.72	0.12	94,114,129,131	0
6	NAG	A	1011	14/15	0.81	0.08	99,107,116,119	0
5	6PX	A	1002	35/35	0.81	0.19	38,65,89,92	0
6	NAG	A	1007	14/15	0.91	0.10	84,92,96,98	0
4	ZN	B	1001	1/1	1.00	0.05	54,54,54,54	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ZN	A	1001	1/1	1.00	0.05	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.