



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:41 PM UTC

PDB ID : 6K17 / pdb_00006k17
Title : Crystal structure of EXD2 exonuclease domain
Authors : Park, J.; Lee, C.
Deposited on : 2019-05-10
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	12% 82% 9% 9%
1	B	220	15% 78% 11% 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3666 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exonuclease 3'-5' domain-containing protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	200	Total	C	N	O	S	0	0	0
			1547	987	262	288	10			
1	B	197	Total	C	N	O	S	0	0	0
			1522	972	256	284	10			

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Na	0	0
			2	2		
2	B	2	Total	Na	0	0
			2	2		



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	304	Total	O	0	0
			304	304		
3	B	289	Total	O	0	0
			289	289		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Exonuclease 3'-5' domain-containing protein 2



- Molecule 1: Exonuclease 3'-5' domain-containing protein 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.95Å 72.61Å 135.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.33 – 1.60 37.33 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (37.33-1.60) 99.5 (37.33-1.60)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.192 , 0.220 (Not available) , 0.253	Depositor DCC
R_{free} test set	2000 reflections (3.30%)	wwPDB-VP
Wilson B-factor (Å ²)	16.5	Xtriage
Anisotropy	0.279	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3666	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.24 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.3606e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/1569	0.80	0/2127
1	B	0.63	0/1544	0.79	0/2094
All	All	0.62	0/3113	0.79	0/4221

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1547	0	1597	25	0
1	B	1522	0	1568	32	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	304	0	0	5	3
3	B	289	0	0	8	3
All	All	3666	0	3165	52	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:224:LEU:CD2	3:B:520:HOH:O	2.07	1.02
1:A:217:PHE:HZ	1:A:249:ILE:HD12	1.34	0.91
1:B:224:LEU:HG	1:B:226:ARG:H	1.38	0.87
1:A:96:ARG:HH21	1:A:155:ILE:HG13	1.46	0.81
1:B:224:LEU:HD11	1:B:242:TYR:CE1	2.18	0.79
1:A:140:LYS:HD2	3:A:672:HOH:O	1.82	0.79
1:B:224:LEU:CD1	1:B:242:TYR:CE1	2.75	0.69
1:B:224:LEU:HD11	1:B:242:TYR:HE1	1.58	0.68
1:B:224:LEU:CD1	1:B:242:TYR:HE1	2.06	0.67
1:A:255:LEU:HD11	1:B:129:PRO:HB3	1.74	0.67
1:B:224:LEU:HD22	3:B:520:HOH:O	1.81	0.66
1:A:96:ARG:HH21	1:A:155:ILE:CG1	2.10	0.64
1:A:259:GLY:O	1:A:260:TYR:HB2	1.99	0.62
1:A:258:LEU:HD13	1:A:280:VAL:HG13	1.80	0.61
1:A:110:GLU:HG2	1:A:125:GLN:HE22	1.65	0.60
1:B:96:ARG:NH2	1:B:154:ASP:OD2	2.32	0.60
1:B:245:ARG:O	1:B:249:ILE:HG12	2.03	0.59
1:A:129:PRO:HB3	1:B:255:LEU:HD11	1.84	0.59
1:A:217:PHE:CZ	1:A:249:ILE:HD12	2.27	0.58
1:B:94:LEU:O	1:B:98:GLU:HG3	2.03	0.58
1:B:254:PHE:CZ	1:B:258:LEU:HD12	2.39	0.57
1:A:223:LEU:HD12	3:A:546:HOH:O	2.05	0.57
1:B:224:LEU:HD21	3:B:520:HOH:O	1.91	0.55
1:A:96:ARG:NH2	3:A:410:HOH:O	2.41	0.54
1:A:140:LYS:CD	3:A:672:HOH:O	2.48	0.53
1:B:278:ARG:HD3	3:B:526:HOH:O	2.09	0.53
1:B:224:LEU:HD21	1:B:226:ARG:HB2	1.91	0.51
1:B:285:GLN:HA	3:B:573:HOH:O	2.10	0.50
1:A:217:PHE:HZ	1:A:249:ILE:CD1	2.15	0.50
1:A:96:ARG:NH2	1:A:155:ILE:CG1	2.74	0.50
1:A:260:TYR:CE2	1:B:256:HIS:HE1	2.29	0.50
1:B:224:LEU:HD11	1:B:242:TYR:CZ	2.47	0.49
1:A:281:LEU:O	1:A:285:GLN:HG3	2.12	0.49
1:B:254:PHE:CZ	1:B:258:LEU:CD1	2.94	0.49
1:B:224:LEU:HD12	1:B:242:TYR:CE1	2.49	0.48
1:A:260:TYR:HE2	1:B:256:HIS:CE1	2.32	0.48
1:B:243:ALA:O	3:B:401:HOH:O	2.20	0.47
1:A:140:LYS:HD3	3:A:612:HOH:O	2.15	0.45
1:A:110:GLU:HG2	1:A:125:GLN:NE2	2.32	0.45
1:A:283:LYS:HB3	1:A:283:LYS:HE3	1.76	0.45
1:B:224:LEU:HD11	1:B:242:TYR:OH	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:TYR:CE2	1:B:256:HIS:CE1	3.04	0.45
1:B:254:PHE:CE1	1:B:258:LEU:CD1	3.00	0.43
1:B:92:GLU:HB3	1:B:93:PRO:HD3	2.00	0.42
1:B:258:LEU:HD13	1:B:280:VAL:HG13	2.01	0.42
1:A:125:GLN:HG3	1:A:135:LEU:HD22	2.02	0.42
1:A:259:GLY:O	1:A:260:TYR:CB	2.68	0.42
1:A:96:ARG:NH2	1:A:155:ILE:HG12	2.35	0.41
1:B:217:PHE:CD1	1:B:218:PRO:O	2.73	0.41
1:B:212:GLU:HG3	3:B:518:HOH:O	2.21	0.41
1:B:281:LEU:O	1:B:285:GLN:HG3	2.21	0.40
1:B:98:GLU:HB3	3:B:473:HOH:O	2.21	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:543:HOH:O	3:B:594:HOH:O[2_564]	2.09	0.11
3:A:656:HOH:O	3:B:676:HOH:O[3_646]	2.14	0.06
3:B:520:HOH:O	3:B:583:HOH:O[2_664]	2.17	0.03
3:A:637:HOH:O	3:A:689:HOH:O[2_564]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/220 (88%)	194 (100%)	0	0	100	100
1	B	191/220 (87%)	191 (100%)	0	0	100	100
All	All	385/440 (88%)	385 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/193 (90%)	173 (99%)	1 (1%)	78	66
1	B	171/193 (89%)	169 (99%)	2 (1%)	63	43
All	All	345/386 (89%)	342 (99%)	3 (1%)	70	55

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	78	LYS
1	B	125	GLN
1	B	280	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	248	GLN
1	B	256	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	200/220 (90%)	0.66	27 (13%) 7 6	10, 19, 35, 48	0
1	B	197/220 (89%)	0.75	33 (16%) 4 3	10, 18, 38, 56	0
All	All	397/440 (90%)	0.70	60 (15%) 5 5	10, 18, 36, 56	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	217	PHE	8.2
1	A	223	LEU	7.4
1	A	260	TYR	7.3
1	B	260	TYR	6.8
1	B	279	LYS	6.7
1	A	217	PHE	6.6
1	B	278	ARG	6.0
1	B	224	LEU	5.9
1	A	293	ARG	5.5
1	B	101	ASP	5.3
1	A	294	SER	5.3
1	A	77	ALA	5.1
1	A	102	PHE	5.1
1	B	275	SER	4.8
1	B	102	PHE	4.6
1	A	224	LEU	4.4
1	A	278	ARG	4.4
1	B	282	GLU	4.4
1	A	279	LYS	4.4
1	B	245	ARG	4.2
1	A	282	GLU	4.1
1	B	98	GLU	4.1
1	A	100	GLU	4.0
1	B	100	GLU	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	101	ASP	4.0
1	A	276	SER	3.7
1	B	218	PRO	3.7
1	A	275	SER	3.7
1	B	130	SER	3.5
1	B	285	GLN	3.5
1	B	276	SER	3.5
1	A	285	GLN	3.3
1	A	109	CYS	3.2
1	A	202	CYS	3.1
1	B	280	VAL	3.1
1	B	258	LEU	3.1
1	A	114	LEU	3.0
1	B	290	ILE	3.0
1	B	131	GLY	3.0
1	B	77	ALA	2.8
1	B	292	PHE	2.7
1	A	218	PRO	2.7
1	B	96	ARG	2.6
1	B	243	ALA	2.6
1	A	201	LEU	2.5
1	B	94	LEU	2.5
1	B	132	LEU	2.5
1	B	97	SER	2.4
1	A	117	LYS	2.4
1	A	205	LEU	2.3
1	B	170	GLU	2.3
1	A	258	LEU	2.2
1	B	277	TRP	2.2
1	A	245	ARG	2.2
1	A	259	GLY	2.1
1	B	99	LEU	2.1
1	A	280	VAL	2.1
1	B	281	LEU	2.1
1	B	143	CYS	2.0
1	B	114	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NA	B	301	1/1	0.89	0.25	49,49,49,49	0
2	NA	A	301	1/1	0.92	0.08	30,30,30,30	0
2	NA	A	302	1/1	0.98	0.03	12,12,12,12	0
2	NA	B	302	1/1	0.98	0.06	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around NA B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NA A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NA A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around NA B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.