



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 07:07 PM EDT

PDB ID : 3K23 / pdb_00003k23
Title : Glucocorticoid Receptor with Bound D-prolinamide 11
Authors : Biggadike, K.B.; McLay, I.M.; Madauss, K.P.; Williams, S.P.; Bledsoe, R.K.
Deposited on : 2009-09-29
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	259	66% 25% • 5%
1	B	259	71% 22% • 5%
1	C	259	4% 64% 29% • 5%
2	D	12	58% 8% 8% 25%
2	E	12	50% 33% 17%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	12	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6158 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucocorticoid receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	0	0	0
			1927	1249	313	350	15			
1	B	247	Total	C	N	O	S	0	0	0
			1966	1278	321	350	17			
1	C	246	Total	C	N	O	S	0	0	0
			1859	1212	308	325	14			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	519	GLY	-	expression tag	UNP P04150
A	520	SER	-	expression tag	UNP P04150
A	602	TYR	PHE	engineered mutation	UNP P04150
A	638	GLY	CYS	engineered mutation	UNP P04150
B	519	GLY	-	expression tag	UNP P04150
B	520	SER	-	expression tag	UNP P04150
B	602	TYR	PHE	engineered mutation	UNP P04150
B	638	GLY	CYS	engineered mutation	UNP P04150
C	519	GLY	-	expression tag	UNP P04150
C	520	SER	-	expression tag	UNP P04150
C	602	TYR	PHE	engineered mutation	UNP P04150
C	638	GLY	CYS	engineered mutation	UNP P04150

- Molecule 2 is a protein called Nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	9	Total	C	N	O	0	0	0
			72	49	10	13			
2	E	10	Total	C	N	O	0	0	0
			74	49	11	14			
2	F	10	Total	C	N	O	0	0	0
			72	49	11	12			


- Molecule 3 is 1-{{3-(4-{{[(2R)-4-(5-fluoro-2-methoxyphenyl)-2-hydroxy-4-methyl-2-(trifluoromethyl)pentyl]amino}-6-methyl-1H-indazol-1-yl]phenyl]carbonyl}-D-prolinamide (CCD ID: JZN) (formula: C₃₄H₃₇F₄N₅O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			47	34	4	5	4		
3	B	1	Total	C	F	N	O	0	0
			47	34	4	5	4		
3	C	1	Total	C	F	N	O	0	0
			47	34	4	5	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	18	Total	O	0	0
			18	18		
4	B	18	Total	O	0	0
			18	18		
4	C	11	Total	O	0	0
			11	11		



- Molecule 2: Nuclear receptor coactivator 2



- Molecule 2: Nuclear receptor coactivator 2



- Molecule 2: Nuclear receptor coactivator 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	184.93Å 65.96Å 71.53Å 90.00° 103.64° 90.00°	Depositor
Resolution (Å)	19.82 – 3.00 19.82 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.82-3.00) 98.8 (19.82-3.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.98Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.224 , 0.289 0.233 , 0.237	Depositor DCC
R_{free} test set	1256 reflections (7.48%)	wwPDB-VP
Wilson B-factor (Å ²)	66.6	Xtriage
Anisotropy	0.036	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 82.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6158	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: JZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	0/1967	0.79	1/2667 (0.0%)
1	B	0.58	0/2010	0.79	0/2726
1	C	0.52	0/1896	0.78	1/2580 (0.0%)
2	D	0.48	0/72	0.67	0/95
2	E	0.66	0/74	0.92	0/98
2	F	0.47	0/72	0.69	0/98
All	All	0.57	0/6091	0.79	2/8264 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	553	PRO	N-CA-CB	7.86	110.17	103.25
1	C	526	PRO	N-CA-CB	6.33	110.23	103.52

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1927	0	1871	64	0
1	B	1966	0	1955	50	0
1	C	1859	0	1779	56	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	72	0	76	2	0
2	E	74	0	69	10	0
2	F	72	0	69	3	0
3	A	47	0	37	14	0
3	B	47	0	37	9	0
3	C	47	0	37	9	0
4	A	18	0	0	0	0
4	B	18	0	0	2	0
4	C	11	0	0	0	0
All	All	6158	0	5930	186	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 186 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:621:LEU:HD21	1:A:646:MET:HE3	1.47	0.94
1:B:570:GLN:HB3	1:B:604:MET:HE1	1.51	0.93
1:C:737:PHE:CE1	1:C:757:ILE:HG23	2.08	0.89
1:B:570:GLN:HB3	1:B:604:MET:CE	2.01	0.89
1:A:567:GLY:HA2	1:A:604:MET:HE1	1.57	0.86

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	237/259 (92%)	215 (91%)	19 (8%)	3 (1%)	9 38
1	B	243/259 (94%)	227 (93%)	15 (6%)	1 (0%)	30 65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	238/259 (92%)	208 (87%)	20 (8%)	10 (4%)	2	13
2	D	7/12 (58%)	7 (100%)	0	0	100	100
2	E	8/12 (67%)	7 (88%)	1 (12%)	0	100	100
2	F	8/12 (67%)	8 (100%)	0	0	100	100
All	All	741/813 (91%)	672 (91%)	55 (7%)	14 (2%)	6	30

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	552	VAL
1	A	617	SER
1	C	617	SER
1	C	748	GLU
1	C	637	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/236 (86%)	188 (93%)	15 (7%)	13	42
1	B	212/236 (90%)	202 (95%)	10 (5%)	23	58
1	C	182/236 (77%)	178 (98%)	4 (2%)	45	74
2	D	7/11 (64%)	6 (86%)	1 (14%)	3	16
2	E	6/11 (54%)	6 (100%)	0	100	100
2	F	6/11 (54%)	6 (100%)	0	100	100
All	All	616/741 (83%)	586 (95%)	30 (5%)	22	56

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	776	GLN
1	C	570	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	549	ASP
1	C	733	LEU
1	B	756	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	760	GLN
1	A	776	GLN
1	C	592	GLN
1	B	570	GLN
1	A	707	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	JZN	C	3	-	50,51,51	1.28	4 (8%)	71,78,78	2.06	20 (28%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	JZN	A	1	-	50,51,51	1.44	7 (14%)	71,78,78	2.62	27 (38%)
3	JZN	B	2	-	50,51,51	1.44	7 (14%)	71,78,78	2.99	29 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	JZN	C	3	-	-	29/45/55/55	0/5/5/5
3	JZN	A	1	-	-	35/45/55/55	0/5/5/5
3	JZN	B	2	-	-	20/45/55/55	0/5/5/5

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2	JZN	F3-C34	4.47	1.48	1.33
3	A	1	JZN	N2-N3	-3.66	1.34	1.38
3	A	1	JZN	C21-C14	-3.57	1.38	1.41
3	A	1	JZN	C21-C19	-3.44	1.38	1.41
3	C	3	JZN	C8-C7	-3.43	1.49	1.54

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2	JZN	F3-C34-C12	-11.94	103.33	112.28
3	B	2	JZN	C11-C12-C13	10.62	118.87	111.44
3	A	1	JZN	C1-O1-C2	-8.75	104.67	117.51
3	A	1	JZN	C33-C32-N4	-8.27	100.11	112.36
3	B	2	JZN	C1-O1-C2	-7.01	107.23	117.51

There are no chirality outliers.

5 of 84 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1	JZN	O3-C28-N4-C29
3	A	1	JZN	C26-C28-N4-C29
3	A	1	JZN	O3-C28-N4-C32
3	A	1	JZN	C26-C28-N4-C32
3	A	1	JZN	C2-C7-C8-C10


There are no ring outliers.

3 monomers are involved in 32 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	3	JZN	9	0
3	A	1	JZN	14	0
3	B	2	JZN	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/259 (94%)	-0.01	2 (0%) 82 64	36, 51, 69, 74	0
1	B	247/259 (95%)	-0.07	3 (1%) 76 55	40, 53, 74, 82	1 (0%)
1	C	246/259 (94%)	0.53	11 (4%) 38 20	57, 76, 94, 103	0
2	D	9/12 (75%)	0.26	0 100 100	74, 75, 77, 77	0
2	E	10/12 (83%)	0.03	0 100 100	60, 61, 62, 64	0
2	F	10/12 (83%)	0.93	2 (20%) 3 2	72, 73, 75, 75	0
All	All	767/813 (94%)	0.16	18 (2%) 61 38	36, 61, 88, 103	1 (0%)

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	636	LEU	3.3
1	C	681	LYS	3.3
1	A	706	GLY	2.9
1	C	761	ILE	2.9
1	C	645	HIS	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	JZN	C	3	47/47	0.85	0.16	74,77,79,79	0
3	JZN	A	1	47/47	0.88	0.13	56,59,60,63	0
3	JZN	B	2	47/47	0.90	0.11	49,55,61,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around JZN A 1:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around JZN B 2:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.