



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2026 – 06:28 AM UTC

PDB ID : 7K2K / pdb_00007k2k
Title : Kelch domain of human KEAP1 bound to Nrf2 cyclic peptide, c[BAL-DEETGE]
Authors : Muellers, S.N.; Allen, K.N.
Deposited on : 2020-09-08
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	5% 82% 13% ••
1	B	301	22% 80% 14% 6%
2	P	7	14% 71% 29%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4687 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kelch-like ECH-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	288	Total	C	N	O	S	0	0	0
			2205	1372	397	420	16			
1	B	283	Total	C	N	O	S	0	0	0
			2167	1350	392	410	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	540	ALA	GLU	conflict	UNP Q14145
A	542	ALA	GLU	conflict	UNP Q14145
A	613	SER	CYS	conflict	UNP Q14145
B	540	ALA	GLU	conflict	UNP Q14145
B	542	ALA	GLU	conflict	UNP Q14145
B	613	SER	CYS	conflict	UNP Q14145

- Molecule 2 is a protein called BAL-ASP-GLU-GLU-THR-GLY-GLU.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	P	7	Total	C	N	O	0	0	0
			51	28	7	16			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	191	Total	O	0	0
			191	191		
3	B	70	Total	O	0	0
			70	70		
3	P	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Kelch-like ECH-associated protein 1



- Molecule 1: Kelch-like ECH-associated protein 1



- Molecule 2: BAL-ASP-GLU-GLU-THR-GLY-GLU



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	162.55Å 68.87Å 77.47Å 90.00° 117.61° 90.00°	Depositor
Resolution (Å)	29.37 – 1.98 29.37 – 1.98	Depositor EDS
% Data completeness (in resolution range)	94.6 (29.37-1.98) 94.5 (29.37-1.98)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.38 (at 1.98Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.197 , 0.220 0.198 , 0.222	Depositor DCC
R_{free} test set	2006 reflections (3.80%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.097	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4687	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/2259	0.90	6/3078 (0.2%)
1	B	0.60	0/2220	0.75	0/3025
2	P	0.68	0/45	0.83	0/59
All	All	0.69	0/4524	0.83	6/6162 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	613	SER	CA-C-N	-6.69	109.67	121.70
1	A	613	SER	C-N-CA	-6.69	109.67	121.70
1	A	348	SER	CA-C-N	5.86	129.22	120.31
1	A	348	SER	C-N-CA	5.86	129.22	120.31
1	A	349	ASP	CB-CA-C	-5.39	100.32	110.67
1	A	550	MET	CG-SD-CE	-5.33	89.18	100.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2205	0	2095	36	1
1	B	2167	0	2062	36	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	P	51	0	36	2	0
3	A	191	0	0	2	0
3	B	70	0	0	4	0
3	P	3	0	0	0	0
All	All	4687	0	4193	73	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:494:ARG:NH1	1:B:498:ARG:NH1	1.64	1.42
1:B:516:HIS:O	3:B:701:HOH:O	1.77	1.01
1:A:518:CYS:SG	1:A:537:TYR:O	2.20	0.98
1:A:494:ARG:HG2	1:A:496:GLU:CD	2.01	0.86
1:A:329:TYR:CE1	1:A:609:THR:HG22	2.11	0.85
1:A:515:LEU:HD22	1:A:566:ILE:HG13	1.59	0.83
1:B:434:CYS:SG	3:B:770:HOH:O	2.32	0.77
1:A:550:MET:HE1	1:A:568:VAL:HG11	1.67	0.76
1:B:515:LEU:HB3	1:B:520:TYR:CE1	2.22	0.74
1:A:338:SER:O	1:A:361:PRO:HB3	1.91	0.71
1:B:515:LEU:HD22	1:B:566:ILE:HG13	1.71	0.70
1:A:329:TYR:CE1	1:A:609:THR:CG2	2.79	0.66
1:A:518:CYS:SG	1:A:538:ASP:HA	2.36	0.66
1:B:466:ALA:HB1	1:B:514:VAL:HG23	1.77	0.65
1:A:518:CYS:HG	1:A:537:TYR:C	2.02	0.63
1:B:550:MET:HE2	1:B:568:VAL:HG21	1.82	0.60
1:A:518:CYS:SG	1:A:537:TYR:C	2.85	0.59
1:A:550:MET:HE1	1:A:568:VAL:CG1	2.33	0.58
1:B:515:LEU:HD22	1:B:566:ILE:CG1	2.33	0.58
1:B:557:LEU:H	1:B:557:LEU:HD23	1.68	0.58
1:B:466:ALA:HB1	1:B:514:VAL:CG2	2.34	0.57
1:A:494:ARG:HG3	1:A:494:ARG:O	2.04	0.57
1:B:522:ALA:HB1	1:B:550:MET:HE3	1.87	0.56
1:B:415:ARG:NH2	2:P:79:GLU:OE2	2.33	0.55
1:B:494:ARG:NH1	1:B:498:ARG:CZ	2.62	0.55
1:B:380:ARG:HD2	3:B:714:HOH:O	2.05	0.55
1:A:515:LEU:HB3	1:A:520:TYR:CE1	2.42	0.55
1:B:515:LEU:HD22	1:B:566:ILE:CD1	2.36	0.55
1:B:550:MET:CE	1:B:568:VAL:HG11	2.36	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:557:LEU:HD23	1:A:557:LEU:H	1.72	0.54
1:A:518:CYS:HG	1:A:538:ASP:HA	1.75	0.52
1:A:548:ALA:O	1:A:591:TRP:NE1	2.40	0.51
1:A:338:SER:HB3	1:A:361:PRO:C	2.36	0.50
1:B:515:LEU:HB3	1:B:520:TYR:HE1	1.74	0.50
1:A:346:ASN:CG	1:A:349:ASP:HB2	2.36	0.50
1:A:329:TYR:CD1	1:A:609:THR:HG22	2.46	0.50
1:A:613:SER:O	1:A:614:ARG:C	2.55	0.49
1:B:470:ARG:NH2	3:B:704:HOH:O	2.44	0.49
1:B:550:MET:HE1	1:B:568:VAL:HG11	1.94	0.48
1:A:338:SER:O	1:A:361:PRO:CB	2.59	0.48
1:B:373:LEU:HB3	1:B:395:CYS:SG	2.53	0.48
1:B:361:PRO:C	1:B:362:ARG:HG3	2.38	0.47
1:A:566:ILE:HB	1:A:584:TYR:HB3	1.97	0.47
1:B:483:ARG:HG2	1:B:508:SER:HB2	1.95	0.47
1:B:455:PRO:O	1:B:497:TRP:CD1	2.69	0.46
1:B:400:THR:HB	1:B:402:GLN:OE1	2.16	0.46
1:B:345:TYR:HB2	1:B:352:TRP:CH2	2.51	0.46
1:B:562:HIS:HB3	1:B:567:TYR:CE2	2.51	0.46
1:B:365:LEU:H	1:B:365:LEU:HD23	1.82	0.45
1:B:515:LEU:HD22	1:B:566:ILE:HD11	1.99	0.45
1:A:472:LEU:HB3	1:A:490:TYR:HB3	1.99	0.45
2:P:78:GLU:H	2:P:78:GLU:HG3	1.65	0.45
1:A:380:ARG:HD3	1:A:382:ASN:OD1	2.17	0.44
1:B:347:PRO:HG2	1:B:562:HIS:CD2	2.52	0.44
1:A:399:MET:HE2	3:A:750:HOH:O	2.18	0.44
1:A:518:CYS:SG	1:A:538:ASP:CA	3.04	0.43
1:A:446:GLU:H	1:A:446:GLU:CD	2.26	0.43
1:B:393:LEU:HD23	1:B:406:CYS:HB2	2.01	0.43
1:B:519:ILE:O	1:B:536:ARG:HA	2.17	0.43
1:B:565:ARG:HD3	1:B:585:ASP:HA	2.00	0.42
1:A:494:ARG:HG2	1:A:496:GLU:OE2	2.19	0.42
1:A:335:PHE:HD2	3:A:859:HOH:O	2.01	0.42
1:B:335:PHE:O	1:B:336:ARG:HB2	2.19	0.42
1:B:566:ILE:O	1:B:583:CYS:HA	2.19	0.42
1:A:519:ILE:O	1:A:536:ARG:HA	2.19	0.42
1:A:388:THR:HG22	1:A:389:ASP:O	2.19	0.41
1:B:345:TYR:HB2	1:B:352:TRP:CZ3	2.54	0.41
1:A:338:SER:OG	1:A:381:ASN:HA	2.20	0.41
1:A:512:VAL:HA	1:A:520:TYR:O	2.20	0.41
1:B:356:ALA:HB2	1:B:401:ASN:OD1	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:381:ASN:O	1:A:387:ASN:HA	2.20	0.41
1:A:554:ARG:HD2	1:A:568:VAL:HG11	2.03	0.41
1:A:515:LEU:HD22	1:A:566:ILE:CG1	2.39	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:434:CYS:SG	1:B:336:ARG:NH2[2_556]	1.58	0.62

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/301 (95%)	275 (96%)	10 (4%)	1 (0%)	36	27
1	B	281/301 (93%)	270 (96%)	10 (4%)	1 (0%)	30	20
2	P	5/7 (71%)	5 (100%)	0	0	100	100
All	All	572/609 (94%)	550 (96%)	20 (4%)	2 (0%)	36	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	516	HIS
1	B	516	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/245 (95%)	229 (99%)	3 (1%)	61	57
1	B	227/245 (93%)	227 (100%)	0	100	100
2	P	5/5 (100%)	5 (100%)	0	100	100
All	All	464/495 (94%)	461 (99%)	3 (1%)	78	76

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	517	ASN
1	A	518	CYS
1	A	610	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	432	HIS
1	A	516	HIS
1	A	552	HIS
1	B	359	GLN
1	B	414	ASN
1	B	517	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BAL	P	76	2	3,4,5	0.22	0	3,3,5	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BAL	P	76	2	-	1/1/2/3	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	P	76	BAL	C-CA-CB-N

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	288/301 (95%)	-0.10	16 (5%) 30 38	14, 22, 43, 65	0
1	B	283/301 (94%)	1.01	67 (23%) 2 2	16, 45, 88, 107	0
2	P	6/7 (85%)	0.51	1 (16%) 4 5	23, 28, 58, 61	0
All	All	577/609 (94%)	0.45	84 (14%) 6 8	14, 30, 82, 107	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	516	HIS	7.6
1	A	335	PHE	7.1
1	A	338	SER	6.2
1	B	349	ASP	5.5
1	A	518	CYS	5.4
1	B	515	LEU	5.3
1	B	517	ASN	5.2
1	A	337	GLN	5.1
1	B	327	LEU	5.1
1	B	353	LEU	4.9
1	B	329	TYR	4.8
1	A	517	ASN	4.7
1	A	610	MET	4.6
1	A	336	ARG	4.6
1	B	373	LEU	4.5
1	B	351	THR	4.3
1	B	370	VAL	4.1
1	A	613	SER	4.1
1	B	608	VAL	4.0
1	A	516	HIS	3.7
1	B	609	THR	3.5
1	B	423	GLY	3.3
1	B	350	GLY	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	345	TYR	3.2
1	A	612	PRO	3.2
1	A	614	ARG	3.2
1	B	347	PRO	3.2
1	B	562	HIS	3.1
1	B	594	VAL	3.1
1	B	399	MET	3.1
1	B	400	THR	3.0
1	B	445	PRO	2.9
1	B	446	GLU	2.9
1	B	575	HIS	2.8
1	B	470	ARG	2.8
1	B	355	LEU	2.8
1	B	595	THR	2.8
2	P	77	ASP	2.8
1	B	567	TYR	2.7
1	A	494	ARG	2.7
1	B	328	ILE	2.7
1	B	369	VAL	2.7
1	B	348	SER	2.7
1	B	352	TRP	2.7
1	B	563	GLN	2.6
1	B	592	SER	2.6
1	B	452	LEU	2.6
1	B	420	VAL	2.6
1	B	590	THR	2.6
1	B	371	GLY	2.5
1	B	514	VAL	2.5
1	B	398	PRO	2.5
1	B	422	ASP	2.4
1	B	576	THR	2.4
1	B	336	ARG	2.4
1	B	493	GLU	2.4
1	B	397	ASN	2.4
1	B	421	ILE	2.4
1	B	561	VAL	2.4
1	B	607	ALA	2.3
1	B	356	ALA	2.3
1	B	596	ARG	2.3
1	B	335	PHE	2.3
1	B	447	ARG	2.3
1	B	585	ASP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	360	VAL	2.3
1	B	395	CYS	2.3
1	B	396	TYR	2.2
1	B	405	PRO	2.2
1	B	451	HIS	2.2
1	B	494	ARG	2.2
1	A	349	ASP	2.2
1	B	346	ASN	2.2
1	A	548	ALA	2.2
1	B	407	ALA	2.2
1	B	424	HIS	2.1
1	B	434	CYS	2.1
1	A	546	PHE	2.1
1	B	443	TYR	2.1
1	B	591	TRP	2.1
1	B	584	TYR	2.0
1	B	372	GLY	2.0
1	B	583	CYS	2.0
1	A	385	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BAL	P	76	5/6	0.91	0.12	42,50,55,57	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.