



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:34 AM UTC

PDB ID : 1K3X / pdb_00001k3x
Title : Crystal structure of a trapped reaction intermediate of the DNA repair enzyme
Endonuclease VIII with Brominated-DNA
Authors : Golan, G.; Zharkov, D.O.; Gilboa, R.; Fernandes, A.S.; Kycia, J.H.; Gerch-
man, S.E.; Rieger, R.A.; Grollman, A.P.; Shoham, G.
Deposited on : 2001-10-04
Resolution : 1.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1730 (1.28-1.24)
Ramachandran outliers	187476	1695 (1.28-1.24)
Sidechain outliers	187428	1694 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	B	13	8% 31% 38% 23%
2	C	13	31% 31% 15% 23%
3	A	262	85% 10% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	BRU	B	404	X	-	-	-
6	GOL	A	511	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 2998 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*GP*CP*(BRU)P*(BRU)P*CP*AP*(BRU)P*CP*CP*(BRU)P*GP*G)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	10	Total	Br	C	N	O	P	0	0	0
			199	4	92	30	63	10			

- Molecule 2 is a DNA chain called 5'-D(*CP*CP*AP*GP*GP*AP*(PED)P*GP*AP*AP*GP*CP*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	P	0	0	0
			202	94	43	55	10			

- Molecule 3 is a protein called Endonuclease VIII.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	253	Total	C	N	O	S	0	8	0
			2059	1316	370	368	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	THR	PRO	SEE REMARK 999	UNP P50465
A	112	ARG	THR	SEE REMARK 999	UNP P50465

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	24	Total	O	0	0
			24	24		
7	C	23	Total	O	0	0
			23	23		
7	A	441	Total	O	0	0
			441	441		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: 5'-D(*GP*GP*CP*(BRU)P*(BRU)P*CP*AP*(BRU)P*CP*CP*(BRU)P*GP*G)-3'

Chain B: 




- Molecule 2: 5'-D(*CP*CP*AP*GP*GP*AP*(PED)P*GP*AP*AP*GP*CP*C)-3'

Chain C: 



- Molecule 3: Endonuclease VIII

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	76.50 Å 76.50 Å 164.31 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.25	Depositor
% Data completeness (in resolution range)	97.4 (10.00-1.25)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.149 , 0.181	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2998	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BRU, PED, GOL, SO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.90	0/129	2.47	11/188 (5.9%)
2	C	1.00	0/215	2.76	22/328 (6.7%)
3	A	0.91	0/2136	1.32	14/2898 (0.5%)
All	All	0.92	0/2480	1.60	47/3414 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	0

There are no bond length outliers.

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	75	ARG	CD-NE-CZ	12.69	142.16	124.40
2	C	429	DA	C4'-C3'-O3'	-11.18	93.22	110.00
2	C	430	DA	C6-N1-C2	9.66	133.30	118.80
2	C	430	DA	P-O5'-C5'	-9.34	105.99	120.00
2	C	429	DA	C2'-C3'-O3'	-8.89	98.16	111.50
2	C	424	DG	C8-N9-C1'	8.43	139.65	127.00
2	C	429	DA	C1'-O4'-C4'	-7.41	98.58	109.70
1	B	412	DG	N3-C4-C5	-7.38	117.33	128.40
3	A	12	ASN	OD1-CG-ND2	7.22	129.82	122.60
3	A	184	HIS	CA-CB-CG	-7.20	106.60	113.80
3	A	239	ARG	CD-NE-CZ	7.18	134.45	124.40
2	C	431	DG	N7-C8-N9	7.06	124.09	113.50
2	C	429	DA	C6-N1-C2	7.05	129.38	118.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	430	DA	O3'-P-O5'	-7.04	93.44	104.00
2	C	422	DC	C6-N1-C1'	7.02	130.23	119.70
1	B	403	DC	C2-N3-C4	6.55	129.82	120.00
2	C	424	DG	N7-C8-N9	6.47	123.21	113.50
1	B	412	DG	N7-C8-N9	6.46	123.20	113.50
2	C	431	DG	N3-C4-C5	-6.38	118.83	128.40
2	C	429	DA	N1-C2-N3	-6.36	119.47	129.00
2	C	422	DC	C2-N3-C4	6.28	129.42	120.00
3	A	97	ASP	CA-CB-CG	6.27	118.87	112.60
1	B	407	DA	N1-C6-N6	-6.01	109.98	119.00
1	B	412	DG	C6-N1-C2	-5.92	116.02	124.90
2	C	425	DG	C8-N9-C1'	5.91	135.87	127.00
1	B	412	DG	C2-N3-C4	5.87	120.61	111.80
2	C	430	DA	N1-C2-N3	-5.87	120.19	129.00
1	B	407	DA	C5-C6-N6	5.87	132.21	123.40
2	C	430	DA	N1-C6-N6	5.79	127.68	119.00
2	C	422	DC	O5'-C5'-C4'	5.71	119.36	110.80
3	A	82	GLU	O-C-N	5.69	127.86	121.32
3	A	84	GLN	CA-C-O	5.64	127.36	121.15
2	C	430	DA	C5-C6-N1	-5.56	109.26	117.60
3	A	114	GLU	CA-CB-CG	5.54	125.18	114.10
1	B	412	DG	N3-C4-N9	5.41	134.12	126.00
1	B	409	DC	O4'-C1'-N1	-5.41	100.29	108.40
3	A	193	GLN	CG-CD-NE2	-5.36	108.36	116.40
2	C	431	DG	C2-N3-C4	5.28	119.72	111.80
3	A	114	GLU	CB-CG-CD	5.27	121.55	112.60
1	B	409	DC	N3-C2-O2	5.26	129.80	121.90
2	C	431	DG	C8-N9-C4	-5.12	98.32	106.00
2	C	422	DC	N1-C1'-C2'	5.12	121.17	113.50
3	A	79	THR	CA-CB-OG1	-5.05	102.03	109.60
3	A	46	HIS	CB-CG-CD2	-5.04	124.64	131.20
1	B	409	DC	C2-N3-C4	5.03	127.54	120.00
3	A	128	ASP	CA-CB-CG	5.02	117.62	112.60
3	A	262	HIS	ND1-CE1-NE2	-5.01	103.39	108.40

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	404	BRU	C4'

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	199	0	103	9	0
2	C	202	0	108	5	0
3	A	2059	0	2057	19	0
4	A	1	0	0	0	0
5	A	25	0	0	0	0
6	A	24	0	32	8	0
7	A	441	0	0	8	0
7	B	24	0	0	1	0
7	C	23	0	0	1	0
All	All	2998	0	2300	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:147:ARG:HH21	6:A:511:GOL:H11	1.26	1.00
1:B:410:DC:H2''	1:B:411:BRU:BR	2.35	0.80
2:C:422:DC:H2'	2:C:422:DC:OP2	1.82	0.79
3:A:69:GLN:HG3	7:A:789:HOH:O	1.85	0.76
3:A:150:ASN:HB2	7:A:894:HOH:O	1.87	0.72
6:A:513:GOL:H12	7:A:737:HOH:O	1.91	0.70
6:A:511:GOL:H12	7:A:773:HOH:O	1.90	0.69
7:C:946:HOH:O	3:A:124[A]:ARG:HD3	1.95	0.67
3:A:125[A]:VAL:HG22	3:A:164:ALA:HB2	1.83	0.61
3:A:147:ARG:NH2	6:A:511:GOL:H11	2.08	0.60
1:B:404:BRU:H2'	1:B:405:BRU:BR	2.57	0.59
1:B:411:BRU:H2''	1:B:412:DG:C8	2.40	0.57
3:A:245:GLU:HG2	7:A:892:HOH:O	2.04	0.57
7:B:942:HOH:O	3:A:86:THR:HG23	2.05	0.55
1:B:410:DC:C2'	1:B:411:BRU:BR	3.10	0.52
1:B:410:DC:H2''	1:B:411:BRU:OP2	2.11	0.50
3:A:147:ARG:HH21	6:A:511:GOL:C1	2.13	0.50
1:B:411:BRU:H2''	1:B:412:DG:N7	2.27	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:430:DA:OP2	3:A:124[A]:ARG:NE	2.47	0.47
3:A:172[B]:VAL:CG1	3:A:229:VAL:HG23	2.45	0.47
3:A:117:THR:HB	6:A:514:GOL:H31	1.97	0.47
3:A:117:THR:CB	6:A:514:GOL:H31	2.45	0.46
3:A:124[A]:ARG:NH2	7:A:644:HOH:O	2.50	0.45
3:A:1:PRO:HG3	7:A:961:HOH:O	2.16	0.44
3:A:60:ASN:O	3:A:61:ASP:HB2	2.18	0.43
3:A:83:PRO:HD2	7:A:779:HOH:O	2.17	0.43
3:A:195:ASP:OD2	6:A:512:GOL:O2	2.30	0.41
2:C:430:DA:OP2	3:A:124[A]:ARG:NH2	2.49	0.41
1:B:404:BRU:H2''	1:B:405:BRU:O5'	2.21	0.41
1:B:412:DG:N9	1:B:412:DG:H5'	2.36	0.41
2:C:422:DC:H2'	2:C:422:DC:P	2.61	0.41
1:B:403:DC:N4	2:C:430:DA:N6	2.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	257/262 (98%)	247 (96%)	10 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	224/226 (99%)	220 (98%)	4 (2%)	51 15

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	1	PRO
3	A	69	GLN
3	A	124[A]	ARG
3	A	124[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	184	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	BRU	B	404	2,1	18,21,22	1.29	3 (16%)	25,30,33	4.55	12 (48%)
1	BRU	B	408	2,1	18,21,22	1.20	2 (11%)	25,30,33	1.65	6 (24%)
1	BRU	B	405	2,1	18,21,22	0.95	1 (5%)	25,30,33	2.31	9 (36%)
1	BRU	B	411	2,1	18,21,22	1.51	1 (5%)	25,30,33	2.72	14 (56%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	BRU	B	404	2,1	1/1/4/4	2/7/21/22	0/2/2/2
1	BRU	B	408	2,1	-	0/7/21/22	0/2/2/2
1	BRU	B	405	2,1	-	0/7/21/22	0/2/2/2
1	BRU	B	411	2,1	-	0/7/21/22	0/2/2/2

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	411	BRU	C2-N3	5.00	1.46	1.38
1	B	404	BRU	C2-N3	2.88	1.43	1.38
1	B	408	BRU	C2-N3	2.65	1.42	1.38
1	B	404	BRU	C4-N3	-2.40	1.34	1.38
1	B	404	BRU	O3'-C3'	2.13	1.47	1.43
1	B	408	BRU	O4'-C1'	2.10	1.47	1.42
1	B	405	BRU	O4'-C1'	2.06	1.47	1.42

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	404	BRU	BR-C5-C4	-14.31	101.55	118.02
1	B	404	BRU	BR-C5-C6	11.30	136.45	120.64
1	B	404	BRU	O2-C2-N1	6.96	131.85	122.80
1	B	411	BRU	C1'-N1-C6	-6.65	109.57	120.74
1	B	405	BRU	C6-C5-C4	6.43	127.20	120.67
1	B	411	BRU	C1'-N1-C2	5.69	128.78	117.66
1	B	404	BRU	O4-C4-C5	-5.59	118.67	125.80
1	B	411	BRU	C4-N3-C2	-5.46	120.19	127.34
1	B	405	BRU	O4'-C1'-N1	-5.15	98.73	107.86
1	B	404	BRU	O2-C2-N3	-4.20	113.75	121.49
1	B	405	BRU	C6-N1-C2	4.06	125.34	121.30
1	B	404	BRU	O4'-C1'-N1	4.05	115.05	107.86
1	B	408	BRU	O4'-C1'-N1	-3.53	101.60	107.86
1	B	408	BRU	C4-N3-C2	-3.50	122.76	127.34
1	B	411	BRU	O2-C2-N3	-3.45	115.13	121.49
1	B	404	BRU	O3'-C3'-C2'	-3.22	99.70	110.88
1	B	411	BRU	N3-C2-N1	2.97	118.76	114.89
1	B	404	BRU	O4-C4-N3	2.97	125.69	120.11
1	B	404	BRU	O4'-C4'-C3'	-2.96	98.90	105.65
1	B	408	BRU	N3-C2-N1	2.83	118.58	114.89
1	B	411	BRU	BR-C5-C6	-2.83	116.69	120.64
1	B	408	BRU	C5-C4-N3	2.72	116.48	113.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	404	BRU	C6-N1-C2	2.71	124.01	121.30
1	B	404	BRU	C1'-N1-C2	-2.67	112.43	117.66
1	B	405	BRU	C5-C6-N1	-2.65	114.72	120.91
1	B	405	BRU	BR-C5-C4	-2.64	114.98	118.02
1	B	404	BRU	O4'-C4'-C5'	2.62	117.71	109.33
1	B	411	BRU	O4'-C4'-C5'	2.57	117.58	109.33
1	B	405	BRU	C4-N3-C2	-2.55	124.00	127.34
1	B	411	BRU	O2-C2-N1	2.53	126.09	122.80
1	B	411	BRU	O3'-C3'-C4'	-2.44	100.83	110.07
1	B	411	BRU	C5-C4-N3	2.39	116.09	113.34
1	B	408	BRU	C2'-C1'-N1	2.29	119.54	113.81
1	B	411	BRU	C3'-C2'-C1'	-2.25	97.10	102.60
1	B	411	BRU	BR-C5-C4	2.15	120.50	118.02
1	B	405	BRU	C1'-N1-C6	-2.14	117.15	120.74
1	B	411	BRU	C6-C5-C4	2.10	122.81	120.67
1	B	408	BRU	O3'-C3'-C4'	-2.07	102.24	110.07
1	B	411	BRU	O3'-C3'-C2'	-2.06	103.73	110.88
1	B	405	BRU	O3'-C3'-C4'	-2.03	102.39	110.07
1	B	405	BRU	BR-C5-C6	-2.02	117.81	120.64

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	404	BRU	C4'

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	404	BRU	C4'-C5'-O5'-P
1	B	404	BRU	C3'-C4'-C5'-O5'

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	404	BRU	2	0
1	B	405	BRU	2	0
1	B	411	BRU	5	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SO4	A	554	-	4,4,4	0.27	0	6,6,6	0.15	0
6	GOL	A	514	-	5,5,5	0.66	0	5,5,5	0.40	0
5	SO4	A	552	-	4,4,4	0.56	0	6,6,6	0.52	0
6	GOL	A	512	-	5,5,5	0.55	0	5,5,5	0.91	0
5	SO4	A	551	-	4,4,4	0.19	0	6,6,6	0.70	0
6	GOL	A	513	-	5,5,5	1.09	1 (20%)	5,5,5	1.43	1 (20%)
6	GOL	A	511	-	5,5,5	1.02	0	5,5,5	2.43	2 (40%)
5	SO4	A	553	-	4,4,4	0.27	0	6,6,6	0.31	0
5	SO4	A	555	-	4,4,4	0.33	0	6,6,6	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	512	-	-	0/4/4/4	-
6	GOL	A	514	-	-	4/4/4/4	-
6	GOL	A	511	-	-	3/4/4/4	-
6	GOL	A	513	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	513	GOL	C1-C2	2.23	1.60	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	511	GOL	O3-C3-C2	4.43	130.34	110.38
6	A	511	GOL	C3-C2-C1	-3.00	100.79	111.80
6	A	513	GOL	O3-C3-C2	2.39	121.16	110.38

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	511	GOL	O1-C1-C2-C3
6	A	513	GOL	O1-C1-C2-C3
6	A	514	GOL	O1-C1-C2-C3
6	A	513	GOL	O1-C1-C2-O2
6	A	514	GOL	C1-C2-C3-O3
6	A	511	GOL	O1-C1-C2-O2
6	A	514	GOL	O1-C1-C2-O2
6	A	514	GOL	O2-C2-C3-O3
6	A	511	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	514	GOL	2	0
6	A	512	GOL	1	0
6	A	513	GOL	1	0
6	A	511	GOL	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.