



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 09:45 PM UTC

PDB ID : 3K3L / pdb_00003k3l
Title : Crystal structure of Siderocalin (NGAL, Lipocalin 2) complexed with apo Enterobactin
Authors : Clifton, M.C.
Deposited on : 2009-10-02
Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	4% 89% 8% ..
1	B	178	79% 76% 10% 13%
1	C	178	4% 86% 11% ..

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 3940 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Neutrophil gelatinase-associated lipocalin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	175	Total	C	N	O	S	0	1	0
			1384	904	226	250	4			
1	B	154	Total	C	N	O	S	0	0	0
			903	566	159	175	3			
1	C	175	Total	C	N	O	S	0	2	0
			1409	914	234	257	4			

There are 3 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	87	SER	CYS	engineered mutation	UNP P80188
B	87	SER	CYS	engineered mutation	UNP P80188
C	87	SER	CYS	engineered mutation	UNP P80188

- Molecule 2 is 2-(2,3-DIHYDROXY-BENZOYLAMINO)-3-HYDROXY-PROPIONIC ACID (CCD ID: DBS) (formula: C₁₀H₁₁NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			17	10	1	6		

- Molecule 3 is 2,3-DIHYDROXY-BENZOIC ACID (CCD ID: DBH) (formula: $C_7H_6O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	7	4		
3	A	1	Total	C	O	0	0
			11	7	4		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Na 1 1	0	0
5	C	1	Total Na 1 1	0	0

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O S 5 4 1	0	0

- Molecule 7 is N-[(2,3-dihydroxyphenyl)carbonyl]-O-[(2S)-2-[(2,3-dihydroxyphenyl)carbonyl]amino]-3-(L-seryloxy)propanoyl]-L-serine (CCD ID: MCK) (formula: $C_{23}H_{25}N_3O_{13}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	C	1	Total	C	N	O	0	0
			39	23	3	13		

- Molecule 8 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	C	1	Total	Cl	0	0
			1	1		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	41	Total	O	0	0
			41	41		
9	B	14	Total	O	0	0
			14	14		
9	C	61	Total	O	0	0
			61	61		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Neutrophil gelatinase-associated lipocalin



- Molecule 1: Neutrophil gelatinase-associated lipocalin



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	114.20Å 114.20Å 119.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.25 – 2.62 41.25 – 2.62	Depositor EDS
% Data completeness (in resolution range)	99.9 (41.25-2.62) 98.6 (41.25-2.62)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.99 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.260 , 0.307 0.261 , 0.303	Depositor DCC
R_{free} test set	2245 reflections (9.39%)	wwPDB-VP
Wilson B-factor (Å ²)	45.4	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for -h,l,k 0.009 for -l,-k,-h	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	3940	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL, NA, DBH, DBS, CL, MCK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/1424	0.63	0/1934
1	B	0.40	0/915	0.76	2/1257 (0.2%)
1	C	0.38	0/1452	0.64	0/1972
All	All	0.40	0/3791	0.67	2/5163 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	161	LEU	CA-C-N	5.73	126.52	120.11
1	B	161	LEU	C-N-CA	5.73	126.52	120.11

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1384	0	1347	12	0
1	B	903	0	572	8	0
1	C	1409	0	1381	12	0
2	A	17	0	8	0	0
3	A	22	0	6	0	0
4	A	12	0	16	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	30	0	40	1	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
6	A	5	0	0	0	0
7	C	39	0	21	0	0
8	C	1	0	0	0	0
9	A	41	0	0	0	0
9	B	14	0	0	0	0
9	C	61	0	0	0	0
All	All	3940	0	3391	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34:VAL:HG23	1:A:137:LEU:HD13	1.65	0.78
1:A:34:VAL:HG21	1:A:144:LEU:HD12	1.72	0.72
1:B:162:PRO:HB2	1:B:163:GLU:HG2	1.73	0.71
1:C:34:VAL:HG21	1:C:144:LEU:HD12	1.72	0.71
1:A:33:VAL:HG21	1:A:52:TYR:CE2	2.29	0.68
1:B:27:PHE:HA	1:B:28:GLN:CB	2.23	0.68
1:C:36:LEU:HD13	1:C:41:ILE:HD12	1.77	0.66
1:C:66:VAL:HG21	1:C:83:PHE:CD1	2.32	0.64
1:A:34:VAL:CG2	1:A:137:LEU:HD13	2.29	0.61
1:B:119:ALA:HB3	1:B:138:TYR:CB	2.33	0.58
1:C:70:LEU:HD12	1:C:70:LEU:N	2.24	0.53
1:C:114:ASN:HD21	1:C:117:GLN:HB2	1.74	0.53
1:B:27:PHE:CA	1:B:28:GLN:CB	2.88	0.51
1:B:61:ASP:O	1:B:62:LYS:CB	2.59	0.51
1:A:119:ALA:HB3	1:A:138:TYR:HB2	1.93	0.51
1:A:140:ARG:HG2	1:C:23:GLN:NE2	2.26	0.50
1:A:34:VAL:HG21	1:A:144:LEU:CD1	2.43	0.48
1:C:69:VAL:HG22	1:C:78:TYR:CD1	2.48	0.48
1:B:162:PRO:CB	1:B:163:GLU:HG2	2.43	0.47
1:A:31:TRP:CE3	1:A:138:TYR:HB3	2.49	0.47
1:B:31:TRP:O	1:B:53:ALA:HB1	2.14	0.47
1:C:31:TRP:CE3	1:C:138:TYR:HB3	2.50	0.46
1:C:34:VAL:CG2	1:C:144:LEU:HD12	2.43	0.46
1:C:33:VAL:HG21	1:C:52:TYR:CZ	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:VAL:HG21	1:A:83:PHE:CD1	2.52	0.45
1:A:31:TRP:CZ3	1:A:140:ARG:HD2	2.52	0.44
1:A:33:VAL:HG21	1:A:52:TYR:CZ	2.52	0.43
1:A:114:ASN:HD21	1:A:117:GLN:HB2	1.83	0.43
1:B:162:PRO:C	1:B:163:GLU:HG2	2.45	0.41
1:C:33:VAL:HG21	1:C:52:TYR:CE2	2.56	0.41
1:C:97:ILE:HD12	4:C:3731:GOL:H11	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	174/178 (98%)	167 (96%)	7 (4%)	0	100	100
1	B	142/178 (80%)	121 (85%)	16 (11%)	5 (4%)	3	4
1	C	175/178 (98%)	172 (98%)	3 (2%)	0	100	100
All	All	491/534 (92%)	460 (94%)	26 (5%)	5 (1%)	12	26

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	28	GLN
1	B	149	LYS
1	B	141	THR
1	B	52	TYR
1	B	95	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/163 (90%)	142 (97%)	5 (3%)	32	58
1	B	40/163 (24%)	37 (92%)	3 (8%)	12	26
1	C	154/163 (94%)	147 (96%)	7 (4%)	24	48
All	All	341/489 (70%)	326 (96%)	15 (4%)	25	48

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	ASP
1	A	109	ARG
1	A	137	LEU
1	A	140	ARG
1	A	154	ARG
1	B	42	LEU
1	B	87	SER
1	B	153	ILE
1	C	25	ASN
1	C	36	LEU
1	C	58	LEU
1	C	70	LEU
1	C	137	LEU
1	C	140	ARG
1	C	147	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	19	GLN
1	C	49	GLN
1	C	151	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 3 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	A	182	-	4,4,4	0.24	0	6,6,6	0.07	0
4	GOL	C	182	-	5,5,5	0.38	0	5,5,5	0.20	0
3	DBH	A	179	-	11,11,11	0.76	0	15,15,15	0.68	0
4	GOL	C	3731	-	5,5,5	0.39	0	5,5,5	0.24	0
2	DBS	A	301	-	17,17,17	0.64	0	23,23,23	0.71	0
7	MCK	C	179	-	39,40,40	1.67	3 (7%)	53,55,55	1.26	7 (13%)
4	GOL	C	180	-	5,5,5	0.42	0	5,5,5	0.17	0
4	GOL	C	181	-	5,5,5	0.37	0	5,5,5	0.15	0
4	GOL	A	180	-	5,5,5	0.45	0	5,5,5	0.23	0
3	DBH	A	303	-	11,11,11	0.74	0	15,15,15	0.69	0
4	GOL	A	3731	-	5,5,5	0.39	0	5,5,5	0.24	0
4	GOL	C	183	-	5,5,5	0.40	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MCK	C	179	-	-	9/40/40/40	0/2/2/2
4	GOL	C	182	-	-	2/4/4/4	-
3	DBH	A	179	-	-	4/4/4/4	0/1/1/1
4	GOL	C	3731	-	-	2/4/4/4	-
2	DBS	A	301	-	-	4/14/14/14	0/1/1/1
4	GOL	C	180	-	-	3/4/4/4	-
4	GOL	C	181	-	-	0/4/4/4	-
4	GOL	A	180	-	-	2/4/4/4	-
3	DBH	A	303	-	-	4/4/4/4	0/1/1/1
4	GOL	A	3731	-	-	2/4/4/4	-
4	GOL	C	183	-	-	2/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	C	179	MCK	O17-C13	6.37	1.46	1.33
7	C	179	MCK	O15-C24	6.04	1.46	1.33
7	C	179	MCK	O23-C21	4.80	1.45	1.30

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	C	179	MCK	O15-C24-C26	4.33	121.69	111.58
7	C	179	MCK	O17-C13-C12	3.63	120.11	111.58
7	C	179	MCK	O15-C24-O25	-3.41	117.91	124.14
7	C	179	MCK	C18-O17-C13	2.51	122.83	116.78
7	C	179	MCK	O17-C13-O16	-2.48	119.61	124.14
7	C	179	MCK	O23-C21-O22	-2.30	118.85	124.08
7	C	179	MCK	O23-C21-C19	2.13	120.70	113.51

There are no chirality outliers.

All (34) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	DBS	N1-C22-C28-O13
4	A	180	GOL	O1-C1-C2-C3
4	C	3731	GOL	O1-C1-C2-C3
4	C	183	GOL	C1-C2-C3-O3
7	C	179	MCK	O25-C24-O15-C14
7	C	179	MCK	C26-C24-O15-C14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	C	179	MCK	O17-C18-C19-N20
7	C	179	MCK	O17-C18-C19-C21
3	A	179	DBH	C3-C18-C21-O9
4	A	3731	GOL	O1-C1-C2-C3
4	C	180	GOL	C1-C2-C3-O3
3	A	179	DBH	C3-C18-C21-O17
4	C	3731	GOL	O1-C1-C2-O2
4	C	180	GOL	O2-C2-C3-O3
4	C	183	GOL	O2-C2-C3-O3
3	A	303	DBH	C3-C18-C21-O9
4	A	3731	GOL	O1-C1-C2-O2
4	A	180	GOL	O1-C1-C2-O2
7	C	179	MCK	O30-C29-C31-C32
3	A	303	DBH	C3-C18-C21-O17
2	A	301	DBS	N1-C22-C25-O10
2	A	301	DBS	N1-C22-C25-O15
7	C	179	MCK	N20-C29-C31-C32
2	A	301	DBS	C25-C22-C28-O13
4	C	182	GOL	O1-C1-C2-O2
7	C	179	MCK	O15-C24-C26-N27
4	C	182	GOL	O1-C1-C2-C3
4	C	180	GOL	O1-C1-C2-O2
3	A	179	DBH	C15-C18-C21-O17
3	A	179	DBH	C15-C18-C21-O9
7	C	179	MCK	O25-C24-C26-N27
3	A	303	DBH	C15-C18-C21-O9
7	C	179	MCK	C14-C12-C13-O16
3	A	303	DBH	C15-C18-C21-O17


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	3731	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	175/178 (98%)	0.59	7 (4%) 42 37	24, 38, 46, 53	2 (1%)
1	B	154/178 (86%)	3.36	140 (90%) 0 0	54, 85, 91, 91	0
1	C	175/178 (98%)	0.22	7 (4%) 42 37	17, 29, 43, 52	2 (1%)
All	All	504/534 (94%)	1.31	154 (30%) 1 1	17, 39, 89, 91	4 (0%)

All (154) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	45	ASP	9.7
1	B	32	TYR	6.6
1	B	8	ILE	6.5
1	B	64	TYR	6.5
1	B	167	VAL	6.4
1	B	65	ASN	6.3
1	B	97	ILE	6.1
1	B	159	LEU	6.0
1	B	170	VAL	5.8
1	B	160	GLY	5.7
1	B	114	ASN	5.5
1	B	116	ASN	5.3
1	B	71	PHE	5.2
1	B	129	ASN	5.1
1	B	95	GLY	5.1
1	B	172	ILE	5.0
1	B	79	TRP	4.7
1	B	104	THR	4.6
1	B	112	SER	4.6
1	B	26	GLN	4.6
1	B	39	ASN	4.6
1	B	78	TYR	4.5
1	B	37	ALA	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	77	ASP	4.5
1	B	88	GLN	4.4
1	B	115	TYR	4.4
1	B	31	TRP	4.2
1	B	54	THR	4.2
1	B	73	LYS	4.2
1	B	140	ARG	4.2
1	B	92	PHE	4.2
1	B	98	LYS	4.1
1	B	132	TYR	4.1
1	B	178	GLY	4.1
1	B	93	THR	4.1
1	B	171	PRO	4.0
1	B	20	GLN	4.0
1	B	25	ASN	4.0
1	B	118	HIS	4.0
1	B	151	ASN	4.0
1	B	56	TYR	3.9
1	B	109	ARG	3.9
1	B	94	LEU	3.9
1	B	141	THR	3.9
1	B	142	LYS	3.8
1	B	22	PHE	3.8
1	B	68	SER	3.8
1	B	33	VAL	3.8
1	B	175	CYS	3.8
1	B	83	PHE	3.8
1	B	121	VAL	3.7
1	B	53	ALA	3.7
1	B	152	PHE	3.7
1	B	36	LEU	3.7
1	B	122	PHE	3.7
1	B	110	VAL	3.7
1	B	14	SER	3.6
1	B	17	PRO	3.6
1	B	107	LEU	3.6
1	B	69	VAL	3.6
1	B	18	LEU	3.6
1	B	67	THR	3.6
1	B	28	GLN	3.6
1	B	128	GLN	3.5
1	B	103	LEU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	176	ILE	3.5
1	B	30	LYS	3.5
1	B	137	LEU	3.5
1	B	70	LEU	3.4
1	B	82	THR	3.4
1	B	162	PRO	3.4
1	B	96	ASN	3.4
1	B	138	TYR	3.4
1	B	174	GLN	3.4
1	B	87	SER	3.3
1	B	57	GLU	3.3
1	B	74	LYS	3.3
1	B	153	ILE	3.3
1	B	158	SER	3.3
1	B	123	PHE	3.3
1	B	55	ILE	3.3
1	C	79	TRP	3.3
1	B	15	LYS	3.2
1	B	76	CYS	3.2
1	B	130	ARG	3.1
1	B	9	PRO	3.1
1	B	34	VAL	3.1
1	B	149	LYS	3.1
1	B	27	PHE	3.1
1	B	42	LEU	3.0
1	A	79	TRP	3.0
1	B	89	PRO	3.0
1	B	35	GLY	3.0
1	B	40	ALA	3.0
1	B	139	GLY	3.0
1	B	106	TYR	3.0
1	B	51	MET	3.0
1	B	16	VAL	3.0
1	B	155	PHE	3.0
1	B	80	ILE	2.9
1	B	113	THR	2.9
1	B	148	LEU	2.9
1	B	131	GLU	2.9
1	B	173	ASP	2.9
1	B	119	ALA	2.9
1	B	125	LYS	2.9
1	B	91	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	23	GLN	2.8
1	B	75	LYS	2.8
1	B	126	VAL	2.8
1	B	13	LEU	2.7
1	B	102	GLY	2.7
1	B	127	SER	2.7
1	B	136	THR	2.7
1	B	41	ILE	2.7
1	B	90	GLY	2.7
1	C	3	SER	2.7
1	B	58	LEU	2.6
1	B	157	LYS	2.6
1	B	111	VAL	2.6
1	C	143	GLU	2.6
1	B	81	ARG	2.5
1	C	46	LYS	2.5
1	B	12	PRO	2.5
1	B	100	TYR	2.5
1	B	66	VAL	2.5
1	A	42	LEU	2.5
1	B	52	TYR	2.5
1	C	174	GLN	2.5
1	B	120	MET	2.4
1	B	99	SER	2.4
1	B	168	PHE	2.4
1	C	96	ASN	2.3
1	B	150	GLU	2.3
1	B	161	LEU	2.3
1	B	156	SER	2.3
1	C	45	ASP	2.3
1	B	117	GLN	2.3
1	B	21	ASN	2.3
1	B	108	VAL	2.2
1	A	178	GLY	2.2
1	B	105	SER	2.2
1	B	163	GLU	2.1
1	B	10	ALA	2.1
1	B	72	ARG	2.1
1	B	84	VAL	2.1
1	B	101	PRO	2.1
1	A	21	ASN	2.1
1	B	147	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	19	GLN	2.1
1	B	133	PHE	2.1
1	A	4	THR	2.0
1	B	29	GLY	2.0
1	A	146	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	C	181	6/6	0.55	0.21	74,74,74,74	0
6	SO4	A	182	5/5	0.72	0.22	92,92,92,92	0
7	MCK	C	179	39/39	0.75	0.21	59,70,73,73	0
4	GOL	A	180	6/6	0.76	0.21	53,54,54,55	0
3	DBH	A	179	11/11	0.76	0.23	101,101,101,102	0
4	GOL	C	183	6/6	0.79	0.17	43,43,44,44	0
4	GOL	A	3731	6/6	0.80	0.16	59,60,60,60	0
4	GOL	C	182	6/6	0.82	0.20	70,71,71,71	0
2	DBS	A	301	17/17	0.83	0.17	58,61,65,65	0
3	DBH	A	303	11/11	0.83	0.16	88,88,88,88	0
4	GOL	C	180	6/6	0.84	0.16	67,67,67,67	0
4	GOL	C	3731	6/6	0.84	0.18	55,56,57,57	0
5	NA	A	181	1/1	0.95	0.16	23,23,23,23	0
5	NA	C	184	1/1	0.95	0.21	33,33,33,33	0
8	CL	C	185	1/1	0.96	0.11	67,67,67,67	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.