



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 26, 2026 – 05:54 AM UTC

PDB ID : 6K3J / pdb_00006k3j
BMRB ID : 36194
Title : Solution structure of APOBEC3G-CD2 with ssDNA, Product A
Authors : Cao, C.; Yan, X.; Lan, W.; Wang, C.
Deposited on : 2019-05-19

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 78%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	188	61% 37% .
2	B	10	40% 50% 10%

2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:197-A:384 (188)	1.24	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 8, 9, 12, 15, 17
2	2, 5, 10, 16, 19
3	3, 7, 11, 14, 20
4	4, 6
Single-model clusters	13; 18

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3332 atoms, of which 1576 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called DNA dC->dU-editing enzyme APOBEC-3G.

Mol	Chain	Residues	Atoms							Trace
1	A	188	Total	C	H	N	O	S		0
			3020	985	1463	275	283	14		

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*TP*TP*CP*UP*(IUR)P*AP*AP*TP*T)-3').

Mol	Chain	Residues	Atoms							Trace
2	B	10	Total	C	H	I	N	O	P	0
			311	97	113	1	30	61	9	

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
3	A	1	Total	Zn
			1	1

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3G




- Molecule 2: DNA (5'-D(*AP*TP*TP*CP*UP*(IUR)P*AP*AP*TP*T)-3')




4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3G



- Molecule 2: DNA (5'-D(*AP*TP*TP*CP*UP*(IUR)P*AP*AP*TP*T)-3')

Chain B:  40% 50% 10%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	refinement	
X-PLOR NIH	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	2221
Number of shifts mapped to atoms	2221
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	78%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5IU, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.43±0.01	0±0/1604 (0.0± 0.0%)	1.30±0.00	1±1/2171 (0.0± 0.0%)
2	B	0.73±0.01	1±0/197 (0.5± 0.0%)	0.88±0.04	0±0/299 (0.0± 0.0%)
All	All	1.37	20/36020 (0.1%)	1.26	14/49400 (0.0%)

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	B	388	DC	O3'-P	6.08	1.62	1.56	15	20

5 of 7 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	350	PHE	CA-CB-CG	-5.39	108.41	113.80	7	2
1	A	241	PHE	CA-CB-CG	-5.26	108.54	113.80	6	5
1	A	343	PHE	CA-CB-CG	-5.25	108.55	113.80	14	1
1	A	206	PHE	CA-CB-CG	-5.24	108.56	113.80	16	2
1	A	355	GLY	N-CA-C	-5.14	108.61	115.40	10	2

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1557	1463	1452	84±9
2	B	198	113	113	15±4
All	All	35120	31520	31300	1734

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

5 of 706 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:253:LEU:HD22	1:A:253:LEU:H	0.93	1.24	5	1
1:A:281:CYS:SG	1:A:309:ILE:HG23	0.86	2.09	6	6
1:A:299:ILE:HD12	1:A:331:ALA:O	0.84	1.73	12	14
1:A:208:ASN:O	1:A:212:VAL:HG13	0.83	1.73	14	3
1:A:209:GLU:O	1:A:212:VAL:HG13	0.81	1.75	15	3

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	186/188 (99%)	172±1 (92±1%)	10±2 (5±1%)	5±1 (2±0%)	6	43
All	All	3720/3760 (99%)	3430 (92%)	198 (5%)	92 (2%)	6	43

5 of 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	199	PRO	20
1	A	200	PRO	20
1	A	382	GLN	17
1	A	320	ARG	13
1	A	212	VAL	9

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/169 (100%)	156±2 (92±1%)	13±2 (8±1%)	13	61
All	All	3380/3380 (100%)	3112 (92%)	268 (8%)	13	61

5 of 48 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	198	ASP	20
1	A	279	VAL	20
1	A	335	ILE	18
1	A	286	SER	17
1	A	281	CYS	16

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	5IU	B	390	2	18,21,22	0.41±0.05	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types,

if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	5IU	B	390	2	25,30,33	0.46±0.03	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5IU	B	390	2	-	0±0,7,21,22	0±0,2,2,2

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 78% for the well-defined parts and 78% for the entire structure.

7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: `data_A3G_PrC.str`

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	2221
Number of shifts mapped to atoms	2221
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	17

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	175	-0.27 ± 0.17	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	165	0.69 ± 0.10	Should be checked
$^{13}\text{C}'$	160	-0.10 ± 0.16	None needed (< 0.5 ppm)
^{15}N	171	-0.19 ± 0.44	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 78%, i.e. 2177 atoms were assigned a chemical shift out of a possible 2784. 0 out of 23 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	864/934 (93%)	358/378 (95%)	335/376 (89%)	171/180 (95%)
Sidechain	1049/1352 (78%)	743/869 (86%)	295/420 (70%)	11/63 (17%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	205/328 (62%)	115/162 (71%)	84/143 (59%)	6/23 (26%)
Sugar	50/108 (46%)	50/63 (79%)	0/45 (0%)	0/0 (—%)
Base	9/62 (15%)	9/35 (26%)	0/18 (0%)	0/9 (0%)
Overall	2177/2784 (78%)	1275/1507 (85%)	714/1002 (71%)	188/275 (68%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	206	PHE	CD2	122.14	125.53 – 137.61	-7.8
1	A	340	TYR	CE2	127.67	111.68 – 124.17	7.8
1	A	206	PHE	CD1	122.14	125.33 – 137.83	-7.5
1	A	310	PHE	CD2	122.67	125.53 – 137.61	-7.4
1	A	340	TYR	CE1	127.67	111.24 – 124.66	7.2
1	A	310	PHE	CD1	122.67	125.33 – 137.83	-7.1
1	A	314	ILE	CG2	26.07	10.93 – 24.12	6.5
1	A	247	PRO	CD	43.58	45.11 – 55.58	-6.5
1	A	249	LYS	CD	22.29	23.50 – 34.42	-6.1
1	A	325	LEU	CG	33.09	21.37 – 32.19	5.8
1	A	222	TYR	CE2	125.05	111.68 – 124.17	5.7
1	A	355	GLY	H	11.65	5.23 – 11.42	5.4
1	A	219	TYR	CD1	125.40	125.84 – 139.60	-5.3
1	A	222	TYR	CE1	125.05	111.24 – 124.66	5.3
1	A	358	PHE	CD2	125.21	125.53 – 137.61	-5.3
1	A	358	PHE	CD1	125.21	125.33 – 137.83	-5.1
1	A	294	GLU	CG	30.13	30.20 – 42.01	-5.1

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	2702
Intra-residue ($ i-j =0$)	1298
Sequential ($ i-j =1$)	571
Medium range ($ i-j >1$ and $ i-j <5$)	358
Long range ($ i-j \geq 5$)	432
Inter-chain	43
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	13.6
Number of long range restraints per residue ¹	2.2

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	49.4	0.2
0.2-0.5 (Medium)	16.4	0.5
>0.5 (Large)	25.6	4.5

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	1298	48.0	73	5.6	2.7	16	1.2	0.6
Backbone-Backbone	198	7.3	2	1.0	0.1	1	0.5	0.0
Backbone-Sidechain	691	25.6	62	9.0	2.3	13	1.9	0.5
Sidechain-Sidechain	409	15.1	9	2.2	0.3	2	0.5	0.1
Sequential (i-j =1)	571	21.1	26	4.6	1.0	1	0.2	0.0
Backbone-Backbone	271	10.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	281	10.4	23	8.2	0.9	1	0.4	0.0
Sidechain-Sidechain	19	0.7	3	15.8	0.1	0	0.0	0.0
Medium range (i-j >1 & i-j <5)	358	13.2	22	6.1	0.8	3	0.8	0.1
Backbone-Backbone	200	7.4	6	3.0	0.2	2	1.0	0.1
Backbone-Sidechain	104	3.8	10	9.6	0.4	1	1.0	0.0
Sidechain-Sidechain	54	2.0	6	11.1	0.2	0	0.0	0.0
Long range (i-j ≥5)	432	16.0	68	15.7	2.5	11	2.5	0.4
Backbone-Backbone	66	2.4	9	13.6	0.3	1	1.5	0.0
Backbone-Sidechain	165	6.1	30	18.2	1.1	3	1.8	0.1
Sidechain-Sidechain	201	7.4	29	14.4	1.1	7	3.5	0.3
Inter-chain	43	1.6	20	46.5	0.7	1	2.3	0.0
Backbone-Backbone	9	0.3	5	55.6	0.2	1	11.1	0.0
Backbone-Sidechain	31	1.1	15	48.4	0.6	0	0.0	0.0
Sidechain-Sidechain	3	0.1	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	2702	100.0	209	7.7	7.7	32	1.2	1.2
Backbone-Backbone	744	27.5	22	3.0	0.8	5	0.7	0.2
Backbone-Sidechain	1272	47.1	140	11.0	5.2	18	1.4	0.7
Sidechain-Sidechain	686	25.4	47	6.9	1.7	9	1.3	0.3

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	39	10	9	32	8	98	0.37	3.63	0.47	0.18
2	39	8	8	33	4	92	0.44	2.38	0.47	0.19
3	43	11	9	33	5	101	0.42	2.89	0.51	0.19
4	38	11	10	36	7	102	0.37	2.65	0.42	0.16
5	39	10	9	32	6	96	0.4	2.28	0.44	0.18
6	39	11	11	34	4	99	0.4	4.5	0.54	0.19
7	38	13	9	32	6	98	0.4	2.6	0.47	0.16
8	39	15	9	33	6	102	0.43	2.68	0.52	0.18
9	37	10	9	33	6	95	0.43	3.05	0.53	0.18
10	36	8	8	32	6	90	0.47	2.55	0.52	0.19

Continued on next page...


Continued from previous page...

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
11	40	11	8	31	6	96	0.4	2.65	0.49	0.18
12	36	8	10	34	8	96	0.42	3.42	0.52	0.18
13	36	11	9	33	8	97	0.46	2.64	0.57	0.19
14	36	6	8	36	7	93	0.44	2.63	0.47	0.18
15	40	8	6	40	6	100	0.39	2.75	0.47	0.16
16	38	8	8	34	5	93	0.42	2.71	0.49	0.19
17	39	13	6	29	7	94	0.41	3.06	0.51	0.18
18	36	8	7	38	6	95	0.43	2.42	0.49	0.18
19	39	7	8	32	6	92	0.41	2.6	0.5	0.18
20	39	5	6	37	5	92	0.41	2.72	0.49	0.18

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model ⓘ



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 2493(IR:1225, SQ:545, MR:336, LR:364, IC:23) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
13	5	6	12	3	39	1	5.0
3	1	3	5	3	15	2	10.0
2	5	0	3	2	12	3	15.0
3	1	1	3	1	9	4	20.0
2	1	1	4	1	9	5	25.0
4	0	1	2	2	9	6	30.0
4	1	2	3	4	14	7	35.0
4	3	0	2	0	9	8	40.0
3	0	0	1	0	4	9	45.0
0	1	1	1	1	4	10	50.0
2	0	1	2	0	5	11	55.0
5	2	0	2	1	10	12	60.0
1	1	2	2	0	6	13	65.0
2	2	0	1	0	5	14	70.0
0	0	0	3	0	3	15	75.0
1	1	0	3	1	6	16	80.0
1	0	0	2	0	3	17	85.0
1	0	0	4	0	5	18	90.0
6	1	1	2	0	10	19	95.0
16	1	3	11	1	32	20	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble [i](#)



9.4 Most violated distance restraints in the ensemble [i](#)

9.4.1 Histogram : Distribution of mean distance violations [i](#)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,611)	1:245:A:GLN:H	1:256:A:ARG:H	20	2.34	0.53	2.6
(1,2670)	2:386:B:DT:H5'	1:366:A:GLU:HA	20	2.22	0.9	2.13
(1,2670)	2:386:B:DT:H5''	1:366:A:GLU:HA	20	2.22	0.9	2.13
(1,567)	1:241:A:PHE:HZ	1:247:A:PRO:HA	20	1.34	0.03	1.35
(1,620)	1:246:A:ALA:HB1	1:252:A:PHE:HB2	20	1.1	0.23	1.19
(1,620)	1:246:A:ALA:HB1	1:252:A:PHE:HB3	20	1.1	0.23	1.19
(1,620)	1:246:A:ALA:HB2	1:252:A:PHE:HB2	20	1.1	0.23	1.19
(1,620)	1:246:A:ALA:HB2	1:252:A:PHE:HB3	20	1.1	0.23	1.19
(1,620)	1:246:A:ALA:HB3	1:252:A:PHE:HB2	20	1.1	0.23	1.19
(1,620)	1:246:A:ALA:HB3	1:252:A:PHE:HB3	20	1.1	0.23	1.19
(1,857)	1:263:A:LEU:HD11	1:298:A:PHE:H	20	1.09	0.18	1.08
(1,857)	1:263:A:LEU:HD12	1:298:A:PHE:H	20	1.09	0.18	1.08
(1,857)	1:263:A:LEU:HD13	1:298:A:PHE:H	20	1.09	0.18	1.08
(1,857)	1:263:A:LEU:HD21	1:298:A:PHE:H	20	1.09	0.18	1.08
(1,857)	1:263:A:LEU:HD22	1:298:A:PHE:H	20	1.09	0.18	1.08
(1,857)	1:263:A:LEU:HD23	1:298:A:PHE:H	20	1.09	0.18	1.08

Continued on next page...

Continued from previous page...


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,182)	1:211:A:TRP:HE1	1:284:A:SER:HB2	20	1.07	0.03	1.08
(1,182)	1:211:A:TRP:HE1	1:284:A:SER:HB3	20	1.07	0.03	1.08
(1,662)	1:249:A:LYS:HD2	1:254:A:GLU:HG2	20	1.06	0.05	1.06
(1,662)	1:249:A:LYS:HD2	1:254:A:GLU:HG3	20	1.06	0.05	1.06
(1,662)	1:249:A:LYS:HD3	1:254:A:GLU:HG2	20	1.06	0.05	1.06
(1,662)	1:249:A:LYS:HD3	1:254:A:GLU:HG3	20	1.06	0.05	1.06
(1,1648)	1:313:A:ARG:HD2	1:337:A:ILE:HD11	20	1.06	0.05	1.06
(1,1648)	1:313:A:ARG:HD2	1:337:A:ILE:HD12	20	1.06	0.05	1.06
(1,1648)	1:313:A:ARG:HD2	1:337:A:ILE:HD13	20	1.06	0.05	1.06
(1,1648)	1:313:A:ARG:HD3	1:337:A:ILE:HD11	20	1.06	0.05	1.06
(1,1648)	1:313:A:ARG:HD3	1:337:A:ILE:HD12	20	1.06	0.05	1.06
(1,1648)	1:313:A:ARG:HD3	1:337:A:ILE:HD13	20	1.06	0.05	1.06
(1,2628)	2:387:B:DT:H6	2:387:B:DT:H5'	20	0.81	0.24	0.84
(1,2628)	2:387:B:DT:H6	2:387:B:DT:H5''	20	0.81	0.24	0.84
(1,1437)	1:301:A:LYS:HG2	1:309:A:ILE:HD11	20	0.77	0.2	0.81

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,2670)	2:386:B:DT:H5'	1:366:A:GLU:HA	6	4.5
(1,2670)	2:386:B:DT:H5''	1:366:A:GLU:HA	6	4.5
(1,2670)	2:386:B:DT:H5'	1:366:A:GLU:HA	1	3.63
(1,2670)	2:386:B:DT:H5''	1:366:A:GLU:HA	1	3.63
(1,2670)	2:386:B:DT:H5'	1:366:A:GLU:HA	12	3.42
(1,2670)	2:386:B:DT:H5''	1:366:A:GLU:HA	12	3.42
(1,611)	1:245:A:GLN:H	1:256:A:ARG:H	17	3.06
(1,2670)	2:386:B:DT:H5'	1:366:A:GLU:HA	9	3.05
(1,2670)	2:386:B:DT:H5''	1:366:A:GLU:HA	9	3.05
(1,611)	1:245:A:GLN:H	1:256:A:ARG:H	3	2.89
(1,2670)	2:386:B:DT:H5'	1:366:A:GLU:HA	15	2.75
(1,2670)	2:386:B:DT:H5''	1:366:A:GLU:HA	15	2.75
(1,611)	1:245:A:GLN:H	1:256:A:ARG:H	20	2.72
(1,611)	1:245:A:GLN:H	1:256:A:ARG:H	16	2.71
(1,611)	1:245:A:GLN:H	1:256:A:ARG:H	8	2.68

10 Dihedral-angle violation analysis ⓘ

No dihedral-angle restraints found