



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:25 PM UTC

PDB ID : 3K40 / pdb_00003k40
Title : Crystal structure of Drosophila 3,4-dihydroxyphenylalanine decarboxylase
Authors : Han, Q.; Ding, H.; Robinson, H.; Christensen, B.M.; Li, J.
Deposited on : 2009-10-05
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	6% 77% 16% • 6%
1	B	475	7% 84% 8% • 6%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7715 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aromatic-L-amino-acid decarboxylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	448	Total	C	N	O	P	S	0	0	0
			3554	2272	610	643	1	28			
1	B	448	Total	C	N	O	P	S	0	0	0
			3554	2272	610	643	1	28			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	344	Total	O	0	0
			344	344		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	257	Total 257	O 257	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aromatic-L-amino-acid decarboxylase



• Molecule 1: Aromatic-L-amino-acid decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.81Å 108.60Å 86.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.88 – 1.75 29.88 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.9 (29.88-1.75) 98.8 (29.88-1.75)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.60 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.197 , 0.226 0.196 , 0.225	Depositor DCC
R_{free} test set	4972 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.009 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7715	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	0/3618	0.90	3/4891 (0.1%)
1	B	0.71	1/3618 (0.0%)	0.87	0/4891
All	All	0.76	1/7236 (0.0%)	0.89	3/9782 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	452	THR	C-O	-5.27	1.17	1.23

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	317	ASP	CA-C-N	8.24	130.13	119.84
1	A	317	ASP	C-N-CA	8.24	130.13	119.84
1	A	461	TRP	CA-CB-CG	-5.17	103.77	113.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3554	0	3499	62	0
1	B	3554	0	3499	33	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	6	0	8	3	0
3	A	344	0	0	10	0
3	B	257	0	0	4	0
All	All	7715	0	7006	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:350:LEU:HD21	3:A:584:HOH:O	1.65	0.97
1:A:313:MET:HE3	1:A:315:LEU:HD11	1.49	0.95
1:A:455:GLU:HG2	3:A:789:HOH:O	1.80	0.79
1:A:352:ARG:HB2	3:A:541:HOH:O	1.83	0.78
1:B:54:GLN:HE21	1:B:54:GLN:H	1.34	0.75
1:B:436:LYS:HE2	1:B:439:ASP:HA	1.71	0.73
1:A:105:TRP:HZ3	1:A:349:PRO:HD2	1.55	0.72
1:A:143:VAL:CG1	1:A:313:MET:HE2	2.22	0.70
1:A:54:GLN:HE21	1:A:54:GLN:H	1.35	0.70
1:A:436:LYS:HE2	1:A:439:ASP:HA	1.73	0.69
3:A:585:HOH:O	1:B:49:LYS:HD3	1.93	0.69
1:A:423:ARG:HH21	1:A:470:GLU:HB3	1.58	0.69
1:A:38:LEU:HD11	1:A:63:VAL:CG1	2.25	0.66
1:B:75:LYS:HE3	3:B:523:HOH:O	1.95	0.66
1:A:101:ILE:HD12	1:B:308:PHE:HB2	1.76	0.65
1:A:308:PHE:HE1	1:B:353:ARG:HE	1.42	0.64
1:A:104:THR:HB	1:B:79:TYR:HD2	1.63	0.63
1:A:38:LEU:CD1	1:A:63:VAL:HG13	2.30	0.62
1:A:79:TYR:HD2	1:B:104:THR:HG23	1.63	0.61
1:A:384:GLN:HE22	1:A:454:SER:HB2	1.66	0.61
1:A:1:MET:N	2:A:476:GOL:H31	2.16	0.60
1:B:384:GLN:NE2	3:B:702:HOH:O	2.34	0.60
1:B:217:ARG:HH12	1:B:255:GLU:CD	2.10	0.60
1:A:383:LYS:NZ	3:A:664:HOH:O	2.35	0.60
1:B:217:ARG:NH1	1:B:255:GLU:OE1	2.35	0.59
1:A:104:THR:HG21	1:B:78:ALA:HB1	1.85	0.58
1:A:39:LYS:HD3	3:A:539:HOH:O	2.04	0.57
1:A:423:ARG:CD	1:A:471:MET:HG2	2.36	0.56
1:A:1:MET:H3	2:A:476:GOL:H31	1.70	0.55
1:A:49:LYS:HE3	1:A:50:PRO:HD2	1.88	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:LEU:HD23	1:A:357:LEU:HD21	1.90	0.54
1:B:34:LYS:H	1:B:34:LYS:CE	2.21	0.53
1:A:377:ARG:HG3	3:A:603:HOH:O	2.09	0.53
1:A:79:TYR:HB2	1:B:104:THR:OG1	2.09	0.52
1:A:84:ASN:HD22	1:A:450:ARG:HE	1.57	0.52
1:A:1:MET:N	2:A:476:GOL:C3	2.72	0.52
1:A:139:LYS:O	1:A:318:PRO:HD2	2.08	0.52
1:B:1:MET:N	3:B:530:HOH:O	2.42	0.51
1:A:261:ASN:HD21	1:A:292:SER:HB2	1.75	0.51
1:A:38:LEU:CD1	1:A:63:VAL:CG1	2.88	0.51
1:A:285:HIS:HD2	3:A:571:HOH:O	1.94	0.50
1:A:473:GLN:C	1:A:475:GLN:H	2.19	0.50
1:B:34:LYS:H	1:B:34:LYS:NZ	2.10	0.50
1:B:54:GLN:H	1:B:54:GLN:NE2	2.06	0.50
1:A:257:GLY:HA3	1:A:292:SER:OG	2.12	0.50
1:B:455:GLU:H	1:B:455:GLU:CD	2.20	0.49
1:A:260:GLY:HA3	1:A:267:ILE:HD11	1.94	0.49
1:A:421:LEU:HD22	1:A:434:PRO:HB3	1.94	0.49
1:A:272:ALA:HB3	1:A:302:LLP:C3	2.43	0.49
1:A:189:ASP:OD1	1:A:210:GLN:NE2	2.47	0.47
1:A:91:ALA:HB2	1:A:361:PHE:CD2	2.50	0.47
1:B:106:ILE:CG2	1:B:108:SER:HB3	2.44	0.47
1:A:427:ARG:HD3	1:A:463:GLU:OE1	2.14	0.47
1:A:423:ARG:CD	1:A:471:MET:CG	2.93	0.47
1:B:94:LEU:HD23	1:B:357:LEU:HD21	1.96	0.47
1:B:272:ALA:HB3	1:B:302:LLP:C3	2.46	0.46
1:A:261:ASN:ND2	1:A:292:SER:HB2	2.30	0.46
1:B:472:GLU:C	1:B:474:GLU:H	2.24	0.46
1:B:162:LYS:HG3	1:B:320:TRP:CZ2	2.51	0.46
1:B:301:HIS:HA	1:B:306:VAL:O	2.15	0.46
1:B:308:PHE:HA	1:B:309:ASP:HA	1.65	0.46
1:A:302:LLP:O3	1:A:302:LLP:NZ	2.49	0.46
1:A:286:LEU:CD1	1:A:403:MET:HE2	2.47	0.45
1:A:111:CYS:HB3	1:A:351:GLY:HA3	1.98	0.45
1:A:145:GLN:HE21	1:A:313:MET:CE	2.30	0.45
1:A:108:SER:HB2	1:B:69:THR:OG1	2.16	0.45
1:A:39:LYS:HB3	1:A:40:PRO:HD3	1.99	0.44
1:A:423:ARG:HG3	3:A:719:HOH:O	2.16	0.44
1:A:423:ARG:HD3	1:A:471:MET:CG	2.47	0.44
1:A:54:GLN:H	1:A:54:GLN:NE2	2.09	0.44
1:A:278:PHE:HB3	1:A:287:MET:HE2	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:261:ASN:HD21	1:A:292:SER:CB	2.31	0.43
1:A:308:PHE:HA	1:A:309:ASP:HA	1.78	0.43
1:A:105:TRP:HZ3	1:A:349:PRO:CD	2.29	0.43
1:B:415:GLU:H	1:B:415:GLU:CD	2.27	0.43
1:B:423:ARG:HG3	3:B:508:HOH:O	2.17	0.43
1:B:34:LYS:HB2	1:B:34:LYS:HE2	1.71	0.42
1:A:184:VAL:HG13	1:A:235:PRO:HA	2.01	0.42
1:B:427:ARG:HD3	1:B:463:GLU:OE1	2.20	0.42
1:A:423:ARG:NH2	1:A:470:GLU:HB3	2.31	0.41
1:A:96:GLY:HA3	3:A:736:HOH:O	2.20	0.41
1:A:286:LEU:CD1	1:A:403:MET:CE	2.99	0.41
1:A:34:LYS:O	1:A:37:TYR:HB3	2.20	0.41
1:A:301:HIS:HA	1:A:306:VAL:O	2.21	0.41
1:B:272:ALA:HA	1:B:298:PHE:HA	2.02	0.41
1:A:65:MET:N	1:A:66:PRO:CD	2.84	0.41
1:B:139:LYS:O	1:B:318:PRO:CD	2.69	0.41
1:B:156:LEU:HD23	1:B:156:LEU:C	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	443/475 (93%)	435 (98%)	8 (2%)	0	100	100
1	B	443/475 (93%)	430 (97%)	13 (3%)	0	100	100
All	All	886/950 (93%)	865 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	371/395 (94%)	360 (97%)	11 (3%)	36	16
1	B	371/395 (94%)	357 (96%)	14 (4%)	29	10
All	All	742/790 (94%)	717 (97%)	25 (3%)	32	13

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	LYS
1	A	54	GLN
1	A	75	LYS
1	A	104	THR
1	A	164	LEU
1	A	169	GLU
1	A	170	LEU
1	A	197	ARG
1	A	390	VAL
1	A	415	GLU
1	A	474	GLU
1	B	34	LYS
1	B	54	GLN
1	B	61	GLU
1	B	75	LYS
1	B	101	ILE
1	B	133	LEU
1	B	164	LEU
1	B	176	GLU
1	B	217	ARG
1	B	231	GLU
1	B	415	GLU
1	B	443	LEU
1	B	455	GLU
1	B	473	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	77	HIS
1	A	84	ASN
1	A	145	GLN
1	A	210	GLN
1	A	261	ASN
1	A	285	HIS
1	A	384	GLN
1	A	473	GLN
1	B	54	GLN
1	B	77	HIS
1	B	84	ASN
1	B	210	GLN
1	B	214	HIS
1	B	261	ASN
1	B	285	HIS
1	B	384	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LLP	B	302	1	23,24,25	1.57	5 (21%)	25,32,34	1.52	5 (20%)
1	LLP	A	302	1	23,24,25	1.53	5 (21%)	25,32,34	1.82	7 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	B	302	1	-	2/16/17/19	0/1/1/1
1	LLP	A	302	1	-	2/16/17/19	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	302	LLP	O3-C3	-4.80	1.25	1.36
1	A	302	LLP	O3-C3	-4.70	1.26	1.36
1	A	302	LLP	C6-N1	2.72	1.40	1.34
1	B	302	LLP	C6-N1	2.64	1.39	1.34
1	B	302	LLP	C4-C4'	2.63	1.52	1.46
1	A	302	LLP	C4'-NZ	2.60	1.35	1.27
1	B	302	LLP	P-OP2	-2.57	1.45	1.54
1	A	302	LLP	P-OP2	-2.37	1.46	1.54
1	B	302	LLP	C2-N1	2.28	1.37	1.33
1	A	302	LLP	C4-C4'	2.08	1.51	1.46

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	302	LLP	C4-C3-C2	-4.03	117.88	120.14
1	A	302	LLP	C5-C6-N1	-3.75	117.74	123.83
1	B	302	LLP	OP3-P-OP2	3.40	120.55	107.80
1	A	302	LLP	CE-NZ-C4'	-3.35	107.98	118.72
1	A	302	LLP	C5'-C5-C6	-3.08	114.35	119.36
1	B	302	LLP	C5-C6-N1	-2.69	119.46	123.83
1	B	302	LLP	C4-C4'-NZ	-2.55	112.26	124.04
1	B	302	LLP	C4-C3-C2	-2.46	118.76	120.14
1	A	302	LLP	C4-C4'-NZ	-2.44	112.78	124.04
1	A	302	LLP	C6-C5-C4	2.39	122.33	118.21
1	B	302	LLP	OP2-P-OP4	-2.28	100.73	106.67
1	A	302	LLP	OP4-C5'-C5	2.10	113.29	109.36

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	302	LLP	C4-C4'-NZ-CE
1	B	302	LLP	C4-C4'-NZ-CE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	302	LLP	C-CA-CB-CG
1	B	302	LLP	C-CA-CB-CG

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	302	LLP	1	0
1	A	302	LLP	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	A	476	-	5,5,5	0.24	0	5,5,5	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	476	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	476	GOL	O1-C1-C2-C3
2	A	476	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	476	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	447/475 (94%)	0.25	30 (6%) 24 28	14, 24, 46, 69	0
1	B	447/475 (94%)	0.55	32 (7%) 21 25	16, 29, 55, 75	0
All	All	894/950 (94%)	0.40	62 (6%) 23 26	14, 27, 50, 75	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	321	VAL	6.5
1	B	105	TRP	6.5
1	B	101	ILE	5.7
1	B	103	PHE	5.5
1	B	104	THR	5.3
1	A	349	PRO	4.9
1	A	105	TRP	4.8
1	B	106	ILE	4.7
1	A	107	ALA	4.5
1	B	170	LEU	4.4
1	A	174	TRP	4.3
1	A	104	THR	4.3
1	B	174	TRP	4.2
1	A	79	TYR	4.1
1	B	107	ALA	3.9
1	A	103	PHE	3.7
1	A	106	ILE	3.6
1	B	321	VAL	3.6
1	A	320	TRP	3.5
1	B	172	PRO	3.5
1	A	350	LEU	3.5
1	B	167	VAL	3.5
1	B	102	GLY	3.2
1	B	232	GLY	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	229	VAL	3.1
1	B	180	LEU	3.0
1	B	475	GLN	3.0
1	B	79	TYR	3.0
1	A	101	ILE	3.0
1	A	475	GLN	3.0
1	A	352	ARG	3.0
1	B	349	PRO	2.9
1	B	350	LEU	2.9
1	B	179	ILE	2.9
1	A	32	GLU	2.7
1	A	439	ASP	2.7
1	B	352	ARG	2.6
1	A	438	LYS	2.6
1	A	170	LEU	2.6
1	A	28	ARG	2.6
1	A	108	SER	2.6
1	A	44	ASP	2.6
1	B	80	PHE	2.6
1	B	318	PRO	2.6
1	B	32	GLU	2.6
1	B	100	CYS	2.5
1	A	318	PRO	2.5
1	B	320	TRP	2.5
1	B	440	VAL	2.4
1	A	208	SER	2.4
1	A	102	GLY	2.3
1	A	38	LEU	2.3
1	B	230	ALA	2.3
1	A	468	ALA	2.2
1	A	178	THR	2.2
1	B	473	GLN	2.1
1	B	234	ILE	2.1
1	A	172	PRO	2.1
1	B	175	ASP	2.0
1	A	171	HIS	2.0
1	B	226	GLU	2.0
1	A	471	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	LLP	B	302	24/25	0.96	0.07	21,24,26,26	0
1	LLP	A	302	24/25	0.97	0.06	14,18,21,22	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	476	6/6	0.67	0.20	54,55,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.