



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 17, 2026 – 07:12 PM UTC

PDB ID : 3K4M / pdb_00003k4m
Title : Pyranose 2-oxidase Y456W mutant in complex with 2FG
Authors : Divne, C.; Tan, T.C.
Deposited on : 2009-10-05
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	623	70% 21% 8%
1	B	623	72% 18% 7%
1	C	623	73% 18% 8%
1	D	623	72% 18% 8%
1	E	623	70% 20% 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	623	 2% 71% 19% 8%
1	G	623	 % 70% 20% 8%
1	H	623	 % 72% 17% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FAD	A	801	X	-	-	-
2	FAD	B	801	X	-	-	-
2	FAD	C	801	X	-	-	-
2	FAD	E	801	X	-	-	-
2	FAD	F	801	X	-	-	-
2	FAD	H	801	X	-	-	-
3	SHG	B	901	X	-	-	-
3	SHG	D	901	X	-	-	-
3	SHG	E	901	X	-	X	-
3	SHG	G	901	X	-	-	-
3	SHG	H	901	X	-	-	-
4	MES	B	902	-	-	X	-
4	MES	C	624	-	-	X	-
4	MES	C	902	-	-	X	-
4	MES	E	902	-	-	X	-
4	MES	H	624	-	-	X	-
4	MES	H	902	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 38766 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyranose 2-oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	574	Total	C	N	O	S	0	0	0
			4528	2861	776	867	24			
1	B	577	Total	C	N	O	S	0	0	0
			4551	2874	779	873	25			
1	C	574	Total	C	N	O	S	0	0	0
			4527	2859	776	868	24			
1	D	574	Total	C	N	O	S	0	0	0
			4527	2859	776	868	24			
1	E	576	Total	C	N	O	S	0	0	0
			4544	2870	778	871	25			
1	F	573	Total	C	N	O	S	0	0	0
			4520	2855	775	866	24			
1	G	573	Total	C	N	O	S	0	0	0
			4520	2855	775	866	24			
1	H	573	Total	C	N	O	S	0	0	0
			4520	2855	775	866	24			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	456	TRP	TYR	engineered mutation	UNP Q7ZA32
B	456	TRP	TYR	engineered mutation	UNP Q7ZA32
C	456	TRP	TYR	engineered mutation	UNP Q7ZA32
D	456	TRP	TYR	engineered mutation	UNP Q7ZA32
E	456	TRP	TYR	engineered mutation	UNP Q7ZA32
F	456	TRP	TYR	engineered mutation	UNP Q7ZA32
G	456	TRP	TYR	engineered mutation	UNP Q7ZA32
H	456	TRP	TYR	engineered mutation	UNP Q7ZA32

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	B	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	C	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	D	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	E	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	F	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	G	1	Total 53	C 27	N 9	O 15	P 2	0	0
2	H	1	Total 53	C 27	N 9	O 15	P 2	0	0

- Molecule 3 is 2-deoxy-2-fluoro-beta-D-glucopyranose (CCD ID: SHG) (formula: $\text{C}_6\text{H}_{11}\text{FO}_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	F	O	0	0
			12	6	1	5		
3	B	1	Total	C	F	O	0	0
			12	6	1	5		
3	C	1	Total	C	F	O	0	0
			12	6	1	5		
3	D	1	Total	C	F	O	0	0
			12	6	1	5		
3	E	1	Total	C	F	O	0	0
			12	6	1	5		
3	F	1	Total	C	F	O	0	0
			12	6	1	5		
3	G	1	Total	C	F	O	0	0
			12	6	1	5		
3	H	1	Total	C	F	O	0	0
			12	6	1	5		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	F	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	H	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	H	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	303	Total	O	0	0
			303	303		
5	B	277	Total	O	0	0
			277	277		
5	C	191	Total	O	0	0
			191	191		
5	D	235	Total	O	0	0
			235	235		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	204	Total 204	O 204	0	0
5	F	199	Total 199	O 199	0	0
5	G	230	Total 230	O 230	0	0
5	H	274	Total 274	O 274	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyranose 2-oxidase





• Molecule 1: Pyranose 2-oxidase






• Molecule 1: Pyranose 2-oxidase




• Molecule 1: Pyranose 2-oxidase




• Molecule 1: Pyranose 2-oxidase







- Molecule 1: Pyranose 2-oxidase




- Molecule 1: Pyranose 2-oxidase





• Molecule 1: Pyranose 2-oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	168.19Å 103.14Å 168.71Å 90.00° 106.47° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 30.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	97.4 (30.00-2.20) 97.4 (30.00-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5	Depositor
R, R_{free}	0.207 , 0.266 0.211 , 0.265	Depositor DCC
R_{free} test set	2798 reflections (1.00%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.022 for l,-k,h	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	38766	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MES, FAD, SHG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.32	7/4645 (0.2%)	1.21	22/6317 (0.3%)
1	B	1.28	16/4668 (0.3%)	1.22	21/6348 (0.3%)
1	C	1.15	5/4644 (0.1%)	1.15	18/6316 (0.3%)
1	D	1.22	10/4644 (0.2%)	1.18	17/6316 (0.3%)
1	E	1.16	7/4661 (0.2%)	1.18	20/6338 (0.3%)
1	F	1.18	3/4637 (0.1%)	1.19	25/6306 (0.4%)
1	G	1.23	4/4637 (0.1%)	1.18	16/6306 (0.3%)
1	H	1.27	11/4637 (0.2%)	1.17	11/6306 (0.2%)
All	All	1.23	63/37173 (0.2%)	1.18	150/50553 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	H	0	1
All	All	0	2

The worst 5 of 63 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	508	ALA	CA-CB	8.63	1.67	1.53
1	D	349	LEU	CA-C	-7.88	1.48	1.53
1	H	422	ASP	C-O	-7.18	1.15	1.24
1	H	167	HIS	C-O	-7.02	1.15	1.24
1	H	589	ALA	CA-CB	-6.74	1.44	1.53

The worst 5 of 150 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	430	ASP	CA-C-N	-10.97	109.67	120.21
1	B	430	ASP	C-N-CA	-10.97	109.67	120.21
1	F	424	LEU	CA-C-N	10.18	131.03	120.04
1	F	424	LEU	C-N-CA	10.18	131.03	120.04
1	A	341	ASN	CA-C-N	9.26	128.87	119.24

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	449	ILE	Peptide
1	H	449	ILE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4528	0	4376	96	0
1	B	4551	0	4396	88	0
1	C	4527	0	4372	97	0
1	D	4527	0	4372	82	0
1	E	4544	0	4389	112	0
1	F	4520	0	4365	96	0
1	G	4520	0	4365	105	0
1	H	4520	0	4365	107	0
2	A	53	0	28	2	0
2	B	53	0	29	5	0
2	C	53	0	26	5	0
2	D	53	0	27	6	0
2	E	53	0	27	9	0
2	F	53	0	26	1	0
2	G	53	0	27	3	0
2	H	53	0	26	3	0
3	A	12	0	10	2	0
3	B	12	0	10	3	0
3	C	12	0	11	5	0
3	D	12	0	9	4	0
3	E	12	0	10	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	12	0	11	1	0
3	G	12	0	10	3	0
3	H	12	0	10	1	0
4	A	12	0	12	4	0
4	B	12	0	12	7	0
4	C	24	0	24	21	0
4	E	12	0	12	12	0
4	F	12	0	12	1	0
4	H	24	0	24	23	0
5	A	303	0	0	9	0
5	B	277	0	0	10	0
5	C	191	0	0	2	0
5	D	235	0	0	6	0
5	E	204	0	0	4	0
5	F	199	0	0	2	0
5	G	230	0	0	6	0
5	H	274	0	0	13	0
All	All	38766	0	35393	766	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 766 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:HIS:HE2	2:B:801:FAD:C8M	1.09	1.59
1:B:167:HIS:NE2	2:B:801:FAD:HM82	1.06	1.38
1:C:133:ALA:CB	4:C:902:MES:H71	1.58	1.32
1:G:82:SER:C	1:H:81:ASP:HA	1.67	1.17
1:E:490:LYS:NZ	1:E:490:LYS:HB3	1.50	1.16

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	572/623 (92%)	548 (96%)	24 (4%)	0	100	100
1	B	575/623 (92%)	544 (95%)	31 (5%)	0	100	100
1	C	572/623 (92%)	545 (95%)	26 (4%)	1 (0%)	43	51
1	D	572/623 (92%)	548 (96%)	24 (4%)	0	100	100
1	E	574/623 (92%)	547 (95%)	26 (4%)	1 (0%)	43	51
1	F	571/623 (92%)	539 (94%)	31 (5%)	1 (0%)	43	51
1	G	571/623 (92%)	545 (95%)	26 (5%)	0	100	100
1	H	571/623 (92%)	555 (97%)	15 (3%)	1 (0%)	43	51
All	All	4578/4984 (92%)	4371 (96%)	203 (4%)	4 (0%)	48	57

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	343	ALA
1	H	187	ASP
1	C	81	ASP
1	F	344	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	502/542 (93%)	476 (95%)	26 (5%)	21	26
1	B	505/542 (93%)	490 (97%)	15 (3%)	36	49
1	C	502/542 (93%)	487 (97%)	15 (3%)	36	49
1	D	502/542 (93%)	482 (96%)	20 (4%)	28	38
1	E	504/542 (93%)	484 (96%)	20 (4%)	28	38
1	F	501/542 (92%)	483 (96%)	18 (4%)	31	42
1	G	501/542 (92%)	484 (97%)	17 (3%)	32	44

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	501/542 (92%)	480 (96%)	21 (4%)	26	36
All	All	4018/4336 (93%)	3866 (96%)	152 (4%)	29	40

5 of 152 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	82	SER
1	H	389	LEU
1	G	185	LYS
1	G	593	ASN
1	H	614	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 71 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	344	ASN
1	G	440	GLN
1	H	276	ASN
1	D	132	GLN
1	D	108	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SHG	E	901	-	12,12,12	1.30	1 (8%)	16,17,17	5.32	9 (56%)
4	MES	E	902	-	12,12,12	2.14	1 (8%)	15,16,16	6.03	10 (66%)
2	FAD	A	801	1	58,58,58	1.36	8 (13%)	85,89,89	2.96	36 (42%)
3	SHG	H	901	-	12,12,12	1.02	1 (8%)	16,17,17	4.89	10 (62%)
2	FAD	C	801	1	58,58,58	1.49	11 (18%)	85,89,89	2.75	36 (42%)
4	MES	B	902	-	12,12,12	1.99	1 (8%)	15,16,16	5.82	10 (66%)
4	MES	H	624	-	12,12,12	2.20	1 (8%)	15,16,16	6.57	9 (60%)
4	MES	A	902	-	12,12,12	2.00	1 (8%)	15,16,16	4.42	9 (60%)
3	SHG	F	901	-	12,12,12	0.86	0	16,17,17	3.33	10 (62%)
2	FAD	H	801	1	58,58,58	1.29	7 (12%)	85,89,89	3.65	38 (44%)
3	SHG	B	901	-	12,12,12	1.21	1 (8%)	16,17,17	4.79	9 (56%)
2	FAD	B	801	1	58,58,58	1.31	6 (10%)	85,89,89	2.62	30 (35%)
3	SHG	C	901	-	12,12,12	0.78	0	16,17,17	3.03	6 (37%)
3	SHG	G	901	-	12,12,12	0.97	1 (8%)	16,17,17	4.97	10 (62%)
4	MES	C	624	-	12,12,12	2.22	2 (16%)	15,16,16	5.96	8 (53%)
3	SHG	D	901	-	12,12,12	1.04	1 (8%)	16,17,17	4.94	8 (50%)
4	MES	H	902	-	12,12,12	2.00	1 (8%)	15,16,16	5.55	8 (53%)
2	FAD	G	801	1	58,58,58	1.35	6 (10%)	85,89,89	2.64	34 (40%)
2	FAD	E	801	1	58,58,58	1.18	5 (8%)	85,89,89	2.57	31 (36%)
3	SHG	A	901	-	12,12,12	0.65	0	16,17,17	3.75	10 (62%)
4	MES	F	902	-	12,12,12	1.58	1 (8%)	15,16,16	5.34	10 (66%)
2	FAD	D	801	1	58,58,58	1.37	6 (10%)	85,89,89	3.00	39 (45%)
2	FAD	F	801	1	58,58,58	1.42	8 (13%)	85,89,89	2.56	33 (38%)
4	MES	C	902	-	12,12,12	1.62	1 (8%)	15,16,16	5.63	9 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SHG	E	901	-	1/1/5/5	0/2/22/22	0/1/1/1
4	MES	E	902	-	-	4/6/14/14	0/1/1/1
2	FAD	A	801	1	1/1/9/9	2/34/50/50	0/6/6/6
3	SHG	H	901	-	1/1/5/5	2/2/22/22	0/1/1/1
2	FAD	C	801	1	2/2/9/9	2/34/50/50	0/6/6/6
4	MES	B	902	-	-	5/6/14/14	0/1/1/1
4	MES	H	624	-	-	4/6/14/14	0/1/1/1
4	MES	A	902	-	-	3/6/14/14	0/1/1/1
3	SHG	F	901	-	-	2/2/22/22	0/1/1/1
2	FAD	H	801	1	2/2/9/9	2/34/50/50	0/6/6/6
3	SHG	B	901	-	1/1/5/5	0/2/22/22	0/1/1/1
2	FAD	B	801	1	1/1/9/9	2/34/50/50	0/6/6/6
3	SHG	G	901	-	1/1/5/5	2/2/22/22	0/1/1/1
3	SHG	C	901	-	-	0/2/22/22	0/1/1/1
4	MES	C	624	-	-	4/6/14/14	0/1/1/1
3	SHG	D	901	-	1/1/5/5	1/2/22/22	0/1/1/1
4	MES	H	902	-	-	2/6/14/14	0/1/1/1
2	FAD	G	801	1	-	4/34/50/50	0/6/6/6
2	FAD	E	801	1	1/1/9/9	4/34/50/50	0/6/6/6
3	SHG	A	901	-	-	0/2/22/22	0/1/1/1
4	MES	F	902	-	-	2/6/14/14	0/1/1/1
2	FAD	D	801	1	-	2/34/50/50	0/6/6/6
2	FAD	F	801	1	2/2/9/9	5/34/50/50	0/6/6/6
4	MES	C	902	-	-	5/6/14/14	0/1/1/1

The worst 5 of 71 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	624	MES	C8-S	-7.24	1.67	1.77
4	C	624	MES	C8-S	-7.13	1.67	1.77
4	E	902	MES	C8-S	-6.95	1.67	1.77
4	A	902	MES	C8-S	-6.62	1.68	1.77
4	B	902	MES	C8-S	-6.51	1.68	1.77

The worst 5 of 422 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	624	MES	O1S-S-C8	-19.79	76.82	106.73
4	E	902	MES	O1S-S-C8	-15.15	83.83	106.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	F	902	MES	O3S-S-C8	-14.53	77.59	106.00
4	C	902	MES	O3S-S-C8	-14.08	78.46	106.00
4	B	902	MES	O1S-S-C8	-13.73	85.98	106.73

5 of 14 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	801	FAD	C4B
2	B	801	FAD	C4B
2	C	801	FAD	C1B
2	C	801	FAD	C2B
2	E	801	FAD	C4B

5 of 59 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	FAD	PA-O3P-P-O5'
2	D	801	FAD	PA-O3P-P-O5'
2	E	801	FAD	PA-O3P-P-O5'
2	F	801	FAD	PA-O3P-P-O5'
4	A	902	MES	C8-C7-N4-C5

There are no ring outliers.

24 monomers are involved in 118 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	901	SHG	6	0
4	E	902	MES	12	0
2	A	801	FAD	2	0
3	H	901	SHG	1	0
2	C	801	FAD	5	0
4	B	902	MES	7	0
4	H	624	MES	7	0
4	A	902	MES	4	0
3	F	901	SHG	1	0
2	H	801	FAD	3	0
3	B	901	SHG	3	0
2	B	801	FAD	5	0
3	C	901	SHG	5	0
3	G	901	SHG	3	0
4	C	624	MES	11	0
3	D	901	SHG	4	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	902	MES	16	0
2	G	801	FAD	3	0
2	E	801	FAD	9	0
3	A	901	SHG	2	0
4	F	902	MES	1	0
2	D	801	FAD	6	0
2	F	801	FAD	1	0
4	C	902	MES	10	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	574/623 (92%)	-0.25	6 (1%) 79 77	11, 18, 31, 52	0
1	B	577/623 (92%)	-0.16	7 (1%) 76 74	11, 19, 33, 45	0
1	C	574/623 (92%)	0.09	12 (2%) 63 60	13, 23, 39, 53	0
1	D	574/623 (92%)	-0.10	8 (1%) 73 71	12, 21, 34, 48	0
1	E	576/623 (92%)	0.14	15 (2%) 57 54	15, 24, 40, 54	0
1	F	573/623 (91%)	0.04	11 (1%) 66 63	13, 23, 36, 49	0
1	G	573/623 (91%)	-0.03	9 (1%) 70 67	14, 21, 36, 52	0
1	H	573/623 (91%)	-0.14	6 (1%) 79 77	12, 19, 32, 46	0
All	All	4594/4984 (92%)	-0.05	74 (1%) 70 67	11, 21, 36, 54	0

The worst 5 of 74 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	619	THR	4.6
1	E	45	ILE	4.3
1	C	619	THR	4.0
1	C	389	LEU	3.6
1	F	389	LEU	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MES	C	902	12/12	0.68	0.25	65,68,69,72	0
4	MES	C	624	12/12	0.69	0.34	99,101,101,101	0
4	MES	H	902	12/12	0.73	0.24	70,77,78,78	0
4	MES	E	902	12/12	0.76	0.25	76,78,79,81	0
4	MES	H	624	12/12	0.78	0.21	68,71,73,73	0
4	MES	B	902	12/12	0.79	0.22	70,73,76,78	0
3	SHG	A	901	12/12	0.89	0.12	32,35,40,43	0
3	SHG	E	901	12/12	0.90	0.10	43,46,48,50	0
3	SHG	F	901	12/12	0.90	0.10	28,31,35,40	0
4	MES	A	902	12/12	0.90	0.13	44,48,53,53	0
3	SHG	C	901	12/12	0.91	0.11	37,39,46,46	0
3	SHG	G	901	12/12	0.91	0.10	41,44,48,49	0
3	SHG	B	901	12/12	0.91	0.11	34,37,42,42	0
3	SHG	H	901	12/12	0.92	0.11	39,41,45,49	0
3	SHG	D	901	12/12	0.92	0.10	38,40,42,44	0
4	MES	F	902	12/12	0.93	0.11	38,40,41,43	0
2	FAD	C	801	53/53	0.95	0.08	20,30,35,41	0
2	FAD	E	801	53/53	0.96	0.07	21,28,34,36	0
2	FAD	B	801	53/53	0.97	0.07	16,26,31,33	0
2	FAD	F	801	53/53	0.97	0.06	13,27,36,41	0
2	FAD	G	801	53/53	0.97	0.06	16,24,28,30	0
2	FAD	H	801	53/53	0.97	0.06	17,25,31,32	0
2	FAD	A	801	53/53	0.97	0.06	16,21,26,27	0
2	FAD	D	801	53/53	0.98	0.05	16,25,29,34	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD C 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD E 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD B 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD F 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD G 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around FAD H 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD D 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.