



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 05:05 AM UTC

PDB ID : 4K4B / pdb_00004k4b
Title : X-ray crystal structure of E. coli YdiI complexed with undeca-2-one-CoA
Authors : Ru, W.; Farelli, J.D.; Dunaway-Mariano, D.; Allen, K.N.
Deposited on : 2013-04-12
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	% 90% 10%
1	B	136	83% 15% .
1	C	136	% 87% 13%
1	D	136	86% 12% .
1	E	136	% 88% 12%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	136	 88% 11% .
1	G	136	%  83% 14% .
1	H	136	 87% 11% .

2 Entry composition


There are 4 unique types of molecules in this entry. The entry contains 10531 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Esterase YdiI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	136	Total	C	N	O	S	0	9	0
			1116	694	207	206	9			
1	B	136	Total	C	N	O	S	0	8	0
			1103	685	205	203	10			
1	C	136	Total	C	N	O	S	0	10	0
			1125	699	211	205	10			
1	D	136	Total	C	N	O	S	0	10	0
			1122	699	206	208	9			
1	E	136	Total	C	N	O	S	0	9	0
			1112	692	206	205	9			
1	F	136	Total	C	N	O	S	0	7	0
			1098	685	202	203	8			
1	G	136	Total	C	N	O	S	0	6	0
			1084	677	199	199	9			
1	H	136	Total	C	N	O	S	0	8	0
			1114	690	207	208	9			

- Molecule 2 is undeca-2-one coenzyme A (CCD ID: UOQ) (formula: C₃₂H₅₆N₇O₁₇P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total 60	C 32	N 7	O 17	P 3	S 1	0	0
2	A	1	Total 60	C 32	N 7	O 17	P 3	S 1	2	0
2	B	1	Total 60	C 32	N 7	O 17	P 3	S 1	0	0
2	B	1	Total 60	C 32	N 7	O 17	P 3	S 1	1	0
2	C	1	Total 60	C 32	N 7	O 17	P 3	S 1	1	0
2	D	1	Total 60	C 32	N 7	O 17	P 3	S 1	3	0
2	E	1	Total 60	C 32	N 7	O 17	P 3	S 1	0	0
2	E	1	Total 60	C 32	N 7	O 17	P 3	S 1	1	0
2	E	1	Total 60	C 32	N 7	O 17	P 3	S 1	0	0
2	F	1	Total 60	C 32	N 7	O 17	P 3	S 1	0	0
2	H	1	Total 60	C 32	N 7	O 17	P 3	S 1	0	0
2	H	1	Total 60	C 32	N 7	O 17	P 3	S 1	3	0
2	H	1	Total 60	C 32	N 7	O 17	P 3	S 1	1	0

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Cl 1	0	0
3	E	1	Total 1	Cl 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	127	Total 127	O 127	0	0
4	B	114	Total 114	O 114	0	0
4	C	119	Total 119	O 119	0	0
4	D	100	Total 100	O 100	0	0
4	E	88	Total 88	O 88	0	0
4	F	89	Total 89	O 89	0	0
4	G	100	Total 100	O 100	0	0
4	H	138	Total 138	O 138	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Esterase YdiI




- Molecule 1: Esterase YdiI




- Molecule 1: Esterase YdiI




- Molecule 1: Esterase YdiI



- Molecule 1: Esterase YdiI





- Molecule 1: Esterase YdiI






• Molecule 1: Esterase YdiI



• Molecule 1: Esterase YdiI



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	57.97Å 68.58Å 81.14Å 78.93° 84.69° 76.46°	Depositor
Resolution (Å)	33.89 – 1.90 33.89 – 1.90	Depositor EDS
% Data completeness (in resolution range)	93.3 (33.89-1.90) 87.8 (33.89-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 1.89Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.169 , 0.216 0.168 , 0.215	Depositor DCC
R_{free} test set	4421 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	11.4	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10531	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.32 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.7081e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UOQ, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/1136	0.78	0/1528
1	B	0.44	0/1122	0.74	0/1507
1	C	0.45	0/1145	0.78	0/1539
1	D	0.45	0/1145	0.72	0/1540
1	E	0.44	0/1132	0.77	1/1523 (0.1%)
1	F	0.42	0/1118	0.74	0/1504
1	G	0.46	0/1106	0.77	1/1487 (0.1%)
1	H	0.47	0/1130	0.75	0/1518
All	All	0.45	0/9034	0.76	2/12146 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	105	LEU	N-CA-C	-5.15	107.54	113.88
1	E	105	LEU	N-CA-C	-5.00	107.73	113.88

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1116	0	1123	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1103	0	1119	20	0
1	C	1125	0	1140	18	0
1	D	1122	0	1130	16	0
1	E	1112	0	1121	12	0
1	F	1098	0	1112	13	0
1	G	1084	0	1105	18	0
1	H	1114	0	1115	21	0
2	A	120	0	104	22	0
2	B	120	0	104	5	0
2	C	60	0	52	8	0
2	D	60	0	52	7	0
2	E	180	0	156	18	0
2	F	60	0	52	10	0
2	H	180	0	155	9	0
3	A	1	0	0	0	0
3	E	1	0	0	0	0
4	A	127	0	0	11	0
4	B	114	0	0	6	0
4	C	119	0	0	8	0
4	D	100	0	0	6	0
4	E	88	0	0	9	0
4	F	89	0	0	4	0
4	G	100	0	0	5	0
4	H	138	0	0	10	0
All	All	10531	0	9640	168	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 168 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:201:UOQ:S1P	4:D:693:HOH:O	2.02	1.14
2:H:202:UOQ:O8A	2:H:202:UOQ:H5B	1.38	1.12
1:E:127[B]:CYS:SG	4:E:749:HOH:O	2.23	0.96
2:A:203:UOQ:S1P	4:B:405:HOH:O	2.24	0.94
1:B:121:GLU:OE2	4:B:404:HOH:O	1.84	0.93

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/136 (105%)	142 (99%)	1 (1%)	0	100	100
1	B	142/136 (104%)	142 (100%)	0	0	100	100
1	C	144/136 (106%)	143 (99%)	1 (1%)	0	100	100
1	D	144/136 (106%)	142 (99%)	2 (1%)	0	100	100
1	E	143/136 (105%)	140 (98%)	3 (2%)	0	100	100
1	F	141/136 (104%)	140 (99%)	1 (1%)	0	100	100
1	G	140/136 (103%)	138 (99%)	2 (1%)	0	100	100
1	H	142/136 (104%)	141 (99%)	1 (1%)	0	100	100
All	All	1139/1088 (105%)	1128 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/112 (108%)	118 (98%)	3 (2%)	42	37
1	B	120/112 (107%)	116 (97%)	4 (3%)	33	26
1	C	122/112 (109%)	119 (98%)	3 (2%)	42	37
1	D	122/112 (109%)	116 (95%)	6 (5%)	22	14
1	E	121/112 (108%)	118 (98%)	3 (2%)	42	37
1	F	119/112 (106%)	115 (97%)	4 (3%)	32	25

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	118/112 (105%)	112 (95%)	6 (5%)	21	13
1	H	120/112 (107%)	116 (97%)	4 (3%)	33	26
All	All	963/896 (108%)	930 (97%)	33 (3%)	34	25

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	92	SER
1	H	48	GLN
1	H	130	ARG
1	D	48	GLN
1	D	31[B]	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	19	ASN
1	E	19	ASN
1	F	19	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UOQ	A	203	-	60,62,62	2.07	16 (26%)	83,88,88	2.11	19 (22%)
2	UOQ	D	501	-	60,62,62	2.07	17 (28%)	83,88,88	2.11	22 (26%)
2	UOQ	E	201	-	60,62,62	2.10	15 (25%)	83,88,88	2.35	22 (26%)
2	UOQ	H	203	-	60,62,62	2.03	14 (23%)	83,88,88	2.15	20 (24%)
2	UOQ	E	204	-	60,62,62	2.02	16 (26%)	83,88,88	2.15	21 (25%)
2	UOQ	H	201	-	60,62,62	2.07	14 (23%)	83,88,88	2.15	20 (24%)
2	UOQ	C	201	-	60,62,62	2.02	16 (26%)	83,88,88	2.13	18 (21%)
2	UOQ	A	201	-	60,62,62	2.19	16 (26%)	83,88,88	2.95	29 (34%)
2	UOQ	H	202	-	60,62,62	3.21	17 (28%)	83,88,88	2.24	25 (30%)
2	UOQ	B	202	-	60,62,62	2.00	16 (26%)	83,88,88	2.11	22 (26%)
2	UOQ	F	201	-	60,62,62	2.06	16 (26%)	83,88,88	2.21	22 (26%)
2	UOQ	E	203	-	60,62,62	2.09	16 (26%)	83,88,88	2.17	17 (20%)
2	UOQ	B	201	-	60,62,62	2.08	15 (25%)	83,88,88	1.95	15 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UOQ	A	203	-	-	21/61/77/77	0/3/3/3
2	UOQ	D	501	-	-	26/61/77/77	0/3/3/3
2	UOQ	E	201	-	-	33/61/77/77	0/3/3/3
2	UOQ	H	203	-	-	11/61/77/77	0/3/3/3
2	UOQ	E	204	-	-	21/61/77/77	0/3/3/3
2	UOQ	H	201	-	-	21/61/77/77	0/3/3/3
2	UOQ	C	201	-	-	20/61/77/77	0/3/3/3
2	UOQ	A	201	-	-	17/61/77/77	0/3/3/3
2	UOQ	H	202	-	-	19/61/77/77	0/3/3/3
2	UOQ	B	202	-	-	16/61/77/77	0/3/3/3
2	UOQ	F	201	-	-	14/61/77/77	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UOQ	E	203	-	-	17/61/77/77	0/3/3/3
2	UOQ	B	201	-	-	14/61/77/77	0/3/3/3

The worst 5 of 204 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	202	UOQ	P3B-O3B	-17.16	1.29	1.59
2	H	202	UOQ	O3B-C3B	-8.69	1.14	1.44
2	F	201	UOQ	OCH-CBX	8.56	1.36	1.21
2	E	201	UOQ	OCH-CBX	8.54	1.36	1.21
2	D	501	UOQ	OCH-CBX	8.46	1.36	1.21

The worst 5 of 272 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	UOQ	C4A-N9A-C1B	-10.58	101.88	126.63
2	A	201	UOQ	C1B-N9A-C8A	10.03	149.36	127.09
2	A	201	UOQ	OCH-CBX-CBW	-9.93	108.01	122.17
2	E	204	UOQ	OCH-CBX-CBW	-9.57	108.52	122.17
2	E	203	UOQ	OCH-CBX-CBW	-9.12	109.17	122.17

There are no chirality outliers.

5 of 250 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	UOQ	C5B-O5B-P1A-O2A
2	A	201	UOQ	S1P-C2P-C3P-N4P
2	A	201	UOQ	C4B-C5B-O5B-P1A
2	A	201	UOQ	N8P-C9P-CAP-OAP
2	A	201	UOQ	O9P-C9P-CAP-OAP

There are no ring outliers.

13 monomers are involved in 79 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	203	UOQ	12	0
2	D	501	UOQ	7	0
2	E	201	UOQ	6	0
2	H	203	UOQ	4	0
2	E	204	UOQ	6	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	201	UOQ	1	0
2	C	201	UOQ	8	0
2	A	201	UOQ	10	0
2	H	202	UOQ	4	0
2	B	202	UOQ	3	0
2	F	201	UOQ	10	0
2	E	203	UOQ	6	0
2	B	201	UOQ	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



































5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	136/136 (100%)	-0.45	1 (0%) 84 86	4, 10, 27, 40	9 (6%)
1	B	136/136 (100%)	-0.37	0 100 100	4, 13, 34, 47	8 (5%)
1	C	136/136 (100%)	-0.33	1 (0%) 84 86	4, 11, 28, 44	10 (7%)
1	D	136/136 (100%)	-0.38	0 100 100	4, 11, 31, 35	10 (7%)
1	E	136/136 (100%)	-0.19	2 (1%) 72 75	5, 14, 36, 48	9 (6%)
1	F	136/136 (100%)	-0.32	0 100 100	5, 14, 34, 47	7 (5%)
1	G	136/136 (100%)	-0.28	2 (1%) 72 75	5, 13, 30, 42	6 (4%)
1	H	136/136 (100%)	-0.44	0 100 100	4, 10, 27, 34	8 (5%)
All	All	1088/1088 (100%)	-0.34	6 (0%) 85 87	4, 12, 31, 48	67 (6%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	136	LEU	2.7
1	E	33	ASP	2.3
1	E	7	ILE	2.3
1	G	136	LEU	2.3
1	G	7[A]	ILE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	UOQ	E	201	60/60	0.81	0.18	16,47,65,76	24
2	UOQ	D	501	60/60	0.82	0.21	6,43,61,73	28
2	UOQ	A	201	60/60	0.82	0.20	15,46,80,89	24
2	UOQ	E	203	60/60	0.84	0.18	10,48,72,83	22
2	UOQ	H	202	60/60	0.85	0.18	7,43,65,76	25
2	UOQ	H	201	60/60	0.86	0.18	11,45,79,81	23
2	UOQ	C	201	60/60	0.86	0.18	9,49,77,90	21
2	UOQ	A	203	60/60	0.88	0.17	11,41,79,81	21
2	UOQ	F	201	60/60	0.88	0.17	15,33,71,79	24
2	UOQ	E	204	60/60	0.89	0.16	10,37,97,97	19
2	UOQ	B	201	60/60	0.90	0.15	15,38,80,88	23
2	UOQ	B	202	60/60	0.93	0.15	8,26,69,84	21
2	UOQ	H	203	60/60	0.94	0.13	11,27,70,76	21
3	CL	E	202	1/1	0.97	0.12	27,27,27,27	0
3	CL	A	202	1/1	0.98	0.04	25,25,25,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UOQ E 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UOQ D 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around UOQ A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UOQ E 203:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UOQ H 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UOQ H 201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UOQ C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around UOQ A 203:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UOQ F 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UOQ E 204:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UOQ B 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UOQ B 202:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UOQ H 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.