



Full wwPDB EM Validation Report ⓘ

Mar 26, 2026 – 10:34 AM UTC

PDB ID : 9K41 / pdb_00009k41
EMDB ID : EMD-62038
Title : Cryo-EM structure of Arabidopsis thaliana H2A.W-nucleosome with Arabidopsis native 147bp DNA 15.2.2 (C2 symmetry)
Authors : Wang, Y.; Dong, A.
Deposited on : 2024-10-21
Resolution : 2.81 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	11740 (2.31 - 3.31)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	136	 65% 5% 29%
1	E	136	 63% 7% 29%
2	B	103	 71% 7% 22%
2	F	103	 73% 5% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	150	
3	G	150	
4	D	148	
4	H	148	
5	I	147	
6	J	147	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11828 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	96	Total	C	N	O	S	0	0
			783	497	149	135	2		
1	E	96	Total	C	N	O	S	0	0
			783	497	149	135	2		

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	80	Total	C	N	O	S	0	0
			644	406	127	110	1		
2	F	80	Total	C	N	O	S	0	0
			644	406	127	110	1		

- Molecule 3 is a protein called Probable histone H2A.7.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	104	Total	C	N	O	S	0	0
			785	501	146	136	2		
3	G	104	Total	C	N	O	S	0	0
			785	501	146	136	2		

- Molecule 4 is a protein called Histone H2B.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	93	Total	C	N	O	S	0	0
			731	468	125	136	2		
4	H	93	Total	C	N	O	S	0	0
			731	468	125	136	2		

- Molecule 5 is a DNA chain called 15.2.2 DNA (147-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	145	Total	C	N	O	P	0	0
			2950	1413	522	870	145		


- Molecule 6 is a DNA chain called 15.2.2 DNA (147-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	145	Total	C	N	O	P	0	0
			2992	1428	552	868	144		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Histone H3.1




• Molecule 1: Histone H3.1




• Molecule 2: Histone H4



• Molecule 2: Histone H4



• Molecule 3: Probable histone H2A.7



LYS
PRO
ALA
GLU
GLU
LYS
LYS
ALA
THR
LYS
SER
PRO
VAL
GLY
LYS
SER
LYS
PRO
LYS
LYS
ALA

• Molecule 3: Probable histone H2A.7

Chain G:  61% 9% 31%

MET GLU SER THR GLY LYS VAL LYS LYS PHE ALA GLY ARG LYS PRO PRO GLY ALA PRO LYS THR LYS LYS S23 V24 S25 R38 R41 M60 V63 A68 S85 R86 L93 E101 V109 T110 T111 P126 LYS LYS SER LYS ALA THR LYS PRO ALA GLU LYS

ALA
THR
LYS
SER
PRO
VAL
LYS
SER
PRO
LYS
LYS
ALA

• Molecule 4: Histone H2B.1

Chain D:  55% 7% 37%

MET ALA PRO ARG ALA GLU LYS LYS PRO ALA ALA LYS LYS THR ALA ALA GLU ARG PRO VAL GLU GLU ASN LYS ALA ALA GLU LYS ALA ALA PRO PRO GLU LYS LYS GLY LYS LEU PRO PRO LYS LYS GLU ALA GLY ASP LYS LYS LYS LYS LYS ARG SER LYS K55 K56 K61

I64 D74 M82 S100 N107 R115 V121 L129 S147

• Molecule 4: Histone H2B.1

Chain H:  57% 6% 37%

MET ALA PRO ARG ALA GLU LYS LYS PRO ALA ALA LYS LYS THR ALA ALA GLU ARG PRO VAL GLU GLU ASN LYS ALA ALA GLU LYS ALA ALA PRO PRO GLU LYS LYS GLY LYS LEU PRO PRO LYS LYS GLU ALA GLY ASP LYS LYS LYS LYS LYS ARG SER LYS K55 E58 T59

I89 N107 P110 R115 V121 R122 L123 K139 S147

• Molecule 5: 15.2.2 DNA (147-MER)

Chain I:  63% 36% .

DA C-72 T-68 A-67 C-62 T-61 T-57 A-56 A-55 T-54 C-49 A-48 A-47 T-46 C-36 G-35 A-34 T-33 T-25 C-24 A-23 C-15 A-14 T3 A4 G7 T8 A9 C14 C15 G16 C17 T25 G26 C27 T28 C34 C35 A36 T45 T46 T49 C50 T51 A54

A55 T56 A57 C64 A65 A66 G67 A69 T70 T71 A72 DA

• Molecule 6: 15.2.2 DNA (147-MER)

Chain J:  51% 48% .

DT T-72 C-67 T-66 T-65 G-64 T-63 G-62 T-59 T-58 T-57 A-56 T-55 T-54 A-53 G-48 A-47 A-46 A-45 A-44 G-43 C-39 G-38 G-37 T-36 G-35 G-34 C-24 T-18 G-17 T-9 A-8 C-7 G-6 A-5 T-4 A-3 C3 T4 T14 G15 A16 A17 G18 T19 T20 A21 T22 T23 G24 A25

C26	A27	G28	T29	T34	C35	G36	A37	G38	C39	C40	G41	A45	A46	T47	T48	G49	G50	T51	A54	T55	T58	G59	G60	A69	A70	G71	G72	DT
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	119108	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	900	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	1.294	Depositor
Minimum map value	-0.357	Depositor
Average map value	-0.002	Depositor
Map value standard deviation	0.052	Depositor
Recommended contour level	0.2	Depositor
Map size ($\text{\AA}$)	274.432, 274.432, 274.432	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.072, 1.072, 1.072	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.16	0/795	0.21	0/1067
1	E	0.16	0/795	0.22	0/1067
2	B	0.18	0/651	0.33	0/872
2	F	0.19	0/651	0.37	0/872
3	C	0.15	0/796	0.24	0/1072
3	G	0.15	0/796	0.25	0/1072
4	D	0.15	0/742	0.26	0/995
4	H	0.14	0/742	0.23	0/995
5	I	0.20	0/3304	0.43	0/5091
6	J	0.19	0/3361	0.38	0/5192
All	All	0.18	0/12633	0.35	0/18295

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	783	0	822	6	0
1	E	783	0	822	7	0
2	B	644	0	686	6	0
2	F	644	0	686	7	0
3	C	785	0	841	13	0
3	G	785	0	841	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	731	0	771	9	0
4	H	731	0	771	10	0
5	I	2950	0	1639	38	0
6	J	2992	0	1640	51	0
All	All	11828	0	9519	129	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (129) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:55:LYS:NZ	5:I:-46:DT:OP1	2.21	0.71
3:G:41:ARG:NH2	4:H:58:GLU:OE1	2.27	0.67
1:E:106:ASP:OD2	1:E:131:ARG:NH1	2.31	0.64
1:A:106:ASP:OD2	1:A:131:ARG:NH1	2.31	0.64
5:I:-36:DC:H2''	5:I:-35:DG:C8	2.33	0.64
3:C:35:PRO:HB2	3:C:38:ARG:HB3	1.81	0.63
1:A:68:GLN:HE21	1:A:72:ARG:HH21	1.46	0.62
2:B:75:HIS:O	4:D:115:ARG:NH2	2.33	0.61
4:D:61:LYS:HA	4:D:82:MET:HE2	1.83	0.60
2:F:75:HIS:CD2	4:H:107:ASN:HD21	2.19	0.60
5:I:-48:DA:H2''	5:I:-47:DA:C8	2.37	0.60
3:G:38:ARG:NH1	4:H:59:THR:O	2.35	0.60
5:I:54:DA:H2''	5:I:55:DA:C8	2.38	0.59
6:J:-48:DG:H2''	6:J:-47:DG:C8	2.38	0.58
3:C:101:GLU:OE1	4:D:129:LEU:N	2.36	0.58
5:I:-47:DA:H2''	5:I:-46:DT:H71	1.86	0.57
6:J:54:DA:H2''	6:J:55:DT:C5	2.39	0.57
2:F:47:SER:HA	5:I:7:DG:H5'	1.87	0.57
3:G:68:ALA:HB2	4:H:89:ILE:HD11	1.86	0.56
5:I:8:DT:H2''	5:I:9:DA:C8	2.41	0.56
4:D:64:ILE:HB	4:D:82:MET:HE1	1.88	0.56
2:F:75:HIS:O	4:H:115:ARG:NH2	2.40	0.55
6:J:-4:DT:H2''	6:J:-3:DA:C8	2.42	0.54
6:J:37:DA:H2''	6:J:38:DG:C8	2.42	0.54
2:B:75:HIS:CD2	4:D:107:ASN:HD21	2.24	0.54
1:E:85:GLN:HG3	6:J:-24:DC:OP1	2.07	0.53
5:I:45:DT:H2''	5:I:46:DT:H71	1.90	0.53
3:C:63:VAL:HG21	4:D:121:VAL:HG21	1.91	0.53
5:I:68:DC:H2''	5:I:69:DA:C8	2.43	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:I:16:DG:H1'	5:I:17:DC:C5	2.44	0.52
3:C:82:ASN:O	3:C:82:ASN:ND2	2.43	0.52
3:C:53:GLY:HA2	6:J:38:DG:H5''	1.91	0.51
3:C:124:LEU:HD11	1:E:108:ASN:HD21	1.76	0.51
6:J:-7:DC:H2''	6:J:-6:DG:N7	2.26	0.51
1:E:42:ARG:HB2	6:J:70:DA:OP1	2.11	0.50
6:J:48:DT:H2''	6:J:49:DG:C8	2.47	0.50
5:I:3:DT:H2''	5:I:4:DA:C8	2.47	0.50
6:J:-54:DT:H2''	6:J:-53:DA:C8	2.47	0.50
6:J:34:DT:H2''	6:J:35:DC:C5	2.47	0.50
6:J:22:DT:H1'	6:J:23:DT:H5'	1.93	0.50
1:A:68:GLN:NE2	1:A:72:ARG:HH21	2.09	0.50
3:G:86:ARG:NH1	6:J:-54:DT:H5'	2.27	0.50
6:J:14:DT:H2''	6:J:15:DG:C8	2.46	0.50
5:I:27:DC:H2''	5:I:28:DT:H71	1.93	0.49
3:G:38:ARG:NH2	5:I:49:DT:OP1	2.45	0.49
5:I:-57:DT:H2''	5:I:-56:DA:N7	2.27	0.49
5:I:56:DT:H1'	5:I:57:DA:C5	2.48	0.49
5:I:55:DA:H2''	5:I:56:DT:C7	2.43	0.49
5:I:14:DC:H2''	5:I:15:DC:C5	2.48	0.48
6:J:-67:DC:H2''	6:J:-66:DT:C6	2.48	0.48
5:I:-57:DT:H2''	5:I:-56:DA:C8	2.48	0.48
3:G:63:VAL:HG21	4:H:121:VAL:HG21	1.95	0.48
1:A:42:ARG:HB2	5:I:70:DT:OP1	2.13	0.48
5:I:34:DC:H2''	5:I:35:DC:C6	2.49	0.47
5:I:-47:DA:H2''	5:I:-46:DT:C7	2.44	0.47
6:J:24:DG:H2''	6:J:25:DA:C8	2.49	0.47
4:D:56:ASN:O	4:D:56:ASN:ND2	2.48	0.47
6:J:18:DG:H2'	6:J:19:DT:H71	1.96	0.46
3:G:93:LEU:HD23	3:G:111:ILE:HD12	1.97	0.46
6:J:-39:DC:H2''	6:J:-38:DG:C8	2.50	0.46
6:J:-63:DT:C2	6:J:-62:DG:N7	2.83	0.46
6:J:-57:DT:H2''	6:J:-56:DA:C8	2.50	0.46
6:J:-54:DT:H2''	6:J:-53:DA:H8	1.80	0.46
6:J:19:DT:H2'	6:J:20:DT:H71	1.96	0.46
2:B:98:TYR:CE2	3:G:109:VAL:HG11	2.51	0.45
5:I:64:DC:H2''	5:I:65:DA:C8	2.52	0.45
5:I:25:DT:H2''	5:I:26:DG:C8	2.52	0.45
6:J:17:DA:H2''	6:J:18:DG:H8	1.82	0.45
6:J:26:DC:H2''	6:J:27:DA:C8	2.51	0.45
5:I:7:DG:C8	5:I:8:DT:H72	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:52:LEU:N	6:J:39:DC:OP1	2.34	0.44
3:G:85:SER:O	3:G:85:SER:OG	2.30	0.44
6:J:-18:DT:H2''	6:J:-17:DG:C8	2.51	0.44
6:J:41:DG:H5'	6:J:41:DG:C8	2.53	0.44
5:I:-15:DC:H2''	5:I:-14:DA:C8	2.53	0.44
2:B:92:ARG:NH1	2:B:92:ARG:HB3	2.33	0.43
2:F:71:THR:HG21	4:H:123:LEU:HG	1.99	0.43
5:I:66:DA:H2''	5:I:67:DG:C8	2.53	0.43
6:J:-65:DT:H2''	6:J:-64:DG:H8	1.84	0.43
5:I:-68:DT:H2''	5:I:-67:DA:C8	2.54	0.43
6:J:-59:DT:H2''	6:J:-58:DT:C6	2.54	0.43
6:J:45:DA:H2''	6:J:46:DA:N7	2.34	0.43
5:I:-25:DT:H2''	5:I:-24:DC:C5	2.54	0.43
5:I:27:DC:H2''	5:I:28:DT:C7	2.49	0.43
5:I:45:DT:H2''	5:I:46:DT:C7	2.49	0.43
5:I:-49:DC:H2''	5:I:-48:DA:C8	2.54	0.42
3:G:60:MET:HE2	3:G:60:MET:HB2	1.94	0.42
6:J:59:DG:H2''	6:J:60:DG:C8	2.54	0.42
6:J:40:DC:C2	6:J:41:DG:N7	2.87	0.42
1:E:61:LEU:HD12	2:F:37:LEU:HD23	2.02	0.42
5:I:-34:DA:H2''	5:I:-33:DT:H71	2.02	0.42
6:J:39:DC:C2	6:J:40:DC:C5	3.08	0.42
6:J:50:DG:H2'	6:J:51:DT:H71	2.01	0.42
1:A:72:ARG:NH2	5:I:-23:DA:OP2	2.53	0.42
2:F:75:HIS:HD2	4:H:107:ASN:HD21	1.67	0.42
5:I:-55:DA:H2''	5:I:-54:DT:C7	2.49	0.42
5:I:35:DC:H2''	5:I:36:DA:C8	2.55	0.42
6:J:54:DA:H2''	6:J:55:DT:C6	2.55	0.42
1:E:63:ARG:HB2	1:E:66:PRO:HD2	2.00	0.42
6:J:16:DA:H2''	6:J:17:DA:C8	2.54	0.42
6:J:69:DA:H2''	6:J:70:DA:C8	2.55	0.42
4:H:110:PRO:HD2	6:J:-34:DG:OP1	2.20	0.41
3:C:52:LEU:O	6:J:38:DG:H3'	2.21	0.41
3:G:101:GLU:H	3:G:101:GLU:HG2	1.68	0.41
5:I:-62:DC:H2'	5:I:-61:DT:H71	2.02	0.41
6:J:-65:DT:H2''	6:J:-64:DG:C8	2.54	0.41
6:J:3:DC:H2''	6:J:4:DT:C5	2.55	0.41
6:J:28:DG:H1'	6:J:29:DT:H5'	2.00	0.41
1:A:63:ARG:HB2	1:A:66:PRO:HD2	2.02	0.41
6:J:-45:DA:H2''	6:J:-44:DA:C8	2.56	0.41
6:J:-36:DT:H2''	6:J:-35:DG:C8	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:75:HIS:HD2	4:D:107:ASN:HD21	1.67	0.41
6:J:-46:DA:H2''	6:J:-45:DA:C8	2.56	0.41
3:C:88:ILE:HG12	3:C:91:HIS:CE1	2.56	0.41
6:J:-9:DT:H2''	6:J:-8:DA:C8	2.56	0.41
3:C:92:LEU:O	3:C:96:ILE:HG13	2.20	0.41
3:G:25:SER:HA	6:J:-43:DG:H5''	2.02	0.41
5:I:50:DC:H2'	5:I:51:DT:H71	2.02	0.41
6:J:58:DT:H2''	6:J:59:DG:C8	2.56	0.41
3:C:122:SER:HA	3:C:125:LEU:HD12	2.03	0.40
1:E:62:ILE:HD11	2:F:37:LEU:HD11	2.04	0.40
5:I:49:DT:H1'	5:I:50:DC:OP2	2.21	0.40
3:C:60:MET:HE2	3:C:60:MET:HB2	1.94	0.40
2:B:96:THR:HB	3:G:109:VAL:HG22	2.04	0.40
3:C:74:LEU:HD12	3:C:102:LEU:HD12	2.01	0.40
6:J:41:DG:H5'	6:J:41:DG:H8	1.87	0.40
4:H:139:LYS:HB3	4:H:139:LYS:HE2	1.89	0.40
6:J:-37:DG:H2''	6:J:-36:DT:C5	2.57	0.40
6:J:59:DG:H2''	6:J:60:DG:H8	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/136 (69%)	92 (98%)	2 (2%)	0	100	100
1	E	94/136 (69%)	92 (98%)	2 (2%)	0	100	100
2	B	78/103 (76%)	77 (99%)	1 (1%)	0	100	100
2	F	78/103 (76%)	76 (97%)	2 (3%)	0	100	100
3	C	102/150 (68%)	101 (99%)	1 (1%)	0	100	100
3	G	102/150 (68%)	100 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	D	91/148 (62%)	90 (99%)	1 (1%)	0	100	100
4	H	91/148 (62%)	90 (99%)	1 (1%)	0	100	100
All	All	730/1074 (68%)	718 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/109 (75%)	82 (100%)	0	100	100
1	E	82/109 (75%)	81 (99%)	1 (1%)	63	86
2	B	66/79 (84%)	63 (96%)	3 (4%)	24	57
2	F	66/79 (84%)	65 (98%)	1 (2%)	57	83
3	C	81/117 (69%)	81 (100%)	0	100	100
3	G	81/117 (69%)	80 (99%)	1 (1%)	63	86
4	D	82/124 (66%)	80 (98%)	2 (2%)	43	75
4	H	82/124 (66%)	82 (100%)	0	100	100
All	All	622/858 (72%)	614 (99%)	8 (1%)	59	85

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	22	LEU
2	B	23	ARG
2	B	24	ASP
4	D	74	ASP
4	D	100	SER
1	E	120	MET
2	F	23	ARG
3	G	24	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	GLN
1	A	68	GLN
1	A	108	ASN
2	B	25	ASN
2	B	27	GLN
2	B	75	HIS
3	C	33	GLN
3	C	82	ASN
4	D	107	ASN
1	E	68	GLN
1	E	108	ASN
2	F	75	HIS
3	G	33	GLN
3	G	77	ASN
4	H	107	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-62038. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128




Y Index: 128




Z Index: 128


6.2.2 Raw map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 106



Y Index: 114




Z Index: 133


6.3.2 Raw map



X Index: 106



Y Index: 142




Z Index: 133


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.2. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_62038_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 186 nm^3 ; this corresponds to an approximate mass of 168 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.356 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.356 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.81	-	-
Author-provided FSC curve	2.81	3.24	2.86
Unmasked-calculated*	3.82	5.72	3.88

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.82 differs from the reported value 2.81 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-62038 and PDB model 9K41. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.2 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.2).

9.4 Atom inclusion ⓘ



At the recommended contour level, 100% of all backbone atoms, 99% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9880	0.4790
A	0.9860	0.5450
B	0.9740	0.5370
C	0.9780	0.5310
D	0.9830	0.5240
E	0.9830	0.5470
F	0.9790	0.5370
G	0.9800	0.5410
H	0.9830	0.5250
I	0.9960	0.4280
J	0.9950	0.4180

1.0

0.0

<0.0