



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:40 AM UTC

PDB ID : 7K5A / pdb_00007k5a
Title : The external aldimine form of Salmonella typhimurium Tryptophan Synthase mutant beta-Q114A in complex with cesium ion at the metal coordination site.
Authors : Hilario, E.; Dunn, M.F.; Mueller, L.J.
Deposited on : 2020-09-16
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	8% 88% 7% 5%
2	B	397	6% 89% 10% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	DMS	A	303	-	-	X	-
7	EDO	B	402	-	-	X	-

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 5755 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan synthase alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	255	Total	C	N	O	S	0	1	0
			1924	1227	330	359	8			


- Molecule 2 is a protein called Tryptophan synthase beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	395	Total	C	N	O	S	0	13	0
			3095	1940	546	588	21			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	114	ALA	GLN	engineered mutation	UNP P0A2K1

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is SERINE (CCD ID: SER) (formula: $C_3H_7NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			7	3	1	3		

- Molecule 5 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Cl	0	0
			2	2		
6	B	2	Total	Cl	0	0
			2	2		

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is (E)-N-({3-hydroxy-2-methyl-5-[(phosphonoxy)methyl]pyridin-4-yl}methylidene)-L-serine (CCD ID: KOU) (formula: C₁₁H₁₅N₂O₈P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	B	1	Total	C	N	O	P	0	0
			22	11	2	8	1		

- Molecule 9 is CESIUM ION (CCD ID: CS) (formula: Cs) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	2	Total	Cs	0	2
			4	4		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	205	Total	O	0	4
			209	209		
10	B	441	Total	O	0	6
			447	447		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tryptophan synthase alpha chain



- Molecule 2: Tryptophan synthase beta chain



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	182.83Å 59.53Å 67.48Å 90.00° 94.88° 90.00°	Depositor
Resolution (Å)	39.29 – 1.50 39.29 – 1.50	Depositor EDS
% Data completeness (in resolution range)	97.8 (39.29-1.50) 97.8 (39.29-1.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.168 , 0.220 0.171 , 0.221	Depositor DCC
R_{free} test set	5713 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.542	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5755	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, DMS, PEG, CL, CS, KOU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/1963	0.97	0/2669
2	B	0.61	0/3153	0.98	3/4255 (0.1%)
All	All	0.59	0/5116	0.98	3/6924 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	47	ASP	CA-CB-CG	6.73	119.33	112.60
2	B	357	HIS	CA-CB-CG	-5.86	107.94	113.80
2	B	241	PHE	CA-CB-CG	5.41	119.21	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1924	0	1916	22	0
2	B	3095	0	3058	37	0
3	A	7	0	10	2	0
4	A	7	0	4	1	0
5	A	8	0	12	6	0
5	B	20	0	30	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	2	0	0	0	0
6	B	2	0	0	0	0
7	B	8	0	12	5	0
8	B	22	0	12	0	0
9	B	4	0	0	0	0
10	A	209	0	0	2	0
10	B	447	0	0	14	0
All	All	5755	0	5054	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:70[B]:ARG:HH11	2:B:70[B]:ARG:CG	1.59	1.11
2:B:70[B]:ARG:NH1	2:B:70[B]:ARG:HG2	1.43	1.10
2:B:70[B]:ARG:CG	2:B:70[B]:ARG:NH1	2.22	0.85
2:B:379[B]:ARG:NH1	10:B:501:HOH:O	2.11	0.83
2:B:136:ALA:HB1	10:B:772:HOH:O	1.79	0.82
1:A:112:ASP:OD1	1:A:146:HIS:HE1	1.63	0.81
2:B:70[B]:ARG:HH11	2:B:70[B]:ARG:HG2	0.68	0.75
1:A:112:ASP:OD2	1:A:145:ARG:NH2	2.20	0.75
1:A:107:PHE:CE2	3:A:301:PEG:H32	2.22	0.74
2:B:279:TYR:CE2	10:B:676:HOH:O	2.40	0.74
2:B:11[B]:GLU:O	2:B:11[B]:GLU:HG2	1.93	0.69
2:B:65:ILE:HD11	2:B:338:THR:HG22	1.75	0.68
2:B:341:ARG:HG3	10:B:756:HOH:O	1.94	0.65
2:B:165:THR:HG23	10:B:696:HOH:O	1.97	0.63
2:B:280:PHE:CE2	10:B:922:HOH:O	2.51	0.62
1:A:1:MET:O	4:A:302:SER:HB2	2.00	0.62
1:A:171:ARG:NH2	5:A:303:DMS:H23	2.16	0.61
2:B:137:LYS:HD2	10:B:558:HOH:O	2.00	0.61
5:A:303:DMS:H11	10:A:529:HOH:O	2.01	0.59
2:B:19:GLN:HE21	7:B:402:EDO:C1	2.16	0.59
2:B:216[A]:ILE:HG21	2:B:224:PRO:HD3	1.87	0.57
2:B:165:THR:HB	10:B:558:HOH:O	2.04	0.57
2:B:334[B]:GLU:HG2	10:B:556:HOH:O	2.06	0.55
5:B:408:DMS:H22	10:B:806:HOH:O	2.08	0.54
1:A:100:LEU:C	1:A:100:LEU:HD13	2.33	0.53
1:A:210:GLN:HE21	1:A:212:PHE:H	1.55	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:279:TYR:HE2	10:B:676:HOH:O	1.85	0.53
1:A:140:ARG:HD3	1:A:169:TYR:HB3	1.90	0.53
2:B:117:VAL:HG13	2:B:152:MET:HE1	1.91	0.53
2:B:19:GLN:HE21	7:B:402:EDO:H11	1.73	0.52
2:B:108:ALA:HB3	2:B:132:ILE:CD1	2.39	0.52
1:A:210:GLN:NE2	1:A:212:PHE:HB2	2.25	0.52
2:B:163:SER:OG	2:B:165:THR:HG22	2.11	0.51
1:A:204:HIS:O	5:A:303:DMS:C1	2.59	0.50
1:A:112:ASP:OD1	1:A:146:HIS:CE1	2.54	0.50
1:A:155:PRO:HA	1:A:177:LEU:HB3	1.95	0.48
2:B:115:HIS:CE1	2:B:189:GLY:HA2	2.48	0.48
2:B:167:LYS:O	2:B:170:CYS:HB2	2.14	0.48
2:B:165:THR:HG21	10:B:748:HOH:O	2.14	0.47
1:A:117:ARG:HA	1:A:117:ARG:HD2	1.71	0.46
2:B:56:PRO:HD2	5:B:404:DMS:H22	1.96	0.45
2:B:137:LYS:O	2:B:141:ARG:HG2	2.17	0.45
1:A:171:ARG:HH21	5:A:303:DMS:H23	1.79	0.44
1:A:171:ARG:HD3	5:A:303:DMS:O	2.18	0.44
1:A:162:LEU:HD22	7:B:402:EDO:H11	2.00	0.44
3:A:301:PEG:H42	2:B:275:ARG:NH2	2.33	0.44
2:B:65:ILE:CD1	2:B:338:THR:HG22	2.44	0.44
2:B:296:GLU:HA	10:B:726:HOH:O	2.18	0.44
1:A:204:HIS:O	5:A:303:DMS:H11	2.18	0.43
2:B:88:THR:HA	2:B:91:VAL:HG22	2.01	0.42
2:B:22:MET:HB2	7:B:402:EDO:H12	2.01	0.42
1:A:117:ARG:NH1	10:A:409:HOH:O	2.53	0.42
2:B:108:ALA:HB3	2:B:132:ILE:HD13	2.00	0.42
2:B:301[A]:SER:OG	2:B:350:GLU:HG3	2.20	0.41
1:A:210:GLN:HE21	1:A:212:PHE:CB	2.33	0.41
1:A:210:GLN:NE2	1:A:212:PHE:CB	2.84	0.41
2:B:106:ILE:HG21	2:B:187[B]:MET:HE2	2.02	0.41
1:A:21:PRO:HD2	1:A:47:ALA:O	2.21	0.41
2:B:300:ILE:HD12	10:B:631:HOH:O	2.20	0.41
2:B:23:PRO:HG3	7:B:402:EDO:H21	2.02	0.41
1:A:20:VAL:HG22	1:A:47:ALA:HB3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/268 (94%)	249 (99%)	3 (1%)	0	100	100
2	B	406/397 (102%)	400 (98%)	6 (2%)	0	100	100
All	All	658/665 (99%)	649 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/208 (95%)	195 (99%)	2 (1%)	68	45
2	B	320/310 (103%)	319 (100%)	1 (0%)	86	75
All	All	517/518 (100%)	514 (99%)	3 (1%)	78	62

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	ASP
1	A	194	HIS
2	B	396	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	ASN
1	A	146	HIS
1	A	210	GLN
2	B	19	GLN
2	B	160	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 8 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	DMS	B	405	-	3,3,3	0.29	0	3,3,3	0.26	0
4	SER	A	302	-	4,6,6	1.02	0	2,7,7	1.24	0
5	DMS	B	408	-	3,3,3	0.26	0	3,3,3	0.16	0
8	KOU	B	403	-	22,22,22	0.93	2 (9%)	27,31,31	1.18	4 (14%)
5	DMS	A	303	-	3,3,3	0.22	0	3,3,3	0.39	0
5	DMS	B	406	-	3,3,3	0.29	0	3,3,3	0.12	0
7	EDO	B	402	-	3,3,3	0.19	0	2,2,2	0.46	0
5	DMS	B	404	-	3,3,3	0.95	0	3,3,3	0.29	0
5	DMS	B	407	-	3,3,3	0.35	0	3,3,3	0.10	0
5	DMS	A	304	-	3,3,3	0.25	0	3,3,3	0.06	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PEG	A	301	-	6,6,6	0.16	0	5,5,5	0.06	0
7	EDO	B	401	-	3,3,3	0.09	0	2,2,2	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SER	A	302	-	-	2/6/6/6	-
3	PEG	A	301	-	-	1/4/4/4	-
8	KOU	B	403	-	-	6/17/17/17	0/1/1/1
7	EDO	B	402	-	-	0/1/1/1	-
7	EDO	B	401	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	403	KOU	C3-C2	2.31	1.43	1.41
8	B	403	KOU	OXT-C	-2.16	1.23	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	B	403	KOU	C4-C3-C2	-2.60	118.68	120.14
8	B	403	KOU	OG-CB-CA	-2.41	98.01	111.42
8	B	403	KOU	CA-N-C4A	2.33	120.23	116.77
8	B	403	KOU	C-CA-N	2.13	113.47	108.22

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	B	403	KOU	O-C-CA-CB
8	B	403	KOU	OXT-C-CA-CB
8	B	403	KOU	CB-CA-N-C4A
8	B	403	KOU	N-CA-CB-OG
8	B	403	KOU	C4-C4A-N-CA
7	B	401	EDO	O1-C1-C2-O2
8	B	403	KOU	C-CA-CB-OG
3	A	301	PEG	O2-C3-C4-O4

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	A	302	SER	N-CA-CB-OG
4	A	302	SER	O-C-CA-N

There are no ring outliers.

6 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	302	SER	1	0
5	B	408	DMS	1	0
5	A	303	DMS	6	0
7	B	402	EDO	5	0
5	B	404	DMS	1	0
3	A	301	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/268 (95%)	0.62	21 (8%) 17 19	14, 35, 54, 100	1 (0%)
2	B	395/397 (99%)	0.07	24 (6%) 27 29	8, 20, 51, 89	13 (3%)
All	All	650/665 (97%)	0.29	45 (6%) 23 25	8, 26, 53, 100	14 (2%)

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	212	PHE	6.1
2	B	139	VAL	4.3
2	B	164	ALA	4.2
1	A	193	LEU	3.9
2	B	135	GLY	3.9
2	B	159	VAL	3.8
2	B	146	VAL	3.7
2	B	147	PHE	3.7
2	B	385	PHE	3.4
1	A	191	LEU	3.4
2	B	171	ASN	3.2
2	B	162	GLY	3.2
1	A	158	ALA	3.1
1	A	58	LEU	3.1
1	A	194	HIS	2.9
2	B	158	PRO	2.8
2	B	62	CYS	2.8
2	B	111	GLY	2.8
1	A	214	ILE	2.8
1	A	192	PRO	2.7
2	B	143	SER	2.7
2	B	138	ASP	2.7
1	A	268	ALA	2.6
1	A	157	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	177	LEU	2.5
2	B	145	ASN	2.5
2	B	169	ALA	2.5
1	A	261	ALA	2.4
1	A	257	SER	2.4
2	B	141	ARG	2.3
1	A	247	SER	2.3
2	B	280	PHE	2.3
1	A	195	HIS	2.3
2	B	134	MET	2.3
1	A	241	ILE	2.2
1	A	217	PRO	2.2
2	B	384	ILE	2.2
1	A	213	GLY	2.2
1	A	252	LEU	2.1
2	B	306	PHE	2.1
2	B	142	GLN	2.1
2	B	173	ALA	2.1
1	A	156	PRO	2.1
1	A	220	VAL	2.0
2	B	174	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	DMS	A	304	4/4	0.82	0.15	62,76,81,85	0
4	SER	A	302	7/7	0.83	0.14	52,58,61,62	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	DMS	A	303	4/4	0.87	0.17	43,55,57,64	0
5	DMS	B	405	4/4	0.88	0.14	37,45,49,62	0
3	PEG	A	301	7/7	0.90	0.10	30,41,49,53	0
5	DMS	B	407	4/4	0.90	0.14	51,53,60,63	0
5	DMS	B	408	4/4	0.90	0.13	52,63,71,72	0
7	EDO	B	402	4/4	0.90	0.11	29,38,38,45	0
6	CL	A	306	1/1	0.92	0.15	72,72,72,72	0
7	EDO	B	401	4/4	0.93	0.09	34,44,46,48	0
5	DMS	B	404	4/4	0.93	0.17	26,29,31,31	0
6	CL	B	412	1/1	0.94	0.17	50,50,50,50	0
6	CL	A	305	1/1	0.94	0.08	52,52,52,52	0
5	DMS	B	406	4/4	0.94	0.12	48,49,49,53	0
8	KOU	B	403	22/22	0.98	0.06	16,21,33,45	0
6	CL	B	411	1/1	0.99	0.04	36,36,36,36	0
9	CS	B	410[A]	1/1	0.99	0.08	28,28,28,28	1
9	CS	B	410[B]	1/1	0.99	0.08	28,28,28,28	1
9	CS	B	409[A]	1/1	1.00	0.06	22,22,22,22	1
9	CS	B	409[B]	1/1	1.00	0.06	25,25,25,25	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around KOU B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CS B 410 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CS B 410 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around CS B 409 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CS B 409 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.