



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 09:06 AM UTC

PDB ID : 8K5R / pdb_00008k5r
Title : CDK9/cyclin T1 in complex with KB-0742
Authors : Zhou, M.; Li, H.; Gao, H.; Trotter, B.W.; Freeman, D.
Deposited on : 2023-07-24
Resolution : 3.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1029 (3.90-3.62)
Clashscore	190562	1061 (3.90-3.62)
Ramachandran outliers	187476	1014 (3.90-3.62)
Sidechain outliers	187428	1009 (3.90-3.62)
RSRZ outliers	180081	1028 (3.90-3.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	 67% 23% 7%
2	B	259	 73% 23% 4% 0%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 9.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	P	S	0	0	0
			2480	1587	428	449	1	15			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ASP	SER	engineered mutation	UNP P50750
A	8	ASN	VAL	engineered mutation	UNP P50750
A	44	ARG	LYS	engineered mutation	UNP P50750
A	138	PHE	TYR	engineered mutation	UNP P50750
A	280	ALA	LYS	engineered mutation	UNP P50750
A	307	GLU	ASP	engineered mutation	UNP P50750
A	311	GLU	ASN	engineered mutation	UNP P50750


- Molecule 2 is a protein called Cyclin-T1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	252	Total	C	N	O	S	0	0	0
			2057	1315	359	374	9			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	26	ALA	ARG	engineered mutation	UNP O60563
B	77	ARG	GLN	engineered mutation	UNP O60563
B	96	GLY	GLU	engineered mutation	UNP O60563
B	106	ARG	LYS	engineered mutation	UNP O60563
B	241	LEU	PHE	engineered mutation	UNP O60563

- Molecule 3 is (1S,3S)-N3-(5-pentan-3-yl)pyrazolo[1,5-a]pyrimidin-7-yl)cyclopentane-1,3-diamine (CCD ID: VQE) (formula: C₁₆H₂₅N₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			21	16	5		

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	171.50Å 171.50Å 96.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	85.75 – 3.75 85.75 – 3.75	Depositor EDS
% Data completeness (in resolution range)	99.7 (85.75-3.75) 99.7 (85.75-3.75)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 3.78Å)	Xtriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.200 , 0.245 0.201 , 0.243	Depositor DCC
R_{free} test set	537 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	186.6	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 190.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4558	wwPDB-VP
Average B, all atoms (Å ²)	207.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, VQE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/2516	0.85	1/3392 (0.0%)
2	B	0.43	0/2110	0.89	0/2875
All	All	0.44	0/4626	0.87	1/6267 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	208	PRO	N-CA-C	5.74	117.70	110.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2480	0	2516	56	0
2	B	2057	0	2041	46	0
3	A	21	0	0	0	0
All	All	4558	0	4557	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:219:MET:HE1	1:A:291:LEU:HD23	1.54	0.87
1:A:248:ILE:HG22	1:A:256:VAL:HG21	1.62	0.81
1:A:248:ILE:HG23	1:A:298:LEU:HD11	1.70	0.74
1:A:73:LEU:HD13	1:A:140:ILE:HG22	1.71	0.72
2:B:70:TYR:CD2	2:B:75:PHE:CZ	2.79	0.71
1:A:82:ILE:HD12	1:A:102:VAL:HG12	1.80	0.63
2:B:12:TRP:CD2	2:B:68:ARG:HD3	2.34	0.63
2:B:48:MET:HE3	2:B:88:LEU:HD12	1.79	0.63
1:A:212:LEU:HD13	1:A:305:ASP:HA	1.82	0.62
2:B:70:TYR:CE2	2:B:75:PHE:CE1	2.90	0.60
1:A:13:CYS:SG	1:A:86:ARG:NE	2.72	0.59
1:A:244:LEU:HG	1:A:275:VAL:HG22	1.85	0.59
2:B:12:TRP:CZ3	2:B:68:ARG:HB3	2.38	0.58
1:A:13:CYS:SG	1:A:86:ARG:CG	2.92	0.58
2:B:126:TYR:HA	2:B:129:GLN:HG2	1.86	0.58
2:B:66:MET:C	2:B:66:MET:SD	2.87	0.57
1:A:253:TRP:CE3	1:A:298:LEU:HD22	2.38	0.57
1:A:320:MET:HE3	1:A:321:PRO:HD2	1.86	0.57
2:B:41:ALA:HB3	2:B:66:MET:CE	2.34	0.57
2:B:101:LEU:HD11	2:B:133:LEU:HD13	1.87	0.57
2:B:41:ALA:HB3	2:B:66:MET:HE2	1.88	0.56
1:A:219:MET:CE	1:A:291:LEU:HD23	2.31	0.55
1:A:77:ASN:HD21	1:A:132:MET:HE2	1.70	0.55
2:B:84:ALA:HB3	2:B:85:PRO:HD3	1.89	0.54
1:A:239:ALA:HA	1:A:268:VAL:HG21	1.89	0.54
2:B:70:TYR:CD2	2:B:75:PHE:CE1	2.96	0.54
1:A:52:MET:HG2	1:A:53:GLU:N	2.22	0.54
2:B:109:HIS:CE1	2:B:118:LEU:HA	2.43	0.54
2:B:176:PHE:C	2:B:176:PHE:CD1	2.85	0.53
1:A:253:TRP:HB2	1:A:298:LEU:HD13	1.90	0.52
2:B:90:LEU:HD11	2:B:141:LEU:HD21	1.91	0.52
2:B:210:TRP:C	2:B:210:TRP:CD1	2.87	0.52
2:B:73:GLN:HB3	2:B:78:PHE:CE1	2.45	0.51
2:B:70:TYR:CG	2:B:75:PHE:CZ	3.00	0.50
2:B:159:LYS:O	2:B:163:LEU:HG	2.12	0.50
1:A:253:TRP:CE3	1:A:298:LEU:CD2	2.95	0.50
1:A:196:PRO:HB3	1:A:213:TRP:CD1	2.47	0.49
1:A:52:MET:HG2	1:A:53:GLU:H	1.77	0.49
1:A:52:MET:SD	1:A:54:ASN:HB2	2.53	0.49
2:B:66:MET:SD	2:B:67:HIS:N	2.85	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:CYS:SG	1:A:86:ARG:HG3	2.53	0.48
2:B:90:LEU:CD1	2:B:141:LEU:HD21	2.43	0.48
1:A:127:LYS:HG2	1:A:314:PHE:CZ	2.48	0.48
2:B:101:LEU:CD1	2:B:133:LEU:HD13	2.43	0.48
2:B:245:LEU:HB3	2:B:252:LEU:HD21	1.97	0.47
1:A:77:ASN:HA	1:A:163:LEU:O	2.13	0.47
1:A:163:LEU:C	1:A:163:LEU:HD23	2.40	0.47
1:A:208:PRO:HB2	1:A:209:PRO:HD3	1.97	0.47
2:B:65:TYR:CD2	2:B:90:LEU:HD23	2.50	0.47
2:B:83:VAL:HG13	2:B:140:ILE:HD11	1.97	0.47
2:B:164:VAL:O	2:B:165:ARG:C	2.57	0.47
1:A:112:GLY:O	1:A:116:ASN:HB2	2.15	0.46
2:B:160:CYS:O	2:B:164:VAL:HG22	2.15	0.46
1:A:55:GLU:HG3	1:A:60:PRO:HG3	1.96	0.46
1:A:181:GLN:HB3	1:A:182:PRO:HD2	1.96	0.46
1:A:70:LEU:HD11	1:A:168:PHE:HB2	1.98	0.46
1:A:318:ASP:N	1:A:318:ASP:OD1	2.49	0.46
2:B:121:THR:HA	2:B:126:TYR:CD1	2.52	0.45
1:A:145:ILE:HD13	1:A:173:ALA:HA	1.98	0.45
1:A:268:VAL:HB	1:A:271:GLN:HE21	1.82	0.45
1:A:187:ASN:HD21	1:A:202:GLY:HA2	1.81	0.45
2:B:73:GLN:HB3	2:B:78:PHE:CD1	2.51	0.44
1:A:189:VAL:C	1:A:190:VAL:HG23	2.42	0.44
2:B:220:HIS:CD2	2:B:222:TRP:CZ2	3.04	0.44
1:A:283:VAL:O	1:A:284:ARG:HB2	2.16	0.44
1:A:196:PRO:HD2	1:A:199:LEU:HD12	1.99	0.44
1:A:128:ARG:O	1:A:129:VAL:C	2.61	0.43
1:A:149:ASP:HB2	1:A:170:LEU:HD12	1.98	0.43
1:A:208:PRO:O	1:A:210:ILE:N	2.51	0.43
2:B:13:TYR:HA	2:B:189:LEU:O	2.18	0.43
2:B:67:HIS:O	2:B:189:LEU:HD11	2.17	0.43
2:B:132:ASP:O	2:B:135:ILE:HB	2.19	0.43
1:A:147:HIS:HB3	1:A:211:ASP:OD1	2.19	0.43
1:A:197:PRO:O	1:A:198:GLU:C	2.62	0.43
1:A:151:LYS:HE2	1:A:154:ASN:HD21	1.84	0.43
1:A:67:ILE:HG23	1:A:84:ILE:HD11	2.01	0.43
2:B:192:THR:O	2:B:196:VAL:HG23	2.19	0.43
1:A:77:ASN:ND2	1:A:132:MET:HE2	2.33	0.42
1:A:52:MET:SD	1:A:54:ASN:CB	3.07	0.42
1:A:108:HIS:CG	1:A:113:LEU:HD21	2.53	0.42
2:B:96:GLY:C	2:B:98:PRO:HD3	2.45	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:20:GLU:O	2:B:25:ARG:NE	2.52	0.42
1:A:280:ALA:HA	1:A:289:LEU:HD21	2.01	0.42
2:B:220:HIS:N	2:B:223:GLU:OE1	2.50	0.42
2:B:57:LEU:HD21	2:B:95:GLU:OE1	2.20	0.41
1:A:96:LYS:HB3	1:A:100:TYR:OH	2.19	0.41
1:A:187:ASN:HD21	1:A:202:GLY:CA	2.33	0.41
1:A:260:GLU:HB3	1:A:266:GLU:OE1	2.19	0.41
2:B:131:GLN:O	2:B:132:ASP:C	2.63	0.41
1:A:13:CYS:SG	1:A:86:ARG:HG2	2.61	0.41
2:B:12:TRP:CZ2	2:B:144:LEU:HA	2.55	0.41
2:B:212:ILE:HG23	2:B:221:TRP:CD1	2.55	0.41
2:B:41:ALA:HB3	2:B:66:MET:HE3	2.03	0.41
2:B:105:ILE:O	2:B:106:ARG:C	2.64	0.41
2:B:156:HIS:CE1	2:B:194:PRO:HG3	2.56	0.41
1:A:51:LEU:HD13	1:A:60:PRO:HD3	2.03	0.40
1:A:309:ALA:O	1:A:315:PHE:CE1	2.74	0.40
2:B:39:GLN:HE22	2:B:244:ILE:HD11	1.84	0.40
1:A:227:PRO:HB2	1:A:230:GLN:HE21	1.86	0.40
1:A:106:CYS:O	1:A:107:GLU:C	2.63	0.40
1:A:295:LEU:HD23	1:A:304:ILE:CG2	2.51	0.40
2:B:97:GLN:N	2:B:98:PRO:HD3	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/330 (89%)	255 (86%)	35 (12%)	5 (2%)	7	33
2	B	250/259 (96%)	232 (93%)	17 (7%)	1 (0%)	30	61
All	All	545/589 (92%)	487 (89%)	52 (10%)	6 (1%)	11	40

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	190	VAL
2	B	165	ARG
1	A	98	SER
1	A	181	GLN
1	A	107	GLU
1	A	209	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/290 (93%)	264 (97%)	7 (3%)	40	59
2	B	229/235 (97%)	223 (97%)	6 (3%)	40	59
All	All	500/525 (95%)	487 (97%)	13 (3%)	40	59

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	CYS
1	A	61	ILE
1	A	76	GLU
1	A	108	HIS
1	A	249	THR
1	A	298	LEU
1	A	318	ASP
2	B	30	ASP
2	B	64	VAL
2	B	66	MET
2	B	67	HIS
2	B	136	LEU
2	B	176	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	181	GLN
1	A	183	ASN
1	A	187	ASN
1	A	230	GLN
2	B	39	GLN
2	B	40	GLN
2	B	50	GLN
2	B	183	HIS
2	B	209	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	186	1	8,10,11	0.84	0	10,14,16	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	186	1	-	2/9/11/13	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	186	TPO	O-C-CA-CB
1	A	186	TPO	CG2-CB-OG1-P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	VQE	A	401	-	22,23,23	1.69	3 (13%)	20,32,32	1.84	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	VQE	A	401	-	-	1/7/21/21	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	VQE	C6-N3	6.35	1.39	1.32
3	A	401	VQE	C3-C6	2.62	1.51	1.49
3	A	401	VQE	C7-C6	-2.14	1.41	1.45

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	VQE	C11-N3-C6	-5.19	113.83	117.10
3	A	401	VQE	N1-C11-N3	5.07	125.17	121.49

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	VQE	C6-C3-C4-C5

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	305/330 (92%)	-0.56	0	100 100	168, 225, 265, 295	0
2	B	252/259 (97%)	-0.67	2 (0%)	82 62	133, 181, 236, 258	0
All	All	557/589 (94%)	-0.61	2 (0%)	88 73	133, 206, 260, 295	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	176	PHE	2.1
2	B	254	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	A	186	11/12	0.86	0.05	245,259,265,273	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	VQE	A	401	21/21	0.96	0.10	151,174,190,200	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.