



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:23 PM UTC

PDB ID : 3K65 / pdb_00003k65
Title : Crystal Structure of Prethombin-2/Fragment-2 Complex
Authors : Adams, T.E.; Huntington, J.A.
Deposited on : 2009-10-08
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	116	
2	B	308	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3172 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prothrombin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	80	Total	C	N	O	S	0	1	0
			618	386	108	118	6			


- Molecule 2 is a protein called Prothrombin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	274	Total	C	N	O	S	0	10	0
			2272	1465	395	398	14			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	525	ALA	SER	engineered mutation	UNP P00734

- Molecule 3 is 1,4-BUTANEDIOL (CCD ID: BU1) (formula: C₄H₁₀O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	4	2		
3	A	1	Total	C	O	0	0
			6	4	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	84	Total	O	0	0
			84	84		
4	B	186	Total	O	0	0
			186	186		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Prothrombin



• Molecule 2: Prothrombin



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	73.53Å 73.53Å 205.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.20 – 1.85 51.20 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.7 (51.20-1.85) 98.7 (51.20-1.85)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.20 (at 1.86Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.213 , 0.229 0.225 , 0.221	Depositor DCC
R_{free} test set	2467 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.587	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 56.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3172	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/634	0.79	0/863
2	B	0.35	0/2330	1.02	16/3151 (0.5%)
All	All	0.35	0/2964	0.97	16/4014 (0.4%)

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	552	ASP	N-CA-C	13.49	127.94	110.53
2	B	553	ARG	N-CA-C	10.59	133.35	110.80
2	B	554	ASP	N-CA-C	-9.78	91.60	110.56
2	B	410	TYR	N-CA-C	-7.48	98.59	109.59
2	B	307	LYS	N-CA-C	7.27	120.07	111.71
2	B	555	GLY	N-CA-C	-7.04	96.48	113.18
2	B	389	THR	N-CA-C	7.02	122.02	113.38
2	B	340	ARG	N-CA-C	-6.48	99.46	109.76
2	B	563	VAL	N-CA-C	6.35	116.52	110.42
2	B	556	LYS	N-CA-C	6.21	119.69	108.69
2	B	379	LEU	N-CA-C	5.81	118.25	109.41
2	B	529	PHE	N-CA-C	-5.76	97.94	108.02
2	B	368	PRO	N-CA-C	5.41	117.30	110.70
2	B	442	ASP	N-CA-C	-5.28	102.19	110.17
2	B	386	HIS	N-CA-C	-5.23	106.84	113.43
2	B	414	GLU	N-CA-C	5.19	119.08	111.34

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	618	0	556	4	0
2	B	2272	0	2217	53	0
3	A	12	0	20	2	0
4	A	84	0	0	1	0
4	B	186	0	0	7	0
All	All	3172	0	2793	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:512:PRO:HA	2:B:555:GLY:HA3	1.52	0.92
2:B:490[B]:ARG:NE	2:B:494:LYS:HE2	1.90	0.86
2:B:490[B]:ARG:HE	2:B:494:LYS:HE2	1.46	0.79
2:B:324:GLY:HA2	2:B:483[B]:VAL:HG22	1.66	0.76
2:B:551:CYS:HA	2:B:556:LYS:HE2	1.67	0.75
2:B:551:CYS:HA	2:B:556:LYS:CE	2.19	0.73
2:B:376:GLU:H	2:B:376:GLU:CD	1.98	0.71
2:B:388:ARG:HB2	2:B:461[A]:TRP:CE3	2.26	0.70
2:B:490[A]:ARG:HB3	2:B:491:PRO:HD3	1.76	0.68
3:A:600:BU1:H31	2:B:373:ASN:HD21	1.59	0.67
2:B:512:PRO:HA	2:B:555:GLY:CA	2.24	0.67
2:B:485:LEU:HD11	2:B:551:CYS:HB2	1.78	0.65
2:B:490[B]:ARG:NH2	2:B:503:ASP:OD1	2.31	0.64
1:A:178:TYR:CZ	1:A:180:GLY:HA3	2.34	0.62
2:B:388:ARG:HB2	2:B:461[B]:TRP:CD1	2.37	0.59
2:B:375:THR:HB	2:B:376:GLU:OE2	2.02	0.59
2:B:573:VAL:O	2:B:577:PHE:HD2	1.89	0.56
2:B:337:MET:HG2	2:B:461[B]:TRP:CZ3	2.40	0.56
2:B:322:VAL:HG13	2:B:322:VAL:O	2.06	0.54
2:B:301:LYS:HE3	4:B:854:HOH:O	2.06	0.54
2:B:380:LEU:C	2:B:380:LEU:HD12	2.33	0.53
2:B:403:LYS:HD3	2:B:405:TYR:OH	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:450[A]:LEU:HD12	2:B:487:ILE:HD13	1.91	0.53
2:B:393:ARG:HG2	2:B:394:ASN:ND2	2.23	0.53
2:B:551:CYS:HB3	4:B:813:HOH:O	2.09	0.52
2:B:490[B]:ARG:HE	2:B:494:LYS:CE	2.20	0.51
2:B:298:LEU:CD1	2:B:433:ASP:HB3	2.42	0.50
2:B:382:ARG:HD2	2:B:396:GLU:OE2	2.12	0.49
2:B:324:GLY:CA	2:B:483[B]:VAL:HG22	2.39	0.49
2:B:499:ILE:HD12	2:B:547:TRP:CZ3	2.48	0.49
1:A:173:ASP:O	1:A:176:GLN:HG3	2.13	0.49
2:B:461[B]:TRP:HZ3	4:B:651:HOH:O	1.95	0.49
2:B:490[B]:ARG:HG2	2:B:494:LYS:HD3	1.95	0.48
2:B:524:ASP:HB3	4:B:622:HOH:O	2.13	0.48
1:A:234:ALA:HB3	1:A:239:ASP:HB3	1.96	0.47
2:B:450[B]:LEU:HD12	4:B:859:HOH:O	2.14	0.47
2:B:577:PHE:CD2	2:B:577:PHE:N	2.83	0.47
2:B:357:TRP:CE3	2:B:422[A]:LEU:HG	2.50	0.47
2:B:365:LEU:HD13	2:B:404:ILE:HG21	1.97	0.46
2:B:459:THR:HG22	2:B:482[A]:VAL:HG22	1.98	0.46
2:B:340:ARG:HD2	2:B:343:PRO:O	2.16	0.46
2:B:324:GLY:HA2	2:B:482[A]:VAL:O	2.17	0.44
2:B:415:ASN:O	2:B:416:LEU:HB2	2.18	0.44
2:B:551:CYS:C	2:B:556:LYS:HD3	2.43	0.44
2:B:455:LYS:NZ	4:B:772:HOH:O	2.52	0.43
2:B:456:GLY:HA3	2:B:529:PHE:CZ	2.54	0.43
2:B:298:LEU:HD11	2:B:433:ASP:HB3	2.01	0.43
2:B:306:ASP:HB2	2:B:331:MET:HE3	2.02	0.42
2:B:558:GLY:HA3	4:B:813:HOH:O	2.18	0.42
1:A:192:LEU:HD13	1:A:233:VAL:HA	2.02	0.42
2:B:403:LYS:HD3	2:B:405:TYR:CZ	2.54	0.42
2:B:485:LEU:HD11	2:B:551:CYS:CB	2.49	0.42
2:B:382:ARG:HG2	2:B:398:ILE:HG12	2.02	0.41
2:B:280:PHE:HE1	2:B:571:GLN:HB2	1.85	0.41
2:B:324:GLY:HA2	2:B:483[B]:VAL:CG2	2.44	0.41
2:B:385:LYS:HE3	2:B:387:SER:O	2.22	0.40
3:A:599:BU1:H11	4:A:738:HOH:O	2.21	0.40
2:B:450[A]:LEU:HD12	2:B:487:ILE:CD1	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	79/116 (68%)	74 (94%)	4 (5%)	1 (1%)	9	2
2	B	275/308 (89%)	262 (95%)	12 (4%)	1 (0%)	30	17
All	All	354/424 (84%)	336 (95%)	16 (4%)	2 (1%)	21	10

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	180	GLY
2	B	553	ARG

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	63/94 (67%)	62 (98%)	1 (2%)	55	44
2	B	239/267 (90%)	233 (98%)	6 (2%)	42	27
All	All	302/361 (84%)	295 (98%)	7 (2%)	43	30

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	215	VAL
2	B	298	LEU
2	B	312	LEU
2	B	376	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	410	TYR
2	B	507	CYS
2	B	577	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	177	GLN
1	A	198	GLN
2	B	373	ASN
2	B	394	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BU1	A	600	-	5,5,5	0.36	0	4,4,4	0.42	0
3	BU1	A	599	-	5,5,5	0.36	0	4,4,4	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BU1	A	600	-	-	0/3/3/3	-
3	BU1	A	599	-	-	0/3/3/3	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	600	BU1	1	0
3	A	599	BU1	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	80/116 (68%)	1.48	22 (27%) 1 1	25, 39, 59, 62	1 (1%)
2	B	274/308 (88%)	0.92	40 (14%) 6 5	13, 34, 63, 83	10 (3%)
All	All	354/424 (83%)	1.04	62 (17%) 4 3	13, 35, 60, 83	11 (3%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	555	GLY	6.9
2	B	551	CYS	6.5
2	B	577	PHE	6.2
2	B	552	ASP	5.1
2	B	576	GLN	4.9
1	A	173	ASP	4.8
2	B	317	ILE	4.7
1	A	215	VAL	4.6
1	A	248	CYS	4.5
2	B	513	ASP	4.4
2	B	554	ASP	4.3
2	B	324	GLY	4.2
2	B	316	TYR	4.1
1	A	218	PHE	4.0
2	B	318	ASP	4.0
1	A	170	CYS	3.9
2	B	553	ARG	3.9
2	B	389	THR	3.8
1	A	245[A]	LEU	3.8
1	A	169	GLN	3.7
2	B	279	THR	3.7
2	B	314	GLU	3.5
2	B	321	ILE	3.5
2	B	319	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	523	GLY	3.4
2	B	575	ASP	3.3
1	A	211	ALA	3.3
2	B	524	ASP	3.2
2	B	322	VAL	3.2
2	B	490[A]	ARG	3.1
2	B	315[A]	SER	3.1
2	B	476	GLN	3.0
2	B	394	ASN	3.0
2	B	493	CYS	2.9
2	B	377	ASN	2.8
1	A	212	VAL	2.8
1	A	183	ALA	2.7
2	B	512	PRO	2.7
1	A	179	GLN	2.7
2	B	494	LYS	2.7
1	A	210	SER	2.6
2	B	280	PHE	2.6
2	B	448	SER	2.6
2	B	442	ASP	2.6
1	A	178	TYR	2.5
2	B	454	TYR	2.4
2	B	311	GLU	2.4
1	A	246	ASN	2.4
1	A	209	ASN	2.4
1	A	172	PRO	2.4
1	A	182	LEU	2.3
2	B	393	ARG	2.3
2	B	398	ILE	2.3
1	A	244	ASP	2.2
1	A	184	VAL	2.2
2	B	451	GLN	2.2
2	B	302[A]	LYS	2.2
2	B	382	ARG	2.1
1	A	180	GLY	2.1
1	A	189	LEU	2.1
1	A	171	VAL	2.1
2	B	477	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BU1	A	600	6/6	0.60	0.24	71,72,73,74	0
3	BU1	A	599	6/6	0.77	0.20	64,66,67,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.