



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 06:59 PM EDT

PDB ID : 3K6J / pdb_00003k6j
Title : Crystal structure of the dehydrogenase part of multifunctional enzyme 1 from *C.elegans*
Authors : Ouyang, Z.; Zhang, K.; Zhai, Y.; Lu, J.; Sun, F.
Deposited on : 2009-10-09
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3633 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein F01G10.3, confirmed by transcript evidence.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	1	0
			3436	2179	588	645	24			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	MET	-	expression tag	UNP O17762
A	-13	GLY	-	expression tag	UNP O17762
A	-12	HIS	-	expression tag	UNP O17762
A	-11	HIS	-	expression tag	UNP O17762
A	-10	HIS	-	expression tag	UNP O17762
A	-9	HIS	-	expression tag	UNP O17762
A	-8	HIS	-	expression tag	UNP O17762
A	-7	HIS	-	expression tag	UNP O17762
A	-6	HIS	-	expression tag	UNP O17762
A	-5	HIS	-	expression tag	UNP O17762
A	-4	SER	-	expression tag	UNP O17762
A	-3	THR	-	expression tag	UNP O17762
A	-2	GLY	-	expression tag	UNP O17762
A	-1	GLU	-	expression tag	UNP O17762
A	0	ASN	-	expression tag	UNP O17762
A	1	LEU	-	expression tag	UNP O17762
A	2	TYR	-	expression tag	UNP O17762
A	3	PHE	-	expression tag	UNP O17762
A	4	GLN	-	expression tag	UNP O17762
A	5	GLY	-	expression tag	UNP O17762
A	6	SER	-	expression tag	UNP O17762
A	427	VAL	-	expression tag	UNP O17762
A	428	SER	-	expression tag	UNP O17762
A	429	THR	-	expression tag	UNP O17762
A	430	GLY	-	expression tag	UNP O17762
A	431	SER	-	expression tag	UNP O17762
A	432	SER	-	expression tag	UNP O17762

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	433	GLY	-	expression tag	UNP O17762
A	434	SER	-	expression tag	UNP O17762
A	435	SER	-	expression tag	UNP O17762
A	436	GLY	-	expression tag	UNP O17762
A	437	GLY	-	expression tag	UNP O17762
A	438	HIS	-	expression tag	UNP O17762
A	439	HIS	-	expression tag	UNP O17762
A	440	HIS	-	expression tag	UNP O17762
A	441	HIS	-	expression tag	UNP O17762
A	442	HIS	-	expression tag	UNP O17762
A	443	HIS	-	expression tag	UNP O17762
A	444	HIS	-	expression tag	UNP O17762
A	445	HIS	-	expression tag	UNP O17762

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Zn 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	176	Total 176	O 176	0	0

- Molecule 1: Protein F01G10.3, confirmed by transcript evidence



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.03Å 90.66Å 66.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.09 – 2.20 31.09 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (31.09-2.20) 99.9 (31.09-2.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.91 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.210 , 0.267 0.213 , 0.221	Depositor DCC
R_{free} test set	1248 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.428	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3633	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	1/3505 (0.0%)	0.89	1/4719 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	50	MET	C-O	-5.92	1.17	1.24

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	164	ARG	N-CA-C	-6.22	104.19	110.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3436	0	3419	64	0
2	A	20	0	0	1	0
3	A	1	0	0	0	0
4	A	176	0	0	2	0
All	All	3633	0	3419	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:ILE:HG22	1:A:290:MET:HE2	1.37	1.05
1:A:281:LEU:CB	1:A:282:GLU:HA	1.96	0.95
1:A:281:LEU:HB2	1:A:282:GLU:HA	1.51	0.92
1:A:353:ASN:HD21	1:A:419:ALA:H	1.24	0.84
1:A:155:ASP:HB3	1:A:158:GLU:HG2	1.64	0.80
1:A:127:MET:CE	1:A:131:LYS:HG3	2.15	0.76
1:A:331:ALA:O	1:A:332:LYS:HB2	1.87	0.75
1:A:127:MET:HE2	1:A:131:LYS:HE3	1.69	0.75
1:A:281:LEU:HB3	1:A:282:GLU:HA	1.74	0.69
1:A:281:LEU:CB	1:A:282:GLU:CA	2.74	0.65
1:A:34:SER:OG	1:A:194:SER:HB2	1.97	0.65
1:A:344:ILE:HG23	1:A:348:LEU:HD22	1.82	0.62
1:A:132:GLU:O	1:A:136:ASN:HB2	2.01	0.61
1:A:127:MET:HE3	1:A:131:LYS:HG3	1.81	0.60
1:A:127:MET:HE1	1:A:159:ILE:HG12	1.85	0.59
1:A:160:SER:O	1:A:162:VAL:N	2.36	0.58
1:A:284:ASN:HD22	1:A:286:ILE:H	1.50	0.58
1:A:76:LYS:HG2	1:A:106:ILE:HD12	1.85	0.58
1:A:281:LEU:CB	1:A:283:PRO:HD3	2.35	0.56
1:A:160:SER:C	1:A:162:VAL:H	2.14	0.56
1:A:247:ILE:HG21	1:A:261:MET:HE2	1.87	0.56
1:A:178:ALA:O	1:A:209:LYS:NZ	2.39	0.55
1:A:281:LEU:HB3	1:A:282:GLU:CA	2.36	0.55
1:A:281:LEU:HB3	1:A:283:PRO:HD3	1.89	0.55
1:A:91:ARG:NH2	1:A:266:MET:HE1	2.22	0.54
1:A:269:PHE:CD1	1:A:272:MET:HG3	2.42	0.54
1:A:286:ILE:HG22	1:A:290:MET:CE	2.26	0.54
1:A:331:ALA:O	1:A:332:LYS:CB	2.57	0.52
1:A:155:ASP:HB3	1:A:158:GLU:CG	2.36	0.52
1:A:127:MET:HE2	1:A:131:LYS:HG3	1.91	0.51
1:A:164:ARG:O	1:A:166:PRO:HD3	2.11	0.51
1:A:14:GLU:OE2	1:A:182:ARG:NH1	2.44	0.51
1:A:306:LYS:HD3	4:A:663:HOH:O	2.11	0.50
1:A:402:MET:CE	1:A:402:MET:HA	2.43	0.49
1:A:320:MET:O	1:A:324:ILE:HG12	2.12	0.49
1:A:173:HIS:HB2	1:A:185:GLU:HB2	1.96	0.47
1:A:127:MET:HE2	1:A:131:LYS:CE	2.43	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:339:ASN:HD22	1:A:341:GLN:H	1.62	0.46
1:A:280:GLY:HA2	1:A:281:LEU:HA	1.67	0.46
1:A:269:PHE:HE2	1:A:290:MET:HE3	1.81	0.45
1:A:91:ARG:HH21	1:A:266:MET:HE1	1.81	0.45
1:A:155:ASP:CB	1:A:158:GLU:HG2	2.42	0.45
1:A:283:PRO:HB3	1:A:291:TRP:CZ3	2.52	0.45
1:A:150:ASN:ND2	1:A:150:ASN:C	2.75	0.44
1:A:209:LYS:HD3	2:A:500:PO4:O2	2.18	0.44
1:A:290:MET:HE1	1:A:304:PHE:CZ	2.52	0.44
1:A:56:ILE:HG23	1:A:100:ILE:HD13	1.99	0.43
1:A:359:ILE:HD11	1:A:369:LEU:HD23	2.00	0.43
1:A:139:ASN:HB3	1:A:164:ARG:HH22	1.82	0.43
1:A:110:PHE:C	1:A:112:LYS:H	2.25	0.43
1:A:272:MET:HE3	1:A:272:MET:HB3	1.95	0.43
1:A:281:LEU:HG	1:A:283:PRO:HD3	2.02	0.42
1:A:121:GLU:OE2	1:A:123:VAL:HG22	2.19	0.42
1:A:160:SER:C	1:A:162:VAL:N	2.77	0.42
1:A:343:VAL:HG12	1:A:347:MET:HE2	2.02	0.41
1:A:420:ASP:HA	1:A:423:LYS:HD3	2.02	0.41
1:A:-2:GLY:HA2	1:A:176:ASN:O	2.20	0.41
1:A:247:ILE:HD13	1:A:261:MET:HE3	2.03	0.41
1:A:137:LEU:HA	1:A:140:ILE:HG22	2.03	0.41
1:A:168:ASN:HB3	4:A:618:HOH:O	2.20	0.41
1:A:122:SER:HA	1:A:150:ASN:ND2	2.36	0.41
1:A:166:PRO:HG2	1:A:191[A]:HIS:CD2	2.55	0.41
1:A:139:ASN:OD1	1:A:164:ARG:NH2	2.54	0.40
1:A:-5:HIS:HB3	1:A:175:PHE:CD1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	427/460 (93%)	409 (96%)	14 (3%)	4 (1%)	14 14

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	310	LYS
1	A	332	LYS
1	A	161	SER
1	A	333	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	380/404 (94%)	363 (96%)	17 (4%)	24 33

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	ARG
1	A	139	ASN
1	A	150	ASN
1	A	244	PRO
1	A	277	LYS
1	A	278	GLU
1	A	284	ASN
1	A	288	LYS
1	A	308	ASP
1	A	309	ASP
1	A	327	VAL
1	A	332	LYS
1	A	348	LEU
1	A	357	ARG
1	A	382	ILE
1	A	402	MET
1	A	417	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	0	ASN
1	A	73	GLN
1	A	150	ASN
1	A	176	ASN
1	A	179	ASN
1	A	234	GLN
1	A	284	ASN
1	A	301	ASN
1	A	312	GLN
1	A	316	ASN
1	A	339	ASN
1	A	341	GLN
1	A	353	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	500	-	4,4,4	0.97	0	6,6,6	0.49	0
2	PO4	A	501	-	4,4,4	0.94	0	6,6,6	0.83	0
2	PO4	A	502	-	4,4,4	0.93	0	6,6,6	0.48	0
2	PO4	A	503	-	4,4,4	0.95	0	6,6,6	0.54	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	430/460 (93%)	0.17	15 (3%)	47 44	13, 34, 68, 81	1 (0%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	327	VAL	4.4
1	A	331	ALA	4.3
1	A	281	LEU	3.9
1	A	280	GLY	3.8
1	A	311	THR	3.6
1	A	324	ILE	3.5
1	A	286	ILE	2.8
1	A	241	GLY	2.7
1	A	162	VAL	2.6
1	A	285	PRO	2.5
1	A	139	ASN	2.3
1	A	325	ARG	2.3
1	A	318	THR	2.1
1	A	334	ASN	2.0
1	A	288	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	A	500	5/5	0.79	0.16	98,98,99,99	0
2	PO4	A	502	5/5	0.79	0.20	107,107,107,107	0
2	PO4	A	503	5/5	0.80	0.12	86,86,87,87	0
2	PO4	A	501	5/5	0.92	0.10	48,51,52,52	0
3	ZN	A	800	1/1	1.00	0.02	19,19,19,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.