



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:53 AM UTC

PDB ID : 4K6J / pdb_00004k6j
Title : Human cohesin inhibitor WapL
Authors : Tomchick, D.R.; Yu, H.; Ouyang, Z.
Deposited on : 2013-04-16
Resolution : 2.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	568	7% 78% 9% 12%
1	B	568	9% 75% 7% 17%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 15557 atoms, of which 7730 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Wings apart-like protein homolog.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	497	Total	C	H	N	O	S	0	0	0
			7887	2451	3963	689	752	32			
1	B	471	Total	C	H	N	O	S	0	0	0
			7485	2330	3761	654	709	31			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLY	-	expression tag	UNP Q7Z5K2
A	-6	PRO	-	expression tag	UNP Q7Z5K2
A	-5	LEU	-	expression tag	UNP Q7Z5K2
A	-4	GLY	-	expression tag	UNP Q7Z5K2
A	-3	SER	-	expression tag	UNP Q7Z5K2
A	-2	GLY	-	expression tag	UNP Q7Z5K2
A	-1	ARG	-	expression tag	UNP Q7Z5K2
A	0	PRO	-	expression tag	UNP Q7Z5K2
B	-7	GLY	-	expression tag	UNP Q7Z5K2
B	-6	PRO	-	expression tag	UNP Q7Z5K2
B	-5	LEU	-	expression tag	UNP Q7Z5K2
B	-4	GLY	-	expression tag	UNP Q7Z5K2
B	-3	SER	-	expression tag	UNP Q7Z5K2
B	-2	GLY	-	expression tag	UNP Q7Z5K2
B	-1	ARG	-	expression tag	UNP Q7Z5K2
B	0	PRO	-	expression tag	UNP Q7Z5K2

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			7	2	3	2		
3	A	1	Total	C	H	O	0	0
			7	2	3	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	58	Total	O	0	0
			58	58		
4	B	48	Total	O	0	0
			48	48		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Wings apart-like protein homolog



• Molecule 1: Wings apart-like protein homolog



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	107.54Å 107.54Å 300.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.68 – 2.62 29.68 – 2.62	Depositor EDS
% Data completeness (in resolution range)	94.8 (29.68-2.62) 94.7 (29.68-2.62)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.42 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.184 , 0.237 (Not available) , 0.233	Depositor DCC
R_{free} test set	2587 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.261	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 57.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	15557	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3979	0.66	0/5369
1	B	0.28	0/3776	0.67	0/5089
All	All	0.29	0/7755	0.67	0/10458

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3924	3963	3943	31	0
1	B	3724	3761	3742	29	0
2	A	35	0	0	0	0
2	B	30	0	0	1	0
3	A	8	6	6	0	0
4	A	58	0	0	0	0
4	B	48	0	0	1	0
All	All	7827	7730	7691	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:715:ASN:ND2	1:B:765:MET:SD	2.65	0.70
1:A:738:ARG:HG2	1:A:818:LEU:HD13	1.81	0.63
1:B:853:ARG:NH2	4:B:1309:HOH:O	2.31	0.63
1:B:-1:ARG:N	1:B:0:PRO:HD2	2.15	0.61
1:B:992:VAL:O	1:B:999:ARG:NH1	2.34	0.61
1:B:644:ASP:O	1:B:646:VAL:N	2.34	0.60
1:A:946:TRP:CD1	1:B:817:LEU:HD13	2.39	0.58
1:A:1051:GLU:HA	1:A:1054:THR:HG22	1.87	0.57
1:A:638:HIS:NE2	1:A:644:ASP:OD2	2.40	0.55
1:A:738:ARG:HG2	1:A:818:LEU:CD1	2.37	0.55
1:A:-1:ARG:N	1:A:0:PRO:CD	2.72	0.52
1:B:1006:GLU:HA	1:B:1031:HIS:HB3	1.92	0.52
1:A:646:VAL:CG1	1:A:647:GLU:N	2.73	0.51
1:A:1030:VAL:HG22	1:A:1031:HIS:N	2.27	0.50
1:B:951:THR:HG22	1:B:957:LEU:HD22	1.93	0.50
1:B:1050:ALA:HB3	1:B:1115:HIS:HB2	1.92	0.50
1:B:1166:MET:HA	1:B:1166:MET:HE2	1.93	0.50
1:A:646:VAL:HG11	1:A:692:MET:HE1	1.95	0.49
1:A:738:ARG:HD3	1:A:818:LEU:HD13	1.95	0.49
1:B:657:ASP:HA	1:B:660:TYR:CE2	2.49	0.48
1:B:968:VAL:N	1:B:969:PRO:CD	2.77	0.47
1:A:968:VAL:N	1:A:969:PRO:CD	2.78	0.47
1:B:765:MET:O	1:B:768:ILE:N	2.48	0.47
1:B:981:ARG:NH2	2:B:1201:SO4:O3	2.48	0.47
1:A:951:THR:HG22	1:A:957:LEU:HD22	1.96	0.46
1:A:891:CYS:HB3	1:A:930:MET:HG3	1.97	0.46
1:B:669:GLN:HG3	1:B:670:PRO:HD2	1.97	0.46
1:B:645:VAL:O	1:B:645:VAL:HG13	2.15	0.46
1:A:669:GLN:HG3	1:A:670:PRO:HD2	1.97	0.45
1:A:1166:MET:HE2	1:A:1166:MET:HA	1.97	0.45
1:A:913:LEU:HB3	1:A:914:PRO:HD2	1.98	0.45
1:A:967:GLN:O	1:A:970:LYS:HG2	2.17	0.45
1:B:1175:THR:HG23	1:B:1176:GLY:N	2.31	0.44
1:A:642:PHE:O	1:A:643:ASN:HB2	2.17	0.44
1:A:900:ARG:HD2	1:A:923:GLY:HA3	1.99	0.44
1:A:1059:LYS:O	1:A:1060:ASP:C	2.59	0.44
1:B:660:TYR:C	1:B:660:TYR:CD1	2.97	0.43
1:B:657:ASP:HA	1:B:660:TYR:CD2	2.54	0.43
1:B:1172:VAL:HG13	1:B:1177:GLN:HG3	2.00	0.43
1:B:1041:LEU:O	1:B:1045:ARG:HB2	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:633:TYR:CE2	1:A:637:GLN:HG3	2.54	0.43
1:A:646:VAL:HG12	1:A:647:GLU:N	2.33	0.42
1:A:651:ASN:O	1:A:654:PHE:HB3	2.19	0.42
1:B:824:ILE:CG2	1:B:854:CYS:HB3	2.49	0.42
1:B:891:CYS:HB3	1:B:930:MET:HG3	2.01	0.42
1:B:714:GLN:HG2	1:B:715:ASN:N	2.35	0.42
1:B:699:MET:O	1:B:703:VAL:HG13	2.20	0.42
1:B:644:ASP:HB3	1:B:646:VAL:HG23	2.02	0.42
1:B:1169:THR:HG22	1:B:1171:ALA:HB2	2.01	0.42
1:B:1172:VAL:HG13	1:B:1177:GLN:CG	2.50	0.42
1:A:646:VAL:HG11	1:A:692:MET:CE	2.50	0.41
1:A:658:ILE:HD11	1:A:694:LEU:HD13	2.01	0.41
1:A:912:PRO:HB2	1:A:919:THR:OG1	2.21	0.41
1:A:1053:LYS:O	1:A:1057:LEU:HD13	2.20	0.41
1:A:1057:LEU:HD21	1:A:1107:LYS:HB3	2.02	0.41
1:A:946:TRP:CD1	1:B:817:LEU:CD1	3.03	0.41
1:A:1151:ASP:CG	1:A:1153:SER:HG	2.28	0.40
1:A:1030:VAL:HG21	1:A:1034:GLN:CD	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/568 (46%)	252 (97%)	9 (3%)	0	100	100
1	B	461/568 (81%)	444 (96%)	15 (3%)	2 (0%)	30	49
All	All	722/1136 (64%)	696 (96%)	24 (3%)	2 (0%)	36	56

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	645	VAL
1	B	641	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	443/504 (88%)	436 (98%)	7 (2%)	55	78
1	B	418/504 (83%)	412 (99%)	6 (1%)	59	80
All	All	861/1008 (85%)	848 (98%)	13 (2%)	57	79

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	653	GLU
1	A	793	HIS
1	A	878	SER
1	A	902	GLU
1	A	957	LEU
1	A	992	VAL
1	A	1120	ILE
1	B	703	VAL
1	B	771	LYS
1	B	957	LEU
1	B	1045	ARG
1	B	1115	HIS
1	B	1185	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	637	GLN
1	A	915	HIS
1	A	967	GLN
1	A	976	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	712	HIS
1	B	879	GLN
1	B	890	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	1201	-	4,4,4	0.25	0	6,6,6	0.08	0
2	SO4	A	1204	-	4,4,4	0.26	0	6,6,6	0.07	0
2	SO4	B	1202	-	4,4,4	0.21	0	6,6,6	0.17	0
2	SO4	B	1205	-	4,4,4	0.25	0	6,6,6	0.09	0
2	SO4	A	1202	-	4,4,4	0.23	0	6,6,6	0.10	0
2	SO4	A	1205	-	4,4,4	0.27	0	6,6,6	0.11	0
3	ACT	A	1209	-	3,3,3	0.83	0	3,3,3	1.26	0
2	SO4	B	1203	-	4,4,4	0.23	0	6,6,6	0.08	0
2	SO4	A	1206	-	4,4,4	0.25	0	6,6,6	0.09	0
2	SO4	A	1201	-	4,4,4	0.24	0	6,6,6	0.12	0
2	SO4	B	1204	-	4,4,4	0.22	0	6,6,6	0.20	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	1206	-	4,4,4	0.26	0	6,6,6	0.10	0
3	ACT	A	1208	-	3,3,3	0.83	0	3,3,3	1.30	0
2	SO4	A	1207	-	4,4,4	0.26	0	6,6,6	0.07	0
2	SO4	A	1203	-	4,4,4	0.23	0	6,6,6	0.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1201	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	497/568 (87%)	-0.01	39 (7%) 19 15	17, 41, 110, 160	0
1	B	471/568 (82%)	0.13	49 (10%) 11 9	17, 47, 113, 161	0
All	All	968/1136 (85%)	0.06	88 (9%) 15 11	17, 44, 111, 161	0

All (88) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	645	VAL	8.8
1	B	642	PHE	8.0
1	A	907	LEU	7.8
1	B	641	HIS	7.4
1	A	1009	CYS	6.3
1	A	644	ASP	6.3
1	A	1058	ILE	5.8
1	A	760	LEU	5.2
1	B	920	ASN	5.2
1	A	906	CYS	4.8
1	B	1109	LEU	4.7
1	B	644	ASP	4.5
1	A	643	ASN	4.5
1	B	655	THR	4.4
1	A	1031	HIS	4.1
1	A	1060	ASP	4.1
1	B	1049	LEU	4.0
1	B	652	GLN	3.9
1	A	641	HIS	3.9
1	A	1008	SER	3.9
1	A	1059	LYS	3.9
1	B	1108	ALA	3.9
1	A	1057	LEU	3.7
1	A	642	PHE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	763	LYS	3.6
1	B	1010	SER	3.4
1	A	1055	ASP	3.4
1	B	653	GLU	3.4
1	B	1009	CYS	3.4
1	A	909	ASP	3.4
1	A	1107	LYS	3.4
1	B	1107	LYS	3.4
1	B	651	ASN	3.4
1	A	838	ASP	3.3
1	A	654	PHE	3.3
1	A	1106	ASN	3.1
1	B	1051	GLU	3.1
1	B	752	GLN	3.1
1	A	837	ARG	3.1
1	B	767	LYS	3.1
1	B	1112	ALA	3.1
1	A	645	VAL	3.1
1	B	714	GLN	3.0
1	A	1109	LEU	2.9
1	B	783	HIS	2.9
1	B	764	ASP	2.9
1	B	654	PHE	2.9
1	B	1048	GLN	2.8
1	B	1052	SER	2.8
1	B	1045	ARG	2.7
1	B	1169	THR	2.7
1	B	1054	THR	2.7
1	A	905	ILE	2.6
1	B	638	HIS	2.6
1	B	649	GLY	2.6
1	A	904	SER	2.6
1	A	763	LYS	2.6
1	B	1113	GLY	2.5
1	A	836	SER	2.5
1	A	840	ASP	2.5
1	B	1008	SER	2.4
1	A	1056	GLU	2.4
1	B	657	ASP	2.4
1	B	838	ASP	2.4
1	B	1053	LYS	2.4
1	B	762	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	651	ASN	2.3
1	B	666	LYS	2.3
1	A	910	SER	2.3
1	B	1007	THR	2.3
1	A	914	PRO	2.3
1	B	1171	ALA	2.3
1	B	640	LYS	2.2
1	B	650	GLU	2.2
1	A	1049	LEU	2.2
1	B	836	SER	2.2
1	B	1050	ALA	2.2
1	A	1054	THR	2.2
1	A	1007	THR	2.1
1	A	1048	GLN	2.1
1	B	1172	VAL	2.1
1	B	656	ASP	2.1
1	B	1011	PHE	2.1
1	B	647	GLU	2.1
1	A	655	THR	2.1
1	A	752	GLN	2.0
1	A	786	LEU	2.0
1	B	1111	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ACT	A	1208	4/4	0.77	0.22	34,40,53,63	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ACT	A	1209	4/4	0.83	0.18	25,30,50,55	0
2	SO4	A	1204	5/5	0.85	0.15	66,79,92,94	0
2	SO4	A	1205	5/5	0.90	0.12	63,66,78,86	0
2	SO4	B	1206	5/5	0.95	0.10	45,59,64,71	0
2	SO4	B	1205	5/5	0.96	0.08	47,53,57,73	0
2	SO4	A	1203	5/5	0.96	0.07	46,50,59,66	0
2	SO4	A	1206	5/5	0.96	0.08	50,57,59,59	0
2	SO4	B	1201	5/5	0.96	0.09	34,40,63,69	0
2	SO4	A	1207	5/5	0.97	0.10	49,53,68,69	0
2	SO4	A	1202	5/5	0.97	0.09	39,47,54,66	0
2	SO4	B	1202	5/5	0.97	0.08	27,35,39,49	0
2	SO4	B	1204	5/5	0.97	0.08	34,45,64,75	0
2	SO4	B	1203	5/5	0.98	0.06	42,53,54,56	0
2	SO4	A	1201	5/5	0.98	0.06	34,38,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.